

6. References

- Akita M., S. Sasaki, S. Matsuyama, S. Mizushima.** 1990. SecA interacts with secretory proteins by recognizing the positive charge at the amino terminus of the signal peptide in *Escherichia coli*. *J. Biol. Chem.* **265**:8164-8169.
- Anagnostopoulos, C. & J. Spizizen.** 1961. Requirements for transformation in *Bacillus subtilis*. *J. Bacteriol.* **81**:741-746.
- Archibald, A. R., Hancock, I. C. and Harwood, C. R.** 1993. Cell wall structure, synthesis, and turnover. In *Bacillus subtilis and other Gram-positive Bacteria*, pp. 381-410. Edited by **J. A. Hoch and R. Losick**. Washington, DC: American Society for Microbiology.
- Ash, C., J.A. Farrow, M. Dorsch, E. Stackebrandt and M.D. Collins.** 1991. Comparative analysis of *Bacillus anthracis*, *Bacillus cereus*, and related species on the basis of reverse transcriptase sequencing of 16S rRNA. *Int. J. Syst. Bacteriol.* **41**:343-346.
- Bairl, A., & P. Müller.** 1998. A second gene for type I signal peptidase in *Bradyrhizobium japonicum*, *sipF*, is located near genes involved in RNA processing and cell division. *Mol. Gen. Genet.* **260**:346-356.
- Banerjee, S., and J. N. Hansen.** 1988. Structure and expression of a gene encoding the precursor of subtilin, a small protein antibiotic. *J. Biol. Chem.* **263**:9508-9514.
- Beissinger, M., and J. Buchner.** 1998. How chaperones fold proteins. *J. Biol. Chem.* **379**:245-259.
- Bersk, B. C.** 1996. A common export pathway for proteins binding complex redox cofactors? *Mol. Microbiol.* **22**:393-404.
- Berks, B. C., F. Sargent, and T. Palmer.** 2000. The Tat protein export pathway. *Mol. Microbiol.* **35**:260-274.
- Birnboim, H.C., J. Doly.** 1979. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acid Res.* **7**:1513-1523.
- Blaak, H. & H. Schrempf.** 1995. Binding and substrate specificities of a *Streptomyces olivaceoviridis* chitinase in comparison with its proteolytically processed form. *Eur J Biochem* **229**:132-139.
- Blackman, S. A., Smith, T. J. and Foster, S. J.** 1998. The role of autolysins during vegetative growth of *Bacillus subtilis* 168. *Microbiology* **144**:73-82.
- Blattner F.R., G. 3rd. Plunkett, C.A. Bloch, N.T. Perna, V. Burland, M. Riley, et al.** 1997. The complete genome sequence of *Escherichia coli* K-12. *Science.* **277**:1453-1474.
- Bohni P.C., R.J. Deshaies, R.W. Schekman.** 1988. SEC11 is required for signal peptide processing and yeast cell growth. *J. Cell Biol.* **106**:1035-1042.
- Boland, F. M., A. Atrih, H. Chirakkal, S. J. Foster, and A. Moir.** 2000. Complete spore-cortex hydrolysis during germination of *Bacillus subtilis* 168 requires SleB and YpeB. *Microbiology* **144**:73-82.
- Bolhuis, A., A. Sorokin, V. Azevedo, S. D. Ehrlich, P. G. Braun, A. de Jong, G. Venema, S. Bron, and J. M. van Dijk.** 1996. *Bacillus subtilis* can modulate its capacity and specificity for protein

- secretion by temporary controlled expression of the *sipS* gene for signal peptidase J. Mol. Microbiol. **22**:605-618.
- Bolhuis, A., C. P. Broekhuizen, A. Sorokin, M. L. van Roosmalen, G. Venema, S. Bron, W. J. Quax, and I. M. van Dijl.** 1998. SecDF of *Bacillus subtilis*, a molecular Siamese twin required for the efficient secretion of proteins. J. Biol. Chem. **273**:21217-21224.
- Bolhuis, A., G. Venema, W. J. Quax, S. Bron, R., and j. M. van Dijl.** 1999. Functional analysis of paralogous thiol-disulfide oxidoreductases in *Bacillus subtilis*. J. Biol. Chem. **274**:24531-24538.
- Bormann, C., D. Baier, I. Hörr, C. Raps, J. Berger, G. Jung & H. Schwarz.** 1999. Characterization of a novel, antifungal, chitin-binding protein from *Streptomyces tendae* Tü901 that interferes with growth polarity. J Bacteriol **181**:7421-7429.
- Bradford, M.M.** 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Any. Biochem. **72**:248-254.
- Briggs, M., and L. M. Gierasch.** 1986. Molecular mechanisms of protein secretion: the role of the signal sequence. Adv. Protein Chem. **38**:109-180.
- Brink S., E.G. Bogsch, A. Mant, C. Robinson.** 1997. Unusual characteristics of amino-terminal and hydrophobic domains in nuclear-encoded thylakoid signal peptides. Eur. J. Biochem. **245**:340-348.
- Brink S., E.G. Bogsch, W.R. Edwards, P.J. Hynds, C. Robinson.** 1998. Targeting of thylakoid proteins by the delta pH-driven twin-arginine translocation pathway requires a specific signal in the hydrophobic domain in conjunction with the twin-arginine motif. FEBS Lett. **434**:425-430.
- Bron S., R. Meima, J.M. van Dijl, A. Wipat & C. Harwood.** 1999. Molecular biology and genetics of *Bacillus* spp. In *Manual of Industrial Microbiology and Biotechnology* (Demain, A.L. & Davies J.E., eds), pp. 392- 416. ASM Press, Washington, D.C.
- Bron S., A. Bolhuis, H. Tjalsma, S. Holsappel, G. Venema and van J.M. Dijl.** 1998. Protein secretion and possible roles for multiple signal peptidases for precursor processing in Bacilli. J. Biotechnol. **64**: 3-13.
- Brüser T., R. Deutsmann, and C. Dahl.** 1998. Evidence against the double-arginine motif as the only determinant for protein translocation by a novel Sec-independent pathway in *Escherichia coli*. FEMS Microbiol. Lett. **164**:329-336.
- Bult C.J., O. White, G.J. Olsen, L. Zhou, R.D. Fleischmann, G.G. Sutton, et al.** 1996. Complete genome sequence of the methanogenic archaeon, *Methanococcus jannaschii*. Science. **273**:1058-1073.
- Chaddock A.M., A. Mant, I. Karnauchoy, S. Brink, R.G. Herrmann, R.B. Klosgen, C. Robinson.** 1995. A new type of signal peptide: central role of a twin-arginine motif in transfer signals for the delta pH-dependent thylakoidal protein translocase. EMBO J. **14**:2715-2722.
- Chang S. and S.N. Cohen.** 1979. High frequency transformation of *Bacillus subtilis* protoplasts by plasmid DNA. Mol. Gen. Genet. **168**:111-115.
- Chung, Y. J., M. T. Steen, and J. N. Hansen,** 1992. The subtilin gene of *Bacillus subtilis* ATCC 6633 is encoded in an operon that contains a homolog of the hemolysin B Transport protein. J. Bacteriol. **174**:1417-1422.

- Chung, Y. S., and D. Dubnau.** 1995. ComC is required for the processing and translocation of ComGC, a pilin-like competence protein of *Bacillus subtilis*. *Mol. Microbiol.* **15**:543-551.
- Chung, Y. S., and D. Dubnau.** 1998. All seven comG open reading frames are required for DNA binding during transformation of competent *Bacillus subtilis*. *J. Bacteriol.* **180**:41-45.
- Chung, Y. S., F. Breidt, and D. Dubnau.** 1998. Cell surface localization and processing of the ComG protein, required for DNA binding during transformation of *Bacillus subtilis*. *Mol. Microbiol.* **29**:905-913.
- Clark, S.A., and S.M. Theg.** 1997. A folded protein can be transported across the chloroplast envelope and thylakoid membranes. *Mol. Biol. Cell.* **8**:923-934.
- Cregg, K.M., I. Wilding & M.T. Black.** 1996. Molecular cloning and expression of the *spsB* gene encoding an essential type I signal peptidase from *Staphylococcus aureus*. *J. Bacteriol.* **178**:5712-5718.
- Cristobal, S., J. -W. de Gier, H. Nielsen, and G. von Heijne.** 1999. Competition between Sec- and Tat-dependent protein translocation in *Escherichia coli*. *EMBO J.* **18**:2982-2990.
- Cutting S. M. and P. B.Vander Horn.** 1990. In: **Harwood, C.R., Cutting, S.M.** (Eds.). *Molecular Biological Methods for Bacillus*. John Wiley & Sons. Ltd. p 27-74.
- Dalbey, R. E., and W. Wickner.** 1985. Leader peptidase catalyzes the release of exported proteins from the outer surface of the *Escherichia coli* plasma membrane. *J. Biol. Chem.* **260**:15925-15931.
- Dalbey, R. E., and G. von Heijne.** 1992. Signal peptidases in prokaryotes and eukaryotes: a new protease family. *Trends Biochem. Sci.* **17**:474-478.
- Dalbey, R. E., and C. Robinson.** 1999. Protein translocation into and across the bacterial plasma membrane and the plant thylakoid membrane. *Trends Biochem. Sci.* **24**:17-22.
- Dalbey, R. E., M. O. Lively, S. Bron, and J. M. van Dijk.** 1997. The chemistry and enzymology of the type I signal peptidases. *Protein Sci.* **6**:1129-1138.
- Date, T.** 1983. Demonstration by a novel genetic technique that leader peptidase is an essential enzyme of *Escherichia coli*. *J. Bacteriol.* **154**:76-83.
- de Gier, J. W., P. A. Scotti, A. Saaf, Q. A. Valent, A. Kuhn, J. Luirink, and G. von Heijne.** 1998. Differential use of the signal recognition particle translocase targeting pathway for inner membrane protein assembly in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **95**:14646-146451.
- Deshaies, R. J., F. Kepes, and P. C. Böhni.** 1989. Genetic dissection of the early stages of protein secretion in yeast. *Trends Genet.* **5**:87-93.
- Dricks, A.** 1999. *Bacillus subtilis* spore coat. *Microbiol. Mol. Biol. Rev.* **63**:1-20.
- Dubnau, D.** 1997. Binding and Transport of transforming DNA by *Bacillus subtilis*: the role of type-IV pilin-like proteins-a review. *Gene* **192**:191-198.
- Duong, F., J. Eichler, A. Price, M. R. Leonard, and W. Wickner.** 1997. Biogenesis of the Gram-negative bacterial envelope. *Cell* **91**:567-573.
- Economou, A.** 1998. Bacterial pre-protein translocase mechanism and conformational dynamic of a processive enzyme. *Mol. Microbiol.* **27**:511-518.

- Economou, A., and W. Wickner.** 1994. SecA promotes pre-protein translocation by undergoing ATP-driven cycles of membrane insertion and deinsertion. *Cell* **78**:835-843.
- Economou, A., J. A. Pogliano, J. Beckwith, D. B. Oliver, and W. Wickner.** 1995. SecA membrane cycling at SecYEG is driven by distinct ATP binding and hydrolysis events and is regulated by SecD and SecF. *Cell* **83**:1171-1181.
- Edens L., L. Heslinga, R. Klok, A.M. Ledebor, J. Maat, M.Y. Toonen, C. Visser & T. Verrips.** 1982. Cloning of cDNA encoding the sweet-tasting plant protein thaumatin and its expression in *Escherichia coli*. *Gene* **18**:1-12.
- Ewalt, K. L., J. P. Hendrick, W. A. Houry, and F. U. Hartl.** 1997. *In vivo* observation of polypeptide flux through the bacterial chaperonin System. *Cell* **90**:491-500.
- Foster, S.J.** 1992. Analysis of the autolysins of *Bacillus subtilis* 168 during vegetative growth and differentiation by using renaturing polyacrylamide gel electrophoresis. *J. Bacteriol.* **174**:464-470.
- Frandsen, N., and P. Stragier.** 1995. Identification and characterization of the *Bacillus subtilis* *spoIIIP* locus. *J. Bacteriol.* **177**:716-722.
- Franke, C. M., J. Tiemersma, G. Venema, and J. Kok.** 1999. Membrane topology of the lactococcal bacteriocin ATP-binding cassette transporter protein LcnC. Involvement of LcnC in lactococcal maturation. *J. Biol. Chem.* **274**:8484-8490.
- Fukamoto, J.** 1943. Studies on the production of bacterial amylase. I. Isolation of bacteria secreting potent amylase and their distribution. *J. Agric. Chem. Soc. Jpn.* **19**:487-503. (In Japanese)
- Goffeau A., J. Park, I.T. Paulsen, J.L. Jonniaux, T. Dinh, P. Mordant, M.H. Jr. Saier.** 1997. Multidrug-resistant transport proteins in yeast: complete inventory and phylogenetic characterization of yeast open reading frames with the major facilitator superfamily. *Yeast.* **13**:43-54.
- Gooday W. G.** 1990. The ecology of chitin degradation. In *Advances in microbial ecology*, pp. 387-430. Edited by K.C. Marshall. Plenum Press, New York.
- Gösseringer R., E. Kuster, A. Galinier, J. Deutscher & W. Hillen.** 1997. Cooperative and non-cooperative DNA binding modes of catabolite control protein CcpA from *Bacillus megaterium* result from sensing two different signals. *J Mol Biol* **266**:665-676.
- Hanahan D.** 1983. Studies on transformation of *Escherichia coli* with plasmids. *J. Mol. Biol.* **166**:557-580.
- Hartl, F. U.** 1998. Highlight: protein folding *in vivo*. *J. Biol. Chem.* **379**:235.
- Harwood C. R.** 1992. *Bacillus subtilis* and its relatives: molecular biological and industrial workhorses. *Trends Biotechnology.* **10**:247-256.
- Havarstein, L. S., D. B. Diep, and I. F. Nes.** 1995. A family of bacteriocin ABC transporters carry out proteolytic processing of their substrates concomitant with export. *Mol. Microbiol.* **16**:229-240.
- Henriques A.O., and C.P. Jr. Moran.** 2000. Structure and assembly of the bacterial endospore coat. *Methods.* **20**:95-110. Review.

- Hirose I., K. Sano, I. Shioda, M. Kumano, K. Nakamura & K. Yamane.** 2000. Proteome analysis of *Bacillus subtilis* extracellular proteins: a two-dimensional protein electrophoretic study. *Microbiology*. **146**:65-75.
- Hoang, V., and J. Hofemeister.** 1995. *Bacillus amyloliquefaciens* possesses a second type I signal peptidase with extensive sequence similarity to other *Bacillus* SPases. *Biochim. Biophys. Acta*. **1269**: 64-68.
- Hofmeister, A.** 1998. Activation of the proprotein transcription factor prosigma E is associated with its progression through three patterns of subcellular localization during sporulation in *Bacillus subtilis*. *J. Bacteriol.* **180**: 2426-2433.
- Hofmeister, A., A. Londono-Vallejo, E. Harry, P. Stragier., and R. Losick.** 1995. Extracellular signal protein triggering the proteolytic activation of a developmental transcription factor in *Bacillus subtilis*. *Cell* **83**:219-226.
- Hofemeister, J., M. Israelis-Reches & D. Dubnau.** 1983. Integration of plasmid pE194 at multiple sites on the *Bacillus subtilis* chromosome. *Mol. Gen. Genet.* **189**:58-68.
- Hueck C.J., W. Hillen & M.H.Jr. Saier.** 1994. Analysis of a cis-active sequence mediating catabolite repression in gram-positive bacteria. *Res Microbiol* **145**:503-518.
- Hynds P.J., D. Robinson, C. Robinson.** 1998. The sec-independent twin-arginine translocation system can transport both tightly folded and malfolded proteins across the thylakoid membrane. *J. Biol. Chem.* **273**:34868-34874.
- Illing, N., and J. Errington.** 1991. Genetic regulation of morphogenesis in *Bacillus subtilis*: roles of σ^E and σ^F in prespore engulfment. *J. Bacteriol.* **173**:3159-3169.
- Inada T., D.L. Court, K. Ito and Y. Nakamura.** 1989. Conditionally lethal amber mutations in the leader peptidase gene of *Escherichia coli*. *J. Bacteriol.* **171**:585-587.
- Ingle M.B. and E.W. Boyer.** 1976. Production of industrial enzymes by *Bacillus* species. In *Microbiology* (Schlessinger D. eds). pp. 420-426. American Society for Microbiology, Washington, D.C. 1976.
- Itoh Y., K. Kanoh, K. Nakamura, K. Takase, K. Yamane.** 1990. Artificial insertion of peptides between signal peptide and mature protein: effect on secretion and processing of hybrid thermostable alpha-amylases in *Bacillus subtilis* and *Escherichia coli* cells. *J. Gen. Microbiol.* **136**:1551-1558.
- Jiang, M., R. Grau and M. Perego.** 2000. Differential processing of propeptide inhibitors of Rap phosphatases in *Bacillus subtilis*. *J. Bacteriol.* **182**:303-310.
- Jones J. D., C. J. McKnight, L. M. Gierasch** 1990. Biophysical studies of signal peptides: implications for signal sequence functions and the involvement of lipid in protein export. *J. Bioenerg Biomembr.* **22**:213-32.
- Klenk, H.P., Clayton, R.A., Tomb, J.F., White, O., Nelson, K.E., et al.** 1997. The complete genome sequence of the hyperthermophilic, sulphate-reducing archaeon *Archaeoglobus fulgidus*. *Nature* **390**: 364-370.

- Klingenberg P., U. Behnke & A. Leuchtenberger.** 1985. Nutzung phänotypischer Merkmale für die Selektion leistungsgesteigerter Stämme des Thermitase-Produzenten *Thermoactinomyces vulgaris*. *Nahrung* **29**:969-978.
- Kolbe, S., S. Fischer, A. Becirevic, P. Hinz & H. Schrempf.** 1998. The *Streptomyces reticuli* α -chitin-binding protein CHB2 and its gene. *Microbiology* **144**:1291-1297.
- Kontinen V.P., P. Saris, M. Sarvas.** 1991. A gene (prsA) of *Bacillus subtilis* involved in a novel, late stage of protein export. *Mol. Microbiol.* **5**:1273-83.
- Kontinen V.P., and M. Sarvas.** 1993. The PrsA lipoprotein is essential for protein secretion in *Bacillus subtilis* and sets a limit for high-level secretion. *Mol. Microbiol.* **8**:727-737.
- Kunst, F., N. Ogasawara, I. Moszer, A. M. Albertini, G. Alloni, et al.** 1997, The complete genome sequence of the Gram-positive bacterium *Bacillus subtilis*. *Nature* **390**:249-256.
- Kuroda, A., Asami, Y. and Sekiguchi, J.** 1993. Molecular cloning of a sporulation-specific cell wall hydrolase gene of *Bacillus subtilis*. *J. Bacteriol.* **175**:6260-6268.
- Lazarevic, V., Margot, P., Soldo, B. and Karamata, D.** 1992. Sequencing and analysis of the *Bacillus subtilis* *lytRABC* divergon: a regulatory unit encompassing the structural genes of the *N*-acetylmuramoyl-L-alanine amidase and its modifier. *J. Gen. Microbiol.* **138**:1949-1969.
- Lazzazera, B. A., J. M. Solomon, and A. D. Grossman.** 1997. An exported peptide functions intracellularly to contribute to cell density signaling in *Bacillus subtilis*. *Cell* **13**:917-925.
- Laemmli, U. K.** 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**:680-685.
- Londono-Vallejo, J. -A., C. Frehel, and P. Stragier.** 1997. *spoIIQ*, a forespore-expressed gene required for engulfment in *Bacillus subtilis*. *Mol. Microbiol.* **24**:29-39.
- Lory, S.** 1994. Leader peptidase of type IV prepilins and related proteins. p. 17-29. *in* G. von Heijne (ed.), Signal peptidases. R. G. Landes Company, Austin, Tex.
- Lowry, O.H., N.J. Rosebrough, A.L. Farr and R.J. Randall.** 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**:265-275.
- Magnuson, R., J. Solomon, and A. D. Grossman.** 1994. Biochemical and genetic characterization of a competence pheromone from *Bacillus subtilis*. *Cell* **77**:207-216.
- Margot, P., Mauel, C. and Karamata, D.** 1994. The gene of the *N*-acetylglucosaminidase, a *Bacillus subtilis* 168 cell wall hydrolase not involved in vegetative cell autolysis. *Mol. Microbiol.* **12**:535-545.
- Margot, P., and D. Karamata.** 1996. The *wprA* gene of *Bacillus subtilis* 168, expressed during exponential growth, encodes a cell-wall-associated protease. *Microbiology* **142**:3437-3444.
- Margot, P., M. Pagni, and D. Karamata.** 1999. *Bacillus subtilis* 168 *lytF* encodes a γ -D-glutamate-meso-diaminopimelate muropeptidase expressed by the alternative vegetative sigma factor, σ^D . *Microbiology* **141**:2379-2389.
- McKenzie T., T. Hoshino, T. Tanaka, N. Sueoka.** 1987. Correction. A revision of the nucleotide sequence and functional map of pUB110. *Plasmid.* **17**:83-85.

- Meens J., M. Herbort, M. Klein and R. Freudl.** 1997. Use of the Pre-Pro part of *Staphylococcus hyicus* lipase as a carrier for secretion of Escherichia coli outer membrane protein A (OmpA) prevents proteolytic degradation of OmpA by cell-associated protease(s) in two different Gram-positive bacteria. *Appl Environ Microbiol.* **63**:2814-2820.
- Meijer, W.J.J., A. de Jong, G. Bea, A. Wisman, H. Tjalsma, G. Venema, Bron, S. and J.M. van Dijl.** 1995. The endogenous *Bacillus subtilis* (natto) plasmids pTA1015 and pTA1040 contain signal peptidase- encoding genes: identification of a new structural module on cryptic plasmids. *Mol. Microbiol.* **17**:621-631.
- Merchante, R., H. M. Pooley, and D. Karamata.** 1995. A periplasm in *Bacillus subtilis*. *J. Bacteriol.* **177**:6176-6183.
- Meyer, T.H., J.F. Menetret, R. Breitling, K.R. Miller, C.W. Akey and T.A. Rapoport.** 1999. The bacterial SecY/E translocation complex forms channel-like structures similar to those of the eukaryotic Sec61p complex. *J. Mol. Biol.* **285**:1789-1800.
- Mizobuchi M. and L.A. Frohman.** 1993. Rapid amplification of genomic DNA ends. *Biotechniques.* **15**:214-216.
- Montgomery M.T. & D.L. Kirchman.** 1994. Induction of chitin-binding proteins during the specific attachment of the marine bacterium *Vibrio harveyi* to chitin. *Appl Environ Microbiol* **60**:4284-4288.
- Moriyama, R., H. Fukuoka, S. Miyata, S. Kudoh, A. Hattori, S. Kozuka, Y. Yasuda, K. Tochikubo, and S. Makino.** 1999. Expression of a germination-specific amidase, SleB, of bacilli in the forespore compartment of sporulating cells and its localization on the exterior side of the cortex in dormant spores. *J. Bacteriol.* **181**:2373-2378.
- Msadek, T., V. Dartois, F. Kunst, ML. Herbaud, F. Denizot, G. Rapoport.** 1998. ClpP of *Bacillus subtilis* is required for competence development, motility, degradative enzyme synthesis, growth at high temperature and sporulation. *Mol. Microbiol.* **27**:899-914.
- Müller, P., K. Ahrens, T. Keller and A. Klucke.** 1995. A *TnphoA* insertion within the *Bradyrhizobium japonicum sipS* gene, homologous to prokaryotic signal peptidases, results in extensive changes in the expression of PBM-specific nodulins of infected soybean (*Glycine max*) cells. *Mol. Microbiol.* **18**:831-840.
- Müller, J., F. Walter, J. M. van Dijl, and D. Behnke.** 1992. Suppression of the growth and export defects of an *Escherichia coli secA(Ts)* mutant by a gene cloned from *Bacillus subtilis*. *Mol. Gen. Genet.* **235**:89-96.
- Müller, J., S. Bron, G. Venema, and J. M. van Dijl.** 2000. Chaperone-like activities of the CsaA protein of *Bacillus subtilis*. *Microbiology* **146**:77-88.
- Müller, J., J. Ozegowski, S. Vettermann, J. Swaving, K. H. van Wely, and A. J. M. Driessen.** 2000. Interaction of *Bacillus subtilis* CsaA with SecA and precursor proteins. *Biochem. J.* **348**:367-373.

- Nakamura K, Yahagi S, Yamazaki T, Yamane K.** 1999. *Bacillus subtilis* histone-like protein, HBSu, is an integral component of a SRP-like particle that can bind the Alu domain of small cytoplasmic RNA. *J. Biol. Chem.* **274**:13569-13576.
- Nakamura K., M. Nishiguchi, K. Honda, K. Yamane.** 1994. The *Bacillus subtilis* SRP54 homologue, Ffh, has an intrinsic GTPase activity and forms a ribonucleoprotein complex with small cytoplasmic RNA *in vivo*. *Biochem Biophys Res Commun.* **199**: 1394-1399.
- Nakano, M. M., and P. Zuber.** 1990. Molecular biology of antibiotic production in *Bacillus*. *Crit. Rev. Biotechnol.* **10**:223-240.
- Navarre, W. W., and O. Schneewind.** 1994. Proteolytic cleavage and cell wall anchoring at the LPXTG motif of surface proteins in Gram-positive bacteria. *Mol. Microbiol.* **14**:115-121.
- Navarre, W. W., S. Daepler, and O. Schneewind.** 1996. Cell Wall sorting of lipoproteins in *Staphylococcus aureus*. *J. Bacteriol.* **178**:441-446.
- Netzer, W. J., und F. U. Hartl.** 1998. Protein folding in the cytosol: chaperonin-dependent and -independent mechanisms. *Trends Biochem. Sci.* **23**:68-73.
- Ng DT, Brown JD, Walter P.** 1996. Signal sequences specify the targeting route to the endoplasmic reticulum membrane. *J. Cell Biol.* **134**:269-278.
- Nicholsen W.L. and P. Setlow.** 1990. Sporulation, germination and outgrowth. In *Molecular Biological Methods for Bacillus* (Harwood, C.R. & Cutting, S.M., eds), pp. 391-450. John Wiley & Sons. Ltd.
- Nielsen H., J. Engelbrecht, S. Brunak, G. von Heijne.** 1997. A neural network method for identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites. *Int. J. Neural. Syst.* **8**:581-599.
- Paetzel M., R.E. Dalbey, N.C. Strynadka.** 1998. Crystal structure of a bacterial signal peptidase in complex with a beta-lactam inhibitor. *Nature.* **396**:186-190.
- Paik, S. H., A. Chakicheria, and J. N. Hansen.** 1998. Identification and characterization of the structural and transporter genes for, and the chemical and biological properties of, sublancin 168, a novel lantibiotic produced by *Bacillus subtilis* 168. *J. Biol. Chem.* **273**:23134-23142.
- Parro, V., S. Schacht, J. Anne & R.P. Mellado.** 1999. Four genes encoding different type I signal peptidases are organized in a cluster in *Streptomyces lividans* TK21. *Microbiology* **145**:2255-2263.
- Perego, M., and J. A. Hoch.** 1996. Cell-cell communication regulates the effects of protein aspartate phosphatases on the phosphorelay controlling development in *Bacillus subtilis*. *Proc. Natl. Acad. Sci. USA* **93**:1549-1553.
- Perego, M.** 1997. A peptide export-import control circuit modulating bacterial development regulates protein phosphatases of the phosphorelay. *Proc. Natl. Acad. Sci. USA* **94**:8612-8617.
- Pohlschröder, M., W. A. Prinz, E. Hartmann, and J. Becwith.** 1997. Protein translocation in the three domains of life: variations on a theme. *Cell* **91**:563-566.

- Popham, D. L., B. Iliades-AGuiar, and P. Setlow.** 1995. The *Bacillus subtilis* *dacB* gene, encoding penicillin-binding protein 5*, is part of a three-gene operon required for proper spore cortex synthesis and spore core dehydration. *J. Bacteriol.* **177**:4721-4729.
- Popham, D. L., J. Meado-Parton, C. E. Costello, and P. Setlow.** 1999. Spore peptidoglycan structure in a *cwld dacB* double mutant of *Bacillus subtilis*. *J. Bacteriol.* **181**:6205-6209.
- Pooley, H. M., R. Merchante, and D. Karamata.** 1996. Overall protein content and induced enzyme components of the periplasm of *Bacillus subtilis*. *Microb. Drug Resist.* **2**:9-15.
- Pragai, Z., H. Tjalsma, A. Bolhuis, J. M. van Dijl, G. Venema, and S. Bron.** 1997. The signal peptidase II (*lsp*) gene of *Bacillus subtilis*. *Microbiology* **143**:1327-1333.
- Priest, F.G.** 1993. Systematic and ecology of *Bacillus*. In *Bacillus subtilis* and Other Gram-Positive Bacteria (Sonenshein, A.L., Hoch, J.A. & Losick, R., eds). pp. 3-16. American Society for Microbiology, Washington, D.C.
- Pugsley, A. P.** 1993. The complete general secretory pathway in gram-negative bacteria. *Microbiol. Rev.* **57**:50-108.
- Quentin, Y., G. Fichant, and F. Denizot.** 1999. Inventory, assembly and analysis of *Bacillus subtilis* ABC transport systems. *J. Mol. Biol.* **287**:467-484.
- Riezman, H.** 1997. The ins and outs of protein translocation. *Science* **278**:1728-1729.
- Robinson, C., D. Cai, A. Hulford, I.W. Brock, D. Michl, L. Hazell, I. Schmidt, R.G. Herrmann, R.B. Klosgen.** 1994. The presequence of a chimeric construct dictates which of two mechanisms are utilized for translocation across the thylakoid membrane: evidence for the existence of two distinct translocation systems. *EMBO J.* **13**:279-285.
- Robinson, C., and A. Mant.** 1997. Targeting of proteins into and across the thylakoid membrane. *Trends in Plant Science* **2**: 431-437.
- Rosenthal, A.L. & Lacks, S.A.** 1977. Nuclease detection in SDS-polyacrylamide gel electrophoresis. *Analytical Biochem.* **80**:76-90.
- Sahl, H. G., R. W. Jack, and G. Bierbaum.** 1995. Biosynthesis and biological activities of lantibiotics with unique post-translational modifications. *Eur. J. Biochem.* **230**:827-853.
- Saito, J., A. Kita, Y. Higuchi, Y. Nagata, A. Ando & K. Miki.** 1999. Crystal structure of chitosanase from *Bacillus circulans* MH-K1 at 1.6-Å resolution and its substrate recognition mechanism. *J Biol Chem* **274**:30818-30825.
- Sambrook J., E.J. Fritsch and T. Maniatis.** 1989. *Molecular Cloning: a Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- Sargent F, Bogsch EG, Stanley NR, Wexler M, Robinson C, Berks BC, Palmer T.** 1998. Overlapping functions of components of a bacterial Sec-independent protein export pathway. *EMBO J.* **17**:3640-3650.
- Schatz, G., and B. Dobberstein.** 1996. Common principles of protein translocation across membranes. *Science* **271**:1519-1526.

- Schneewind, O., A. Fowler, and K. F. Faull.** 1995. Structure of the cell wall anchor of surface proteins in *Staphylococcus aureus*. *Science* **268**:103-106.
- Schneewind, O., D. Mihaylova-Petkov, and P. Model.** 1993. cell wall sorting signals in surface proteins of Gram-positive bacteria. *EMBO J.* **12**:4803-4811.
- Schneewind, O., P. Model, and V. A. Fischetti.** 1992. Sorting of protein A to the staphylococcal cell wall. *Cell* **70**:267-281.
- Schnellmann, J., A. Zeltins, H. Blaak & H. Schrempf.** 1994. The novel lectin-like protein CHB1 is encoded by a chitin-inducible *Streptomyces olivaceoviridis* gene and binds specifically to crystalline α -chitin of fungi and other organisms. *Mol Microbiol* **13**:807-819.
- Schrempf, H.** 1999. Chitin-binding proteins in streptomycetes. In *Chitin and Chitinases*, pp. 99-108. Edited by P. Jollès & R. A. A. Muzzarelli. Birkhäuser Verlag.
- Sekiguchi, J., K. Akeo, H. Yamamoto, F. K. Khasanov, J. C. Alonso, A. Kuroda.** 1995. Nucleotide sequence and regulation of a new putative cell wall hydrolase gene, *cwlD*, which affects germination in *Bacillus subtilis*. *J. Bacteriol.* **177**:5582-5589.
- Serrano, M., R. Zilhao, E. Ricca, A. J. Ozin, C. P. Moran Ir, and A. O. Henriques.** 1999. A *Bacillus subtilis* secreted protein with a role in endospore coat assembly and function. *J. Bacteriol.* **181**:3632-3643.
- Settles A.M., A. Yonetani, A. Baron, D.R. Bush, K. Cline, R. Martienssen.** 1997. Sec-independent protein translocation by the maize Hcf106 protein. *Science.* **278**:1467-1470.
- Simonen, M., and I. Palva.** 1993. Protein secretion in *Bacillus* species. *Microbiol. Rev.* **57**:109-137.
- Smith, T. J., Blackman, S. A. and Foster, S. J.** 1996. Peptidoglycan hydrolases of *Bacillus subtilis* 168. *Microb. Drug Resist* **2**:113-118.
- Smith, T. J., Blackman, S. A. and Foster, S. J.** 2000. Autolysins of *Bacillus subtilis*: multiple enzymes with multiple functions. *Microbiology* **146**:249-262.
- Smith D.R., L.A. Doucette-Stamm, C. Deloughery, H. Lee, J. Dubois, et al.** 1997. Complete genome sequence of *Methanobacterium thermoautotrophicum* deltaH: functional analysis and comparative genomics. *J. Bacteriol.* **179**:7135-7155.
- Solomon, J. M., R. Magnuson, A. Srivastava, and A. D. Grossman.** 1995. Convergent sensing pathways mediate **response to two** extracellular competence factors in *Bacillus subtilis*. *Genes Dev.* **9**:547-558.
- Stackebrandt, E., W. Ludwig, M. Weizenegger, S. Dorn, T. McGill, G.E. Fox, C.R. Woese, W. Schubert & K.-H. Schleifer.** 1987. Comparative 16S rRNA oligonucleotide analyses and murein types of round-spore-forming bacilli and non-spore-forming relatives. *J. Gen. Microbiology* **133**:2523-2529.
- Stein, T., S. Borchert, B. Conrad, J. Feesche, B. Hofemeister, K. –D. Entian and J. Hofemeister.** 2001. Two ericins originate from the lantibiotic gene cluster of *Bacillus subtilis* A1/3. *J. Bacteriol.* (in submitting).

- Steinmetz, M., F. Kunst and R. Dedonder.** 1976. Mapping of mutations affecting synthesis of extracellular enzymes in *Bacillus*. Identity of *sacU*, *amyB* and *pap* mutations. *Mol. Gen. Genet.* **148**:281-285.
- Stöver, A. G., and A. Driks.** 1999a. Secretion, localization, and antibacterial activity of TasA, a *Bacillus subtilis* spore-associated protein. *J. Bacteriol.* **181**:1664-1672.
- Stöver, A. G., and A. Driks.** 1999b. Regulation of synthesis of the *Bacillus subtilis* transition-phase, spore-associated antibacterial protein TasA. *J. Bacteriol.* **181**:5476-5481.
- Stöver, A. G., and A. Driks.** 1999c. Control of synthesis and secretion of the *Bacillus subtilis* protein YqxM. *J. Bacteriol.* **181**:7065-7069.
- Stragier, P., and R. Losick.** 1996. Molecular genetics of sporulation in *Bacillus subtilis*. *Annu. Rev. Genet.* **30**:297-341.
- Stragier P., C. Bonamy and C. Karmazyn-Campelli.** 1988. Processing of a sporulation sigma factor in *Bacillus subtilis*: How morphological structure could control gene expression. *Cell* **52**:697-704.
- Stülke J. & W. Hillen.** 2000. Regulation of carbon catabolism in *Bacillus* species. *Annu Rev Microbiol* **54**:849-880.
- Suetake T., S. Tsuda, S. Kawabata, K. Miura, S. Iwanaga, K. Hikichi, K. Nitta & K. Kawano.** 2000. Chitin-binding proteins in invertebrates and plants comprise a common chitin-binding structural motif. *J Biol Chem.* **275**:17929-17932.
- Sung M. and R.E. Dalbey.** 1992. Identification of potential active-site residues in the *Escherichia coli* leader peptidase. *J. Biol Chem.* **267**:13154-13159.
- Suzuki, K., M. Suzuki, M. Taiyoji, N. Nikaidou & T. Watanabe.** 1998. Chitin binding protein (CBP21) in the culture supernatant of *Serratia marcescens* 2170. *Biosci Biotechnol Biochem* **62**:128-135.
- Svergun D.I., A. Becirevic, H. Schrempf, M.H. Koch and G. Gruber.** 2000. Solution structure and conformational changes of the *Streptomyces* chitin-binding protein (CHB1). *Biochemistry.* **39**:10677-10683.
- Swofford, D. L.** 2001. PAUP*. Phylogenetic analysis using parsimony (*and other methods). Version 4. Sinauer Associates, Sunderland, Massachusetts.
- Takami H, K. Nakasone, Y. Takaki, G. Maeno, R. Sasaki, N. Masui, F. Fuji, et al.** 2000. Complete genome sequence of the alkaliphilic bacterium *Bacillus halodurans* and genomic sequence comparison with *Bacillus subtilis*. *Nucleic Acids Res.* **28**:4317-4331.
- Takase K., H. Mizuno, K. Yamane.** 1988. NH₂-terminal processing of *Bacillus subtilis* alpha-amylase. *J. Biol. Chem.* **263**:11548-11553.
- Taura T., Y. Akiyama, K. Ito.** 1994. Genetic analysis of SecY: additional export-defective mutations and factors affecting their phenotypes. *Mol. Gen. Genet.* **243**:261-269.
- Tjalsma, H., M. A. Noback, S. Bron, G. Venema, K. Yamane, and J. M. van Dijl.** 1997. *Bacillus subtilis* contains four closely related type I signal peptidase with overlapping substrate specificities: constitutive and temporally controlled expression of different genes. *J. Biol. Chem.* **272**:25983-25992.

- Tjalsma, H., A. Bolhuis, M. L van Roosmalen, T. Wiegert, W. Schumann, C. P. Broekhuizen, W. Quax, G. Venema, S. Bron, and J. M. van Dijl.** 1998. Functional analysis of the secretory precursor processing machinery of *Bacillus subtilis*: identification of an eubacterial homolog of archaeal and eukaryotic signal peptidases. *Genes Dev.* **12**:2318-2331.
- Tjalsma, H., V. P. Kontinen, Z. Pragai, H. Wu, R. Meima, G. Venema, S. Bron, M. Sarvas, and J. M. van Dijl.** 1999a. The role of lipoprotein processing by signal peptidase II in the Gram-positive eubacterium *Bacillus subtilis* signal peptidase II is required for the efficient secretion of α -amylase, a non-lipoprotein. *J. Biol. Chem.* **274**:1698-1707.
- Tjalsma, H., J. van den Dolder, W. J. Meijer, G. Venema, S. Bron, and J.M. van Dijl.** 1999b. The plasmid-encoded type I signal peptidase SipP can functionally replace the major signal peptidases SipS and SipT of *Bacillus subtilis*. *J. Bacteriol.* **181**:2448-2454.
- Tjalsma, H., A. G. Stöver, A. Driks, G. Venema, S. Bron, and J. M. van Dijl.** 2000a. Conserved serine and histidine residues are critical for activity of the ER-type signal peptidase SipW of *Bacillus subtilis*. *J. Biol. Chem.* **275**:25102-25108.
- Tjalsma, H., A. Bolhuis, J. Jongbloed, S. Bron, and J. M. van Dijl.** 2000. Signal peptide-dependent protein transport in *Bacillus subtilis*: a genome-base survey of the secretome. *Microbiol. Mol. Biol. Rev.* **64**:515-547.
- Tschantz, W. R., M. Sung, V. M. Delgado-Partin, and R. E. Dalbey.** 1993. A serine and a lysine residue implicated in the catalytic mechanism of the *Escherichia coli* leader peptidase. *J. Biol. Chem.* **268**:27349-27354.
- Valent, Q. A., P. A. Scotti, S. High, J. W. de Gier, G. von Heijne, G. Lentzen, W. Wintermeyer, B. Oudega, and J. Luirink.** 1998. The *Escherichia coli* SRP and SecB targeting pathways converge at the translocon. *EMBO J.* **17**:2504-2512.
- van Dijl, J. M., A. de Jong, G. Venema, and S. Bron.** 1995. Identification of the Potential active site of the signal peptidase SipS of *Bacillus subtilis*: structural and functional similarities with LexA-like proteases. *J. Biol. Chem.* **270**:3611-3618.
- van Dijl, J. M., A. de Jong, J. Vehmannperä, G. Venema, and S. Bron.** 1992. Signal peptidase I of *Bacillus subtilis*: patterns of conserved amino acids in prokaryotic and eukaryotic type I Signal peptides. *EMBO J.* **11**:2819-2828.
- van Roosmalen, M.L., J.D.H. Jongbloed, A. Kuipers, G. Venema, S. Bron and J.M. van Dijl.** 2000. A truncated soluble *Bacillus* signal peptidase produced in *Escherichia coli* is subject to self-cleavage at its active site. *J. Bacteriol.* **182**:5765-5770.
- van Roosmalen, M.L., J.D.H. Jongbloed, J.-Y. F. Dubois, G. Venema, S. Bron & J.M. van Dijl.** 2001. Distinction between major and minor *Bacillus* signal peptidases based on phylogenetic and structural criteria. *J. Biol. Chem.* **276**:25230-25235.
- van Sinderen, D., R. Kiewiet, and G. Venema.** 1995. Differential expression of two closely related deoxyribonuclease genes, *nucA* and *nucB*, in *Bacillus subtilis*. *Mol. Microbiol.* **15**:213-223.

- Vaupel M. & R.K. Thauer.** 1995. Coenzyme F420-dependent N5,N10-methylenetetrahydromethanopterin reductase (Mer) from *Methanobacterium thermoautotrophicum* strain Marburg. Cloning, sequencing, transcriptional analysis, and functional expression in *Escherichia coli* of the *mer* gene. *Eur J Biochem* **231**:773-778.
- Vehmaanperä J.** 1989. Transformation of *Bacillus amyloliquefaciens* by electroporation. *FEMS Microbiol. Lett.* **61**:165-170.
- Vehmaanperä J., G. Steinborn & J. Hofemeister.** 1991. Genetic manipulation of *Bacillus amyloliquefaciens*. *J. Biotechnol.* **19**:221-240.
- von Heijne, G.** 1989. The structure of signal peptides from bacterial lipoproteins. *Protein Eng.* **2**:531-534.
- von Heijne, G.** 1990a. Protein targeting Signals. *Curr. Opin. Cell Biol.* **2**:604-608.
- von Heijne, G.** 1990b. The signal peptide. *J. Membr. Biol.* **115**:195-201.
- von Heijne, G.** 1998. Life and death of the Signal peptide. *Nature* **396**:111-113.
- von Heijne, G., and L. Abrahmsen.** 1989. Species-specific variation in signal peptide design. Implications for protein secretion in foreign hosts. *FEBS Lett.* **27**:439-446.
- Wallin E., and G. von Heijne.** 1998. Genome-wide analysis of integral membrane proteins from eubacterial, archaean, and eukaryotic organisms. *Protein Sci.* **7**:1029-1038.
- Watanabe T., W. Oyanagi, K. Suzuki, K. Ohnishi & H. Tanaka.** 1992. Structure of the gene encoding chitinase D of *Bacillus circulans* WL-12 and possible homology of the enzyme to other prokaryotic chitinases and class III plant chitinases. *J Bacteriol* **174**:408-414.
- Weiner, J. H., P. T. Bilous, G. M. Shaw, S. P. Lubitz, L. Frost, G. H. Thomas, J. A. Cole, R. J. Turner.** 1998. A novel and ubiquitous system for membrane targeting and secretion of cofactor-containing proteins. *Cell* **93**:93-101.
- Wexler, M., E. Bogsch, R. B. Klösgen, T. Palmer, C. Robinson, and B. C. Berks.** 1998. Targeting signals for a bacterial Sec-independent export system direct plant thylakoid import by the Δ pH pathway. *FEBS Lett.* **431**:339-342.
- Youngman P.** 1990. In: **Harwood, C.R., Cutting, S.M.** (Eds.). *Molecular Biological Methods for Bacillus*. John Wiley & Sons. Ltd. p 221-266.
- Zeltins, A. and H. Schrempf.** 1995. Visualization of α -chitin with a specific chitin-binding protein (CHB1) from *Streptomyces olivaceoviridis*. *Anal Biochem* **231**:287-294.
- Zeltins, A. & H. Schrempf.** 1997. Specific interaction of the *Streptomyces* chitin-binding protein CHB1 with α -chitin: the role of individual tryptophan residues. *Eur J Biochem* **246**:557-564.
- Zhang, Y.B., Greenberg, B. & S.A. Lacks.** 1997. Analysis of a *Streptococcus pneumoniae* gene encoding signal peptidase I and overproduction of the enzyme. *Gene* **194**:249-255.
- Zheng, G., L. Z. Yan, J. C. Wederas, and P. J. Zuber.** 1999. Genes of the *sbo-alb* locus of *Bacillus subtilis* are required for production of the antilisterial bacteriocin subtilosin. *J. Bacteriol.* **181**:7346-7355.

Zuber, P., M. M. Nakano, and M. A. Marahiel. 1993. Peptide antibiotics. In *Bacillus subtilis and other Gram-positive Bacteria*, pp. 381-410. Edited by **J. A. Hoch and R. Losick**. Washington, DC: American Society for Microbiology.