

Matrix-bound peptides modeling protein- protein interactions

Dissertation

zur Erlangung des akademischen Grades

Doctor rerum naturalium

Vorgelegt dem Fachbereich Biochemie / Biotechnologie
der Mathematisch-Naturwissenschaftlich-Technischen Fakultät
der Martin-Luther-Universität Halle-Wittenberg

von

Chao Yu

geb. am 25.06.1971 in Shanghai, P. R. China

Halle/S., Juni 2003

1. Gutachter: Prof. Dr. G. Fischer (MPG FS für Enzymologie der Proteinfaltung, Halle)
2. Gutachter: Prof. Dr. S. Reißmann (Friedrich-Schiller-Universität Jena)
3. Gutachter: Prof. Dr. K. Neubert (Martin-Luther-Universität Halle-Wittenberg)

Datum der Verteidigung: 20.10.2003

Table of contents

1 Introduction	1
1.1 Sterically constrained oligopeptides with induced conformations	1
1.1.1 Peptides bound to scaffolds	2
1.1.2 Peptides bound to solid phase	3
1.2 Synthesis of peptide arrays on planar supports — SPOT technology	6
1.2.1 SPOT-synthesis technique	7
1.2.2 The supports used in SPOT synthesis	9
1.2.3 Analysis of protein-protein/peptide contact sites based on SPOT synthesis	9
1.2.4 Mapping linear and discontinuous binding sites by standard SPOT strategy	11
1.3 Prolyl <i>cis/trans</i> isomerization – a probe for structural dynamics of proline-containing polypeptides	12
1.3.1 Prolyl <i>cis/trans</i> isomerization in peptides containing proline mimetics	14
1.3.2 Properties of <i>cis/trans</i> isomerization of peptide bond preceding fluorinated prolines	15
1.4 Aims	18
2 Results and discussion	20
2.1 Prerequisites relating to Janus-peptide arrays by SPOT synthesis	20
2.1.1 Scheme of screening matrix-bound peptide-peptide interaction in mapping protein-protein interaction sites — Janus-peptide array	20
2.1.2 Amino group loading on planar polymeric supports	22
2.1.3 Peptide quality on planar polymeric supports	22
2.1.4 Templates used in Janus-peptide arrays	25
2.2 The examples of protein-peptide interaction	26
2.2.1 Example I: Streptavidin/ <i>Strep</i> -tag II interaction	26
2.2.1.1 Self-recognition in streptavidin subunit association	27
2.2.1.2 Mapping binding epitopes on streptavidin to <i>Strep</i> -tag II by Janus-peptide arrays	29
2.2.1.3 Biotin blocks the binding of streptavidin to <i>Strep</i> -tag II in the Janus-peptide array	31
2.2.1.4 Comparison of two different templates used for Janus-peptide arrays	32
2.2.1.5 The quality of the synthetic Janus-peptide pairs	34
2.2.1.6 Peptide length variation for the minimized binding epitopes by Janus-peptide array	35
2.2.1.7 Failure in analyzing the binding epitopes on streptavidin by fluorescence labeled <i>Strep</i> -tag II using standard SPOT strategy	36
2.2.1.8 Janus-peptide arrays based on the reverse sequence of streptavidin and <i>Strep</i> -tag II	38
2.2.1.9 Using polypropylene membrane for Janus-peptide arrays	39
2.2.1.10 Conclusions	42

2.2.2	Example II: Protein 14-3-3/phosphopeptides interaction	43
2.2.2.1	Mapping binding epitopes on 14-3-3 to RQRSTpSTPNV (Raf peptide) by Janus-peptide arrays	45
2.2.2.2	Mapping binding epitopes on 14-3-3 to ARSHpSYPA (mT peptide) by Janus-peptide arrays	47
2.2.2.3	Trouble shooting for the synthesis of phosphopeptides on cellulose	48
2.2.2.4	Conclusions	49
2.3	The examples of protein-protein interaction	51
2.3.1	Example I: FKBP12/FAP48 interaction	53
2.3.1.1	Mapping binding epitopes on FAP48 to FKBP12 by standard SPOT strategy	53
2.3.1.2	Mapping binding epitopes on FKBP12 to FAP48 using Janus-peptide arrays	54
2.3.1.3	Conclusions	58
2.3.2	Example II: FKBP12/EGFR cytosolic domain (residues 645-1186) interaction	61
2.3.2.1	Mapping binding epitopes on EGFR (Arg ⁶⁴⁵ -Ala ¹¹⁸⁶) to FKBP12 by standard SPOT strategy	61
2.3.2.2	The inhibitory activities (IC ₅₀) of several peptide epitopes derived from EGFR on the PPIase activity of FKBP12	64
2.3.2.3	Mapping binding epitopes on FKBP12 to EGFR (Arg ⁶⁴⁵ -Ala ¹¹⁸⁶) by using Janus-peptide arrays	65
2.3.2.4	Interactions between Janus-peptide pairs are sequence dependent	66
2.3.2.5	Conclusions	68
2.4	General properties of Janus-peptide arrays	68
2.4.1	The properties of Janus-peptide arrays	69
2.4.2	The potential drawbacks of Janus-peptide arrays	71
2.4.3	The evaluation of the results from Janus-peptide arrays	72
2.5	Thermodynamic and kinetic parameters of the <i>cis/trans</i> isomerization of (4)-fluoroproline containing peptides and the influence of PPIases	74
2.5.1	<i>Cis/trans</i> isomerization of (4)-fluoroproline containing peptides	74
2.5.2	Catalysis of the <i>cis/trans</i> isomerization of fluoroproline containing peptide substrates by PPIases	77
2.5.3	The influence of urea on the activity of different PPIases	80
2.5.4	Conclusions	83
3	Materials and Methods	85
3.1	Materials	85
3.2	Methods	91
3.2.1	Expression and purification of hCyp18	91
3.2.2	Expression and purification of hFKBP12	92
3.2.3	MALDI-TOF analysis of cellulose bound peptide spots	92
3.2.4	Modification of cellulose membrane with (β-Ala) ₂ anchor functions	93

3.2.5	The synthesis of templates used in Janus-peptide arrays	94
3.2.6	The preparation of Janus-peptide membranes	95
3.2.7	Western blot analysis	96
3.2.8	Membrane regeneration	97
3.2.9	SDS-Polyacrylamide Gel Electrophoreses (SDS-PAGE)	97
3.2.10	Determination of urea concentration	97
3.2.11	Determination of protein concentration with UV/VIS-spectroscopy	98
3.2.12	Determination of protein concentration with Bradford method	98
3.2.13	The catalytic efficiency of peptidyl-prolyl <i>cis/trans</i> isomerases	99
3.2.14	Influence of PPIases on the <i>cis/trans</i> isomerization of (4)-fluoroproline containing peptides (protease-coupled assay)	100
3.2.15	Solvent-jump method (protease-free assay)	101
3.2.16	The inhibition of soluble peptides on the PPIase activity of <i>hFKBP12</i>	101
	Summary	102
	References	104
	Acknowledgements	
	Appendix A	
	Appendix B	

Abbreviations

(4)-diF-Pro	4 <i>R</i> , 4 <i>S</i> - L-di-fluoroproline
(4 <i>R</i>)-FPro	(4 <i>R</i>)-L-fluoroproline
(4 <i>S</i>)-FPro	(4 <i>S</i>)- L-fluoroproline
4-Oxa	(<i>S</i>)-oxazolidine-4-carboxylic acid
4-Thz	(<i>R</i>)-thiazolidine-4-crboxylic acid
Abz	2-aminobenzoyl
Ac-	Acetyl-
APEG	amino PEG (Poly-ethylene-glycol) spacer
AU	Absorption Unit
b*pwt	Barstar pseudo-wildtype
Boc	<i>tert</i> -butyloxycarbonyl
BPB	bromophenol blue
BSA	Bovines Serumalbumin
Bzl	Benzyl
Cyp	Cyclophilin
DCC	N,N'-Dicyclohexylcarbodiimide
DCM	Dichloromethane
DIC	N,N'-Diisopropylcarbodiimide
Dde	1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)ethyl
DEAE	Diethylaminoethyl-
Dhbt	3-hydroxy-2,3-dihydroxy-4-oxo-benzotriazolyl
DIEA	Diisopropylethylamine
DIPCDI	N,N'-Diisopropylcarbodiimide
DMF	Dimethylformamide
DMSO	Dimethylsulfoxide
DTT	Dithiothreitol
ECL	Enhanced chemiluminescence
<i>E. coli</i> Par	<i>E. coli</i> Parvulin
<i>E. coli</i> TF	<i>E. coli</i> Trigger Factor
EDTA	Ethylenediamine-tetraaceticacid
EGFR	EGF receptor
ESI-MS	Electrospray ionization mass spectrometry
EtOH	Ethanol
FAP	FKBP-associated protein
FKBP	FK506-binding Protein
Flp	4(<i>R</i>)-fluoro-L-proline
flp	4(<i>S</i>)-fluoro-L-proline
FMDV	foot-and mouth disease virus
Fmoc	9-fluorenylmethoxycarbonyl
FT-IR	Fourier Transform Infrared Spectroscopy
HBTU	2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
HEPES	4-(2-Hydroxyethyl)-1-piperazine-1-ethanesulfonic acid
HF	hydrogen fluoride
HOBt	1-hydroxybenzotriazole
HRMAS	high-resolution magic angle spinning
HRP	Horse-Radish Peroxidase
Hyp	4(<i>R</i>)-hydroxy-L-proline

IPTG	Isopropyl β -D-thiogalactopyranoside
ISP	Isomer-Specific Proteolysis
k_{cat}	catalytic constant
K_d	Dissociation constant
K_m	Michaelis-Menten constant
MALDI-TOF	Matrix assisted laser desorption ionisation time-of-flight
MBHA	Methylbenzylamine
MES	2-(<i>N</i> -Morpholin-o)ethanesulfonic acid
MLU	Martin-Luther-University
MS	Mass spectrometry
NMI	<i>N</i> -methylimidazole
NMP	<i>N</i> -methylpyrrolidin
NMR	Nuclear Magnetic Resonance
OD	optical density
PAGE	Polyacrylamide gelelectrophoresis
Par	Parvulin
Pbf	2,2,4,6,7-Pentamethyl-dihydrobenzofurane-5-sulfonyl
PEG	Polyethyleneglycol
PEGA	acrylamidopropyl-PEG- <i>N,N</i> -dimethylacrylamide
Pfp	Pentafluorophenyl
PMSF	Phenylmethylsulfonylfluoride
-pNA	- <i>para</i> -nitroaniline
POEPOP	polyoxyethylene-polyoxypropylene
PP	Polypropylene
PPIase	Peptidyl-Prolyl- <i>cis/trans</i> -Isomerase
PTFE	polytetrafluoroethylene
PyBOP	benzotriazole-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate
Rink	4-(2',4'-Dimethoxyphenyl-hydroxymethyl)-phenoxyacetic acid
RT	room temperature
SDS	Sodium dodecylsulfate
Sip	Silaproline
SPPS	Solid Phase Peptide Synthesis
Suc-	Succinyl-
TASP	template-assembled synthetic protein
TBS	Tris buffered saline
TBT	Tris buffered saline with Tween 20
TBTU	2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate
tBu	t-Butyl
TEMED	<i>N,N,N',N'</i> -Tetramethylethylenediamine
TF	Trigger factor
TFA	Trifluoroacetic acid
TOF-MS	Time-of-flight mass spectrometry
Tricine	<i>N</i> -Tris(hydroxymethyl)-methylglycine
Tris	2-Amino-2-(hydroxymethyl)-1,3-propane-1,3-diol
Trt	Trityl
UV/VIS	Ultraviolet/Visible
wt	Wild type
Xaa	amino acids