

Literature

- Adler K, Scholz G** (1986) Die Chloroplastenstrukturen der Mutante chloronerva von *Lycopersicon esculentum* Mill. im Mutantenstatus und nach Normalisierung mit Nicotianamin. *Kulturpflanze* **34**: 185-194
- Alcántara E, Romera FJ, Canete M, De la Guardia MD** (1994) Effects of heavy metals on both induction and function of root Fe (III) reductase in Fe-deficient cucumber (*Cucumis sativus* L.) plants. *Journal of Experimental Botany* **45**: 1893-1998
- Becker R, Fritz E, Manteuffel R** (1995) Subcellular localization and characterization of excessive iron in the nicotianamine-less tomato mutant *chloronerva*. *Plant Physiol* **108**: 269-275
- Bienfait HF** (1988) Mechanisms in Fe-efficiency reactions of higher plants. *J Plant Nutr* **11**: 605–629.
- Bintrim SB, Donohue TJ, Handelsman J, Roberts GP, Goodman RM** (1997) Molecular phylogeny of Archaea from soil. *Proc Natl Acad Sci USA* **94**: 277-282
- Buckhout TJ, Bell PF, Luster DG, Chaney RL** (1989) Iron-stress induced redox activity in tomato (*Lycopersicon esculentum* Mill.) is localized on the plasma membrane. *Plant Physiology* **90**: 151–156.
- Buděšinský M, Budesinsky H, Prochazka Z, Ripperger H, Römer A, Scholz G, Schreiber K** (1980) Nicotianamine, a possible phytosiderophore of general occurrence. *Phytochemistry* **19**: 2295 – 2297
- Chosack E, Rubinstein B, Reinhold L** (1991) Effect of iron deficiency on electron transport processes of plasmalemma-enriched vesicles isolated from cotton roots. *Plant Science* **77**: 163–172.
- Church GM and Gilbert W** (1984) Genomic Sequencing. *Proc Natl Acad Sci USA* **81**: 1991-1995
- Clemens S** (2000) Molecular mechanisms of plant metal tolerance and homeostasis. *Planta* **212**: 475-486
- Christensen AH and Quail PH** (1996) Ubiquitin promoter-based vectors for high-level expression of selectable and/or screenable marker genes in monocotyledonous plants. *Transgen Res* **5**: 213-218
- Dell'Orto M, Santi S, De Nisi P, Cesco S, Varanini Z, Zocchi G, Pinton R** (2000) Development of Fe-deficiency responses in cucumber (*Cucumis sativus* L.) roots: involvement of plasma membrane H(+)-ATPase activity. *J Exp Bot* **51**(345): 695-701
- Douchkov D, Herbig A, Koch G, Mock H-P, Melzer M, Stephan UW, Bäumlein H** (2002) Nicotianamine synthase: Gene isolation, gene transfer and application for manipulation of plant iron assimilation. *Plant and Soil* **241**: 115-119

- Drechsel H, Jung G** (1998) Peptide Siderophores. *J of Peptide Science* **4**:147-181
- Eide D, Broderius M, Fett J, Guerinot ML** (1996) A novel iron-regulated metal transporter from plants identified by functional expression in yeast. *Proc Natl Acad Sci USA* **93**: 5624-5628
- Fujiki Y, Ito M, Nishida I, Watanabe A** (2001) Leucine and its keto acid enhance the coordinated expression of genes for branched-chain amino acid catabolism in Arabidopsis under sugar starvation. *FEBS Lett* **499**(1-2): 161-165
- Fujiki Y, Sato T, Ito M, Watanabe A** (2000) Isolation and characterization of cDNA clones for the e1beta and E2 subunits of the branched-chain alpha-ketoacid dehydrogenase complex in Arabidopsis. *J Biol Chem* **275**(8): 6007-6013
- Guerinot ML** (2000) The ZIP family of metal transporters. *Biochim Biophys Acta* **1465**(1-2): 190-198
- Guerinot ML** (2001) Improving rice yields-iron out the details. *Nature Biotechnology* **19**: 417-418
- Halliwell B, Gutteridge JMC** (1984) Oxygen toxicity, oxygen radicals, transition metals and disease. *Biochem J* **219**: 1-14
- Hentze MW, Caughman SW, Rouault TA, Barriocanal JG, Dancis A, Harford JB, Klausner RD** (1987) Identification of the iron-responsive element for the translational regulation of human ferritin mRNA. *Science* **238**(4833): 1570-3
- Herbik A** (1997) Proteinchemische und molekularbiologische Charakterisierung der Tomatenmutante *chloronerva*. *Dissertation, Humboldt-Universität zu Berlin*
- Herbik A, Koch G, Mock H-P, Dushkov D, Czihal A, Thielmann J, Stephan UW, Bäumlein H** (1999) Isolation, characterization and cDNA cloning of nicotianamine synthase from barley - a key enzyme for iron homeostasis in plants. *Eur J Biochem* **264**: 1-10
- Higuchi K, Kanazawa K, Nishizawa N-K, Chino M, Mori S** (1994) Purification and characterization of nicotianamine synthase from Fe-deficient barley roots. *Plant and Soil* **165**: 173-179
- Higuchi K, Nishizawa N-K, Yamaguchi H, Römheld V, Marschner H, Mori S** (1995) response of nicotianamine synthase activity to Fe-deficiency in tobacco plants as compared with barley. *J Exp Botany* **46**(289) 1061-1063
- Higuchi K, Suzuki K, Nakanishi H, Yamaguchi H, Nishizawa N-K, Mori S** (1999) Cloning of Nicotianamine Synthase Genes, Novel Genes Involved in the Biosynthesis of Phytosiderophores. *Plant Physiology* **119**: 471-479
- Higuchi K, Watanabe S, Takahashi M, Kawasaki S, Nakanishi H, Nishizawa N-K, Mori S** (2001) Nicotianamine synthase gene expression differs in barley and rice under Fe-deficient conditions. *The Plant Journal* **25**(2): 159-167
- Joshi CP, Chang VL** (1998) Conserved sequence motifs in plant S-adenosyl-L-methionine-dependent methyltransferases. *Plant Mol Biol* **37**: 663-674

- Kawai S, Itoh K, Takagi S** (1993) Incorporation of ^{15}N and ^{14}C of methionine into the mugineic acid family of phytosiderophores in iron-deficient barley roots. *Physiologia Plantarum* **88**: 668-674
- Kawai S, Takagi S, Sato Y** (1988) Mugineic acid-family phytosiderophores in root-secretions of Barley, Corn and Sorghum varieties. *J Plant Nutrition* **11** (6-11): 633-642
- Kobayashi T, Nakanishi H, Takahashi M, Kawasaki S, Nishizawa NK, Mori S.** (2001) In vivo evidence that *Ids3* from *Hordeum vulgare* encodes a dioxygenase that converts 2'-deoxymugineic acid to mugineic acid in transgenic rice. *Planta* **212**: 864-871
- Korshunova YO, Eide D, Clark WG, Guerinot ML, Pakrasi HB** (1999) The IRT1 protein from *Arabidopsis thaliana* is a metal transporter with a broad substrate range. *Plant Mol Biol* **40**(1): 37-44
- Kristensen I, Larsen PO** (1974) Azetidine-2-carboxylic Acid Derivatives from Seeds of *Fagus Silvatica* L. and a Revised Structure for Nicotianamine. *Phytochemistry* **13**: 2791-2798
- Krüger C, Hell R, Stephan UW** (2001) A metal-binding LEA protein traffics micronutrients in the phloem of *Ricinus communis* L. in *Plant Nutrition* (Horst et al. Eds.) *Kluwer Academic Publishers* 54-55
- Kruger C, Berkowitz O, Stephan UW, Hell R** (2002) A metal-binding member of the late embryogenesis abundant protein family transports iron in the phloem of *Ricinus communis* L.. *J Biol Chem* **277** (28): 25062-25069
- Ling HQ, Koch G, Baumlein H, Ganai MW** (1999) Map-based cloning of *chloronerva*, a gene involved in iron uptake of higher plants encoding nicotianamine synthase. *Proc Natl Acad Sci USA* **96**(12):7098-7103
- Ling HQ, Bauer P, Berczky Z, Keller B Ganai MW** (2002) The tomato fer gene encoding a bHLH protein controls iron-uptake responses in roots. *Proc Natl Acad Sci USA*, published online before print **10.1073/pnas.212448699**
- Luna CM, Gonzalez CA, Trippi VS** (1994) Oxidative Damage Caused by an Excess of Copper in Oat Leaves. *Plant Cell Physiol* **35**(1): 11-15
- Ma JF, Nomoto K** (1993) Inhibition of mugineic acid-ferric complex uptake in barley by copper, zinc and cobalt. *Physiologia Plantarum* **89**: 331-334
- Ma JF, Nomoto K** (1996) Effective regulation of iron acquisition in gramineous plants. The role of mugineic acids as phytosiderophores. *Physiologia Plantarum* **97**: 609-617
- Mori S** (1999) Iron acquisition by plants. *Curr Opin Plant Biol* **2**(3): 250-3
- Mori S, Nishizawa N** (1987) Methionine as a Dominant Precursor of Phytosiderophores in Gramineae Plants. *Plant Cell Physiol* **28**(6): 1081-1092

- Nakanishi H, Bughio N, Matsushashi S, Ishioka N-S, Uchida H, Tsuji A, Osa A, Sekine T, Kume T, Mori S** (1999) Visualizing real time [¹¹C]methionine translocation in Fe-sufficient and Fe-deficient barley using a positron emitting tracer imaging system (PETIS). *J of Exp Botany* **50**: 637-643
- Nakanishi H, Okumura N, Umehhara Y, Nishizawa N-K, Chino M, Mori S** (1993) Expression of a Gene Specific for Iron Deficiency (Ids3) in the Roots of *Hordeum vulgare*. *Plant Cell Physiol* **34**(3): 401-410
- Nakanishi H, Yamaguchi H, Sasakuma T, Nishizawa N-K, Mori S** (2000) Two dioxygenase genes, Ids3 and Ids2, from *Hordeum vulgare* are involved in the biosynthesis of mugineic acid family phytosiderophores. *Plant Mol Biol* **44**: 199-207
- Noma M, Noguchi M, Tamaki E** (1971) A New Amino Acid, Nicotianamine, From Tobacco Leaves. *Tetrahedron Letters* **22**: 2017-2020
- Pich A, Manteuffel R, Hillmer S, Scholz G, Schmidt W** (2001) Fe homeostasis in plant cells: Does nicotianamine play multiple roles in the regulation of cytoplasmic Fe concentration? *Planta* **213**: 967-976
- Pich A, Scholz G** (1993) The relationship between the activity of various iron-containing and iron-free enzymes and the presence of nicotianamine in tomato seedlings. *Physiologia Plantarum* **88**: 172-178
- Pich A, Scholz G, Stephan UW** (1994) Iron dependant changes of heavy metals, nicotianamine and citrate in different plant organs and in xylem exudates of two tomato genotypes. Nicotianamine as possible copper translocator. *Plant Soil* **165**: 189-196
- Rabotti G, Zocchi G** (1994) Plasma-membrane bound H⁺-ATPase and reductase activities in Fe-deficient cucumber roots. *Physiologia Plantarum* **90**: 779-785.
- Ripperger H, Schreiber K** (1982) Nicotianamine and analogous amino acids, endogenous iron carriers in higher plants. *Heterocycles* **17**: 447-461
- Robinson NJ, Procter CM, Connolly EL, Guerniot ML** (1999) A ferric-chelate reductase for iron uptake from soils. *Nature* **329**: 694-697
- Römheld V and Marschner H** (1986) Mobilization of iron uptake by dicotyledonous and monocotyledonous plants. II. The reduction of Fe³⁺ as influenced by root and inhibitors. *J. Plant Nutr.* **2**: 647-660
- Rudolph A, Becker R, Scholz G, Prochazka Z** (1985) The occurrence of the amino acid nicotianamine in plants and microorganisms. A Reinvestigation. *Biochem Physiol Pflanzen* **180**: 557-563
- Samuelson AI, Martin RC, Mok DWS, Mok MC** (1998) Expression of yeast FRE genes in transgenic tobacco. *Plant Physiol* **118**: 51-58
- Schmidt W, Bartels M, Tittel J, Fuhner C** (1997) Physiological effects of copper on iron acquisition processes in *Plantago*. *New Phytologist* **135**: 659-666.

- Schmidt W, Bartels M, Tittel J, Fühner C** (1997) Physiological effects of copper on iron acquisition processes in *Plantago*. *New Phytol* **135**: 659-666
- Scholz G, Becker R, Pich A, Stephan UW** (1992) Nicotianamine - A common constituent of strategies I and II of iron acquisition by plants: A Review. *J of Plant Nutrition* **15**(10): 1647-1665
- Scholz G, Becker R, Stephan UW, Rudolph A, Pich A** (1988) The regulation of iron uptake and possible functions of nicotianamine in higher plants. *Biochem Physiol Pflanzen* **183**: 257-269
- Scholz G, Faust J, Ripperger H, Schreiber K** (1988) Structure-Funktion Relationships of Nicotianamine Analogues. *Phytochemistry* **27**(9): 2749-2754
- Scholz G, Rudolph A** (1968) A Biochemical mutant of *Lycopersicon esculentum* Mill. Isolation and properties of the ninhydrin-positive "Normalizing Factor". *Phytochemistry* **7**: 1759-1764
- Scrimshaw NS** (1986) Consequences of hunger for individuals and societies. *Fed Proc* **45** (10): 2421-6
- Shojima S , Nishizawa N-K, Fushiya S, Nozoe S, Irifune T, Mori S** (1990) Biosynthesis of Phytosiderophores. In vitro biosynthesis of 2'-deoxymugineic acid from L-methionine and nicotianamine. *Plant Physiol* **93**: 1497-1503
- Shojima S, Nishizawa N-K, Mori S** (1989) Establishment of a Cell-Free System for the Biosynthesis of Nicotianamine. *Plant Cell Physiol* **30**(5): 673-677
- Stephan UW, Grün M** (1989) Physiological disorders of the nicotianamine-auxotroph tomato mutant *chloronerva* at different levels of iron nutrition II. Iron deficiency response and heavy metal metabolism. *Biochem Physiol Pflanzen* **185**: 189-200
- Stephan UW, Scholz G** (1993) Nicotianamine: mediator of transport of iron and heavy metals in the phloem? *Physiologia Plantarum* **88**: 522-529
- Stephan UW, Scholz G, Rudolph A** (1990) Distribution of nicotianamine, a presumed symplast iron transporter, in different organs of sunflower and of a tomato wild type and its mutant *chloronerva*. *Biochem Physiol Pflanzen* **186**: 81-88
- Suzuki K, Higuchi K, Nakanishi H, Nishizawa H-K, Mori S** (1999) Cloning of nicotianamine synthase genes from *Arabidopsis thaliana*. *Soil Sci Plant Nutr* **45**(4): 993-1002
- Suzuki K, Shimada K, Nozoe S, Tanzawa K, Ogita T** (1996) Isolation of nicotianamine as a gelatinase inhibitor. *The Journal of Antibiotics* **49**(12): 1284-1285
- Vijayaraghavan K** (1995) Strategies for control of micronutrient malnutrition. *Indian J Med Res* **102**: 216-22

von Wirén N, Khodr H, Hider RC (2000) Hydroxylated phytosiderophore species possess and enhanced chelate stability and affinity for iron *Plant Physiology* **124**: 1149-1157

von Wirén N, Klair S, Bansal S, Briat J-F, Khodr H, Shioiri T, Leigh RA, Hider RC (1999) Nicotianamine chelates both FeIII and FeII. Implications for metal transport in plants. *Plant Physiology* **119**: 1107-1114

Wallace A, Wallace GA, Cha JW (1992) Some modifications in trace metal toxicities and deficiencies in plant resulting from interactions with other elements and chelating agents. The special case of iron. *J Plant Nutr* **15**: 1589-1598

Yeaman SY (1986) the mammalian 2-oxoacid dehydrogenases: a complex family. *Trends Biochem Sci* **11**: 283-296

Yi Y, Guerinot ML (1996) Genetic evidence that induction of root Fe(III) chelate reductase activity is necessary for iron uptake under iron deficiency. *The Plant Journal* **10**: 835–84