

5 Literatur

1. Albert ED, Scholz S: Juvenile arthritis: genetic update. *Baillieres Clin Rheumatol* 12 (1998) 209-218
2. Alsaeid KM, Haider MZ, al-Awadhi AM, Srivastva BS, Ayoub EM: Role of human leukocyte antigen DRB1*0307 and DRB1*0308 in susceptibility to juvenile rheumatoid arthritis. *Clin Exp Rheumatol* 21 (2003) 399-402
3. Afkarian M, Sedy JR, Yang J, Jacobson NG, Cereb N, Yang SY, Murphy TL, Murphy KM: T-bet is a STAT1-induced regulator of IL-12R expression on naive CD4+ T cells. *Nat Immunol* 3 (2002) 549-557
4. Akita K, Ohtsuki T, Nukada Y, Tanimoto T, Namba M, Okura T, Takakura-Yamamoto R, Torigoe K, Gu Y, Su MS, Fujii M, Satoh-Itoh M, Yamamoto K, Kohno K, Ikeda M, Kurimoto M: Involvement of caspase-1 and caspase-3 in the production and processing of mature human interleukin-18 in monocytic THP 1 cells. *J Biol Chem* 272 (1997) 26595-26603
5. al-Janadi M, al-Balla S, al-Dalaan A, Raziuddin S: Cytokine profile in systemic lupus erythematosus, rheumatoid arthritis and other rheumatic disease. *J Clin Immunol* 13 (1993) 58-67
6. Alphey L: DNA-Sequenzierung. Spektrum Akademischer Verlag GmbH, Heidelberg, Berlin, 1998, S.32
7. Amital H, Schoenfeld Y: Autoimmunity and autoimmune disease such as systemic lupus erythematosus. In: Lahita RG (Ed): *Systemic lupus erythematosus* Academic Press New York 1999, S.1-11
8. Arend WP, Dayer JM: Cytokines and cytokine inhibitors or antagonists in rheumatoid arthritis. *Arthritis Rheum* 33 (1990) 305-315

9. Block SR, Winfield JB, Lockshin MD, D'Angelo WA, Christian CL: Studies of twins with systemic lupus erythematosus: a review of the literature and presentation of 12 additional sets. *Am J Med* 59(1975) 533-552
10. Bombardier C, Gladman DD, Urowitz MB, Caron D, Chang CH: Derivation of the SLEDAI: a disease activity index for lupus patients. The Committee on Prognosis Studies in SLE. *Arthritis Rheum* 35 (1992) 630-640
11. Born TL, Thomassen E, Bird TA, Sims JE: Cloning a novel receptor subunit, AcPL, required for interleukin-18 signaling. *J Biol Chem* 273 (1998) 29445-29450
12. Botto M, Dell'Agnola C, Bygrave AE, Thompson EM, Cook HT, Petry F, Loos M, Pandolfi PP, Walport MJ: Homozygous C1q deficiency causes glomerulonephritis associated with multiple apoptotic bodies. *Nat Genet* 19 (1998) 56-59
13. Brunner HI, Silverman ED, To T, Bombardier C, Feldman BM: Risk factors for damage in childhood-onset systemic lupus erythematosus: cumulative disease activity and medication use predict disease damage. *Arthritis Rheum* 46 (2002) 436-444
14. Calvani N, Tucci M, Richards HB, Tartaglia P, Silvestris F: Th1 cytokines in the pathogenesis of lupus nephritis: the role of IL-18. *Autoimmunity Reviews* 4 (2005) 542-548
15. Cao Z, Xiong J, Takeuchi M, Kurama T, Goeddel DV: TRAF6 is a signal transducer for interleukin-1. *Nature* 383 (1996) 443-446
16. Cervera R, Khamashta MA, Font J, Sebastiani GD, Gil A, Lavilla P, Domenech I, Aydintug AO, Jedryka-Goral A, de Ramon E, et al.: Systemic lupus erythematosus: clinical and immunologic patterns of disease expression in a cohort of 1000 patients. The European Working Party on Systemic Lupus Erythematosus. *Medicine (Baltimore)* 72 (1993) 113-124

17. Chen DY, Lan JL, Lin FJ, Hsieh TY, Wen MC: Predominance of Th1 cytokine in peripheral blood and pathological tissues of patients with active untreated adult onset Still's disease. *Ann Rheum Dis* 63 (2004) 1300-1306
18. Clemens LE, Albert E, Ansell BM: HLA studies in IgM rheumatoid-factor-positive arthritis in childhood. *Ann Rheum Dis* 42 (1983) 431-434
19. Croston GE, Cao Z, Goeddel DV: NF- κ B activation by interleukin-1 (IL-1) requires an IL-1 receptor-associated protein kinase activity. *J Biol Chem* 270 (1995) 16514-16517
20. Date Y, Seki N, Kamizono S, Higuchi T, Hirata T, Miyata K, Ohkuni M, Tatsuzawa O, Yokota S, Joo K, Ueda K, Sasazuki T, Kimura A, Itoh K, Kato H: Identification of a genetic risk factor for systemic juvenile rheumatoid arthritis in the 5'-flanking region of the TNF α gene and HLA genes. *Arthritis Rheum* 42 (1999) 2577-2582
21. Davas EM, Tsirogianni A, Kappou I, Karamitsos D, Economidou I, Dantis PC: Serum IL-16, TNF- α , p55 sr TNF- α , p75 sr TNF- α , srIL-12 α levels and disease activity in systemic lupus erythematosus. *Clin Rheumatol* 18 (1999) 17-22
22. Davies EJ, Snowden N, Hillarby MC, Carthy D, Grennan DM, Thomson W, Ollier WE: Mannose-binding Protein gene polymorphism in systemic lupus erythematosus. *Arthritis Rheum* 38 (1995) 110-114
23. Deapen D, Escalante A, Weinrib L, Horwitz D, Bachmann B, Roy-Burman P, Walker A, Mack TM: A revised estimate of twin concordance in systemic lupus erythematosus. *Arthritis Rheum* 35 (1992) 311-318
24. De Benedetti F, Maezza C, Vivarelli M, Rossi F, Pistorio A, Lamb R, Lunt M, Thomson W, Ravelli A, Donn R, Martini A; British Paediatric Rheumatology Study Group: Functional and prognostic relevance of the -173 polymorphism of the macrophage migration inhibitory factor gene in systemic-onset juvenile idiopathic arthritis. *Arthritis Rheum* 48 (2003) 1398-1407

25. DiDonato JA, Hayakawa M, Rothwarf DM, Zandi E, Karin M: A cytokine-responsive I κ B kinase that activates the transcription factor NF- κ B. *Nature* 388 (1997) 548-554
26. Dinarello CA: IL-18: A TH1-inducing proinflammatory cytokine and new member of the IL-1 family. *J Allergy Clin Immunol* 103 (1999) 11-24
27. Dinarello CA, Moldawer LL: Proinflammatory and Anti-inflammatory Cytokines in Rheumatoid Arthritis. *Amgen*. 2002
28. Dolhain RJ, van der Heiden AN, ter Haar NT, Breedveld FC, Miltenburg AM: Shift toward T lymphocytes with a T helper 1 cytokine-secretion profile in the joints of patients with rheumatoid arthritis. *Arthritis Rheum* 39 (1996) 1961-1969
29. Dubois EL, Tufanelli DL: Clinical manifestations of systemic lupus erythematosus. Computer analysis of 520 cases. *JAMA* 190 (1964) 104.113
30. Esfandiari E, McInnes IB, Lindop G, Huang FP, Field M, Komai-Koma MK, Wei X, Liew FY: A proinflammatory role of IL-18 in the development of spontaneous autoimmune disease. *J Immunol* 167 (2001) 5338-5347
31. Eskdale J, Wordworth P, Bowman S, Field M, Gallagher G: Association between polymorphisms at the human IL-10 locus and systemic lupus erythematosus. *Tissue Antigens* 49 (1997) 635-639
32. Fernandez-Vina MA, Fink CW, Stastny P: HLA antigens in juvenile arthritis. Pauciarticular and polyarticular juvenile arthritis are immunogenetically distinct. *Arthritis Rheum* 33 (1990) 1787-1794
33. Font J, Cervera R, Espinosa G, Pallarés L, Ramos-Casals M, Jiménez S, Garcia-Carrasco M, Seisdedos L, Ingelmo M: Systemic lupus erythematosus (SLE) in childhood: analysis of clinical and immunological findings in 34 patients and comparison with SLE characteristic adults. *Ann Rheum Dis* 57 (1998) 456-459

34. Forre O, Dobloug JH, Hoyeraal HM, Thorsby E: HLA antigens in juvenile arthritis. Genetic bases for the different subtypes. *Arthritis Rheum* 26 (1983) 35-38
35. Fujikawa S, Okuni M: Clinical analysis of 570 cases with juvenile rheumatoid arthritis: results of a nationwide retrospective survey in Japan. *Acta Paediatr Jpn* 39 (1997) 245-249
36. Gattorno M, Facchetti P, Ghiotto F, Vignola S, Buoncompagni A, Prigione I, Picco P, Pistoia V: Synovial fluid T cell clones from oligoarticular juvenile arthritis patients display a prevalent Th1/Th0-type pattern of cytokine secretion irrespective of immunophenotype. *Clin Exp Immunol* 109 (1997) 4-11
37. Gracie JA: Interleukin-18 as a potential target in inflammatory arthritis. *Clin Exp Immunol* 136 (2004) 402-404
38. Gracie JA, Forsey RJ, Chan WL, Gilmour A, Leung BP, Greer MR, Kennedy K, Carter R, Wei XQ, Xu D, Field M, Foulis A, Liew FY, McInnes IB: A proinflammatory role for IL-18 in rheumatoid arthritis. *J Clin Invest* 104 (1999) 1393-1401
39. Gracie JA, Koyama N, Murdoch J, Field M, McGarry F, Crilly A, Schobel A, Madhok R, Pons-Kühnemann J, McInnes IB, Möller B: Disease association of two distinct interleukin-18 promoter polymorphisms in Caucasian rheumatoid arthritis patients. *Genes Immun* 6 (2005) 211-216
40. Gracie JA, Robertson SE, McInnes IB: Interleukin-18. *J Leukoc Biol* 73 (2003) 213-224
41. Gerdes N, Sukhova GK, Libby P, Reynolds RS, Young JL, Schonbeck U: Expression of interleukin (IL)-18 and functional IL-18 receptor on human vascular endothelial cells, smooth muscle cells, and macrophages: implications for atherogenesis. *J Exp Med* 195 (2002) 245-257

42. Gibofsky A, Winchester RJ, Patarroyo M, Fotino M, Kunkel HG: Disease association of the Ia-like human alloantigens. Contrasting patterns in rheumatoid arthritis and systemic lupus erythematosus. *J Exp Med* 148 (1978) 1728-1732
43. Giedraitis V, He B, Huang WX, Hillert J: Cloning and mutation analysis of the human Il-18 promotor: a possible role of polymorphism in expression regulation. *J Neuroimmunol* 112 (2001) 146-152
44. Grumet FC, Coukell A, Bodmer JG, Bodmer WF, McDevitt HO: Histocompatibility (HL-A) antigens associated with systemic lupus erythematosus. A possible genetic predisposition to disease. *N Engl J Med* 285 (1971) 193-196
45. Gu Y, Kuida K, Tsutsui H, Ku G, Hsiao K, Fleming MA, Hayashi N, Higashino K, Okamura H, Nakanishi K, Kurimoto M, Tanimoto T, Flavell RA, Sato V, Harding MW, Livingston DJ, Su MS: Activation of interferon- γ inducing factor mediated by interleukin-1 β converting enzyme. *Science* 275 (1997) 206-209
46. Haas JP, Nevinny-Stickel C, Schoenwald U, Truckenbrodt H, Suschke J, Albert ED: Susceptible and protective major histocompatibility complex class II alleles in early-onset pauciarticular juvenile chronic arthritis. *Hum Immunol* 41(1994) 225-233
47. Haus-Seuffert P, Meisterernst M: Mechanisms of transcriptional activation of cAMP-responsive element-binding-protein CREB. *Mol Cell Biochem* 212 (2000) 5-9
48. Heinzmann A, Gerhold K, Ganter K, Kurz T, Schuchmann L, Keitzer R, Berner R, Deichmann KA.: Association study of polymorphisms within interleukin-18 in juvenile idiopathic arthritis and bronchial asthma. *Allergy* 59 (2004) 845-849
49. Heninger E, Treszl A, Kocsis I, Derfalvi B, Tulassay T, Vasarhelyi B: Genetic variants of the interleukin-18 promotor region (-607) influence of course of necrotising enterocolitis in very low birth weight neonates. *Eur J Pediatr* 161 (2002) 410-411

50. Hess EV: Lupus- the Clinical Entity. In Lupus: Kammer GM, Tsokos GC (Eds): Molecular and cellular pathogenesis. Humana Press, Totowa, 1999, S 1-12
51. Higa S, Hirano T, Mayumi M, Hiraoka M, Ohshima Y, Nambu M, Yamaguchi E, Hizawa N, Kondo N, Matsui E, Katada Y, Miyatake A, Kawase I, Tanaka T: Association between interleukin-18 gene polymorphism 105 A/C and asthma. Clin Exp Immunol 33 (2003) 1097-1102
52. Ho CT, Mok CC, Lau CS, Wong RW: Late onset systemic lupus erythematosus in southern Chinese. Ann Rheum Dis 57 (1998) 437-440
53. Horneff G: Ätiopathogenese des systemischen Lupus erythematoses. Monatsschrift Kinderheilkunde 154 (2006) 1177-1188
54. Horwitz DA, Stohl W, Gray JD: T-lymphocytes, natural killer cells, cytokines, and immune regulation. In: Wallace DJ, Hahn BJ (Eds): Duboi's lupus erythematosus. Williams & Willkins, Baltimore 1997, S.155-194
55. Hoshino K, Tsutsui H, Kawai T, Takeda K, Nakanishi K, Takeda Y, Akira S: Cutting edge: generation of IL-18 receptor-deficient mice: evidence for IL-1 receptor-related protein as an essential IL-18 binding receptor. J Immunol 162 (1999) 5041-5044
56. Hyodo Y, Matsui K, Hayashi N, Tsutsui H, Kashiwamura S, Yamauchi H, Hiroishi K, Takeda K, Tagawa Y, Iwakura Y, Kayagaki N, Kurimoto M, Okamura H, Hada T, Yagita H, Akira S, Nakanishi K, Higashino K.: IL-18 up-regulates perforin-mediated NK activity without increasing perforin messenger RNA expression by binding to constitutively expressed IL-18 receptor. J Immunol 162 (1999) 1662-1668
57. Ide A, Kawasaki E, Abiru N, Sun F, Kobayashi M, Fukushima T, Takahashi R, Kuwahara H, Kita A, Oshima K, Uotani S, Yamasaki H, Yamaguchi Y, Eguchi K: Association between IL-18 gene promotor polymorphisms and CTLA-4 gene 49A/G polymorphism in Japanese patients with type 1 diabetes. J Autoimmunity 22 (2004) 73-78

58. Kawashima M, Yamamura M, Tani ai M, Yamauchi H, Tanimoto T, Kurimoto M, Miyawaki S, Amano T, Takeuchi T, Makino H: Levels of interleukin-18 and its binding inhibitors in the blood circulation of patients with adult-onset Still's disease. *Arthritis Rheum* 46 (2001) 2539-2541
59. Kjeldsen-Kragh J, Quayle A, Kalvenes C, Forre O, Sorskaar D, Vinje O, Thoen J, Natvig JB: T gamma delta cells in juvenile rheumatoid arthritis and rheumatoid arthritis. In the juvenile rheumatoid arthritis synovium the T gamma delta cells express activation antigens and are predominantly V delta 1+, and a significant proportion of these patients have elevated percentages of T gamma delta cells. *Scand J Immunol* 32 (1990) 651-659
60. Koene HR, Kleijer M, Swaak AJG, Sullivan E, Bijl M, Petri MA, Kallenberg CGM, Roos D, von dem Borne AEGK, de Haas M: The FcγRIIIa-158F allele is a risk factor for systemic lupus erythematosus. *Arthritis Rheum* 41 (1998) 1813-1818
61. Kretowski A, Mironczuk K, Karpinska A, Bojaryn U, Kinalski M, Puchalski Z, Kinalska I: Interleukin-18 promoter polymorphism in type 1 diabetes. *Diabetes* 51 (2002) 3347-3349
62. Kruse S, Kuehr J, Moseler M, Kopp MV, Kurz T, Deichmann KA, Foster PS, Mattes J: Polymorphisms in the IL 18 gene are associated with specific sensitization to common allergens and allergic rhinitis. *J Allergy Clin Immunol* 111 (2003) 117-122
63. Kuo CT, Leiden JM: Transcriptional regulation of T lymphocyte development and function. *Annu Rev Immunol* 17 (1999) 149-187
64. Lee WK, Song CH, Park YB, Lee CW, Suh CH, Lee CH, Lee J, Kim DS, Lee SK: Correlation of serum interleukin-8 with clinical activity in systemic lupus erythematosus (SLE). *Korean J Med* 54 (1998) 770-777
65. Leung BP, Culshaw S, Gracie JA, Hunter D, Canetti CA, Campbell C, Cunha F, Liew FY, McInnes IB.: A role for IL-18 in neutrophil activation. *J Immunol* 167 (2001) 2879-2886

66. Maeno N, Takei S, Nomura Y, Imanaka H, Hokonohara M, Miyata K: High elevated serum levels of interleukin-18 in systemic juvenile idiopathic arthritis but not in other juvenile idiopathic arthritis subtypes or in Kawasaki disease: comment on the article by Kawashima et al. *Arthritis Rheum* 46 (2002) 2539-2541; author reply 2541-2542
67. Malinin NL, Boldin MP, Kovalenko AV, Wallach D: MAP3K-related kinase involved in NK- κ B induction by TNF, CD95 and IL-1. *Nature* 385 (1997) 540-544
68. McMurray RW, May W: Sex hormones and systemic lupus erythematosus: review and meta-analysis. *Arthritis Rheum* 48 (2003) 2100-2110
69. McDowell TL, Symons JA, Ploski R, Forre O, Duff GW: A genetic association between juvenile rheumatoid arthritis and a novel interleukin-1 alpha polymorphism. *Arthritis Rheum* 38 (1995) 221-228
70. Miller ML, Aaron S, Jackson J, Fraser P, Cairns L, Hoch S, Borel Y, Larson M, Glass DN: HLA gene frequencies in children and adults with systemic onset juvenile rheumatoid arthritis. *Arthritis Rheum* 28 (1985) 146-150
71. Miltenburg AM, van Laar JM, de Kuiper R, Daha MR, Breedveld FC: T cells cloned from human rheumatoid synoviale membrane functionally represent the Th1 subset. *Scand J Immunol* 35 (1992) 603-610
72. Mitchell DA, Pickering MC, Warren J, Fossati-Jimack L, Cortes-Hernandez J, Cook HT, Botto M, Walport MJ: C1q deficiency and autoimmunity: the effects of genetic background on disease expression. *J Immunol* 168 (2002) 2538-2543
73. Mitterski B, Drynda S, Boschow G, Klein W, Oppermann J, Kekow J, Epplen JT: Complex genetic predisposition in adult and juvenile rheumatoid arthritis. *BMC Genet* 5 (2004) 1-14
74. Mok CC, Mak A, Chu WP, To CH, Wong SN: Long-term survival of southern chinese patients with systemic lupus erythematosus. A prospective study of all age-groups. *Medicine (Baltimore)* 84 (2005) 218-224

75. Monteleone G, Trapasso F, Parrello T, Biancone L, Stella A, Iuliano R, Luzzza F, Fusco A, Pallone F: Bioactive IL-18 expression is up-regulated in Crohn's disease. *J Immunol* 163 (1999) 143-147
76. Morel JC, Park CC, Zhu K, Kumar P, Ruth JH, Koch AE: Signal transduction pathways involved in rheumatoid arthritis synovial fibroblast interleukin-18-induced vascular cell adhesion molecule-1 expression. *J Biol Chem* 277 (2002) 34679-3491
77. Moroldo MB, Chaudhari M, Shear E, Thompson SD, Glass DN, Giannini EH: Juvenile rheumatoid arthritis affected sibpairs. *Arthritis & Rheumatism* 50 (2004) 1928-1934
78. Mosmann TR, Coffman RL: Th1 and th2 cells: different patterns of lymphokine secretion lead to different functional properties. *Annu Rev Immunol* 7 (1989) 145-173
79. Nakanishi K, Yoshimoto T, Tsutsui H, Okamura H: Interleukin-18 is a unique cytokine that stimulates both Th1 and Th2 responses depending on its cytokine milieu. *Cytokine Growth Factor Rev* 12 (2001) 53-72
80. Nakamura K, Okamura H, Wada M, Nagata K, Tamura T: Endotoxin-induced serum factor that stimulates gamma interferon production. *Infect Immun* 57 (1989) 590-595
81. Netea MG, Fantuzzi G, Kullberg BJ, Stuyt RJ, Pulido EJ, McIntyre RC Jr, Joosten LA, Van der Meer JW, Dinarello CA: Neutralization of IL-18 reduces neutrophil tissue accumulation and protects mice against lethal *Echerichia coli* and *Salmonella typhimurium* endotoxemia. *J Immunol* 164 (2000) 2644-2649
82. Newton CR, Graham A: PCR. 2. Aufl. Spektrum Akademischer Verlag GmbH, Heidelberg, Berlin, Oxford, 1994, S.28
83. Novak N, Kruse S, Potreck J, Maintz L, Jenneck C, Weidinger S, Fimmers R, Bieber T: Single nucleotide polymorphisms of the IL 18 gene are associated with atopic eczema. *J Allergy Clin Immunol* 115 (2005) 828-833

84. Oen K, Petty RE, Schroeder ML: An association between HLA-A2 and juvenile rheumatoid arthritis in girls. *J Rheumatol* 9 (1982) 916-920
85. Park MC, Park YB, Lee SK: Elevated interleukin-18 levels correlated with disease activity in systemic lupus erythematosus. *Clin Rheumatol* 23 (2004) 225-229
86. Park YB, Lee SK, Kim DS, Lee J, Lee CH, Song CH: Elevated IL-10 levels correlated with disease activity in systemic lupus erythematosus. *Clin Exp Rheumatol* 16 (1998) 283-288
87. Paul C, Yao Z, Nevinny-Stickel C, Keller E, Schoenwald U, Truckenbrodt H, Hoza J, Suschke HJ, Albert ED: Immunogenetics of juvenile chronic arthritis. HLA interactions between A2, DR5/8-DR/DQ, and DPB1*0201 is a general feature of all subsets of early onset pauciarticular juvenile chronic arthritis. *Tissue Antigens* 45 (1995) 280-283
88. Petty RE, Southwood TR, Manners P, Baum J, Glass DN, Goldenberg J, He X, Maldonado-Cocco J, Orozco-Alcala J, Prieur AM, Suarez-Almazor ME, Woo P; International League of Associations for Rheumatology.: International League of Association for Rheumatology classification of juvenile idiopathic arthritis: second revision, Edmonton, 2001. *J Rheumatol* 31 (2004) 390-392
89. Pickering MC, Walport MJ: Links between complement abnormalities and systemic lupus erythematosus. *Rheumatology (Oxford)* 39 (2000) 133-141
90. Pirhonen J: Regulation of IL-18 expression in virus infection. *Scand J Immunol* 53 (2001) 533-539
91. Ploski R: Immunogenetic polymorphism and disease mechanism in juvenile chronic arthritis. *Rev Rhum Engl Ed* 64 (1997) 127-130
92. Ploski R, Flato B, Vinje O, Maksymowych W, Forre O, Thorsby E: Association to HLA-DRB1*08, HLA-DPB1*0301 and homozygosity for an HLA-linked proteasome gene in juvenile ankylosing spondylitis. *Hum Immunol* 44 (1995) 88-96

93. Ploski R, Vinje O, Ronningen KS, Spurkland A, Sorskaar D, Vartdal F, Forre O: HLA class II alleles and heterogeneity of juvenile rheumatoid arthritis. DRB1*0101 may define a novel subset of disease. *Arthritis Rheum* 36 (1993) 465-472
94. Rachelefsky GS, Terasaki PI, Katz R, Stiehm ER: Increased prevalence of W27 in juvenile rheumatoid arthritis. *N Engl J Med* 290 (1974) 892-893
95. Ray A, Cohn L: Th2 cells and GATA-3 in asthma: new insights into the regulation of airway inflammation. *J Clin Invest* 104 (1999) 985-993
96. Reinertsen JL, Klippel JH, Johnson AH, Steinberg AD, Decker JL, Mann DL: B-lymphocyte alloantigens associated with systemic lupus erythematosus. *N Engl J Med* 299 (1978) 515-518
97. Rothe H, Hausmann A, Casteels K, Okamura H, Kurimoto M, Burkart V, Mathieu C, Kolb H: IL-18 inhibits diabetes development in nonobese diabetic mice by counterregulation of Th1-dependent destructive insulinitis. *J Immunol* 163 (1999) 1230-1236
98. Rueda B, González-Gay MÁ, Mataran L, López-Nevot MÁ, Martín J: Interleukin-18-promotor polymorphisms are not relevant in rheumatoid arthritis *Tissue Antigens* 65 (2005) 544-548
99. Schur PH, Meyer I, Garovoy M, Carpenter CB: Associations between systemic lupus erythematosus and the major histocompatibility complex: clinical and immunological considerations. *Clin Immunol Immunopathol* 24 (1982) 263-275
100. Seya T, Matsumoto M, Shiratori I, Fukumori Y, Toyoshima K: Protein polymorphism of human IL-18 identified by monoclonal antibodies. *Int J Mol Med* 8 (2001) 585-590
101. Shigehara K, Shijubo N, Ohmichi M, Yamada G, Takahashi R, Okamura H, Kurimoto M, Hiraga Y, Tatsuno T, Abe S, Sato N: Increased levels of interleukin-18 in patients with pulmonary sarcoidosis. *Am J Respir Crit Care Med* 162 (2000) 1979-1982

102. Sivalingam SP, Yoon KH, Koh DR, Fong KY: Single-nucleotide polymorphism of the interleukin-18 gene promoter region in rheumatoid arthritis patients: protective effect of AA genotype. *Tissue Antigens* 62 (2003) 498-504
103. Smeltz RB, Chen J, HU-LI J, Shevach EM: Regulation of interleukin (IL)-18 receptor alpha chain expression on CD4(+) T cells during T helper(Th)1/ Th2 differentiation. Critical downregulatory role of IL-4. *J Exp Med* 194 (2001) 143-153
104. Spronk PE, ter Borg EJ, Kallenberg CG: Plasma concentration of IL-6 in systemic lupus erythematosus: an indicator of disease activity? *Clin Exp Immunol* 90 (1992) 106-110
105. Sugiura T, Maeno N, Kawaguchi Y, Takei S, Imanaka H, Kawano Y, Terajima. Ichida H, Hara M, Kamatani N: A promoter haplotype of interleukin-18 gene is associated with juvenile idiopathic arthritis in the Japanese population. *Arthritis Res Ther* 8 (2006) 1-9
106. Takada T, Suzuki E, Morohashi K, Gejyo F: Association of single nucleotide polymorphism in IL-18 gene with sarcoidosis in a Japanese population. *Tissue Antigens* 60 (2002) 36-42
107. Tan EM, Cohen AS, Fries JF, Masi AT, McShane DJ, Rothfield NF, Schaller JG, Talal N, Winchester RJ: The 1982 revised criteria for the classification of systemic lupus erythematosus. *Arthritis and Rheumatism* 25 (1982) 1271-1277
108. Tokano Y, Morimoto S, Kaneko H, Amano H, Nozawa K, Takasaki Y, Hashimoto H: Levels of IL-12 in sera of patients with systemic lupus erythematosus (SLE)-relation to Th1 and Th2 derived cytokines. *Clin Exp Immunol* 116 (1999) 169-173
109. Tominaga K, Yoshimoto T, Torigoe K, Kurimoto M, Matsui K, Hada T, Okamura H, Nakanishi K: IL-12 synergizes with IL-18 or IL-1 beta for IFN-gamma production from human T cells. *Int Immunol* 12 (2000) 151-160

110. Tsao BP, Grossman JM, Riemekasten G, Strong N, Kalsi J, Wallace DJ, Chen CJ, Lau CS, Ginzler EM, Goldstein R, Kalunian KC, Harley JB, Arnett FC, Hahn BH, Cantor RM: Familiality and co-occurrence of clinical feature of systemic lupus erythematosus. *Arthritis & Rheum* 46 (2002) 2678-2685
111. Tsao BP, Cantor RM, Grossman JM, Shen N, Teophilov NT, Wallace DJ Arnett FC, Hartung K, Goldstein R, Kalunian KC, Hahn BH, Rotter JL: PARP alleles within the linked chromosomal region are associated with systemic lupus erythematosus. *J Clin Invest* 103 (1999) 1135-1140
112. van Kerckhove C, Luyrink L, Taylor J, Melin-Aldana H, Balakrishnan K, Maksymowych W, Elma M, Lovell D, Choi E, Glass DN: HLA-DQA1*0101 haplotypes and disease outcome in early onset pauciarticular juvenile rheumatoid arthritis. *J Rheumatol* 18 (1991) 874-879
113. Williams J, Ceccarelli A, Spurr N: Genetic Engineering. BIOS Scientific Publishers Limited, Oxford, 1993, S.2
114. Wong CK, Li EK, Ho CY, Lam CW: Elevation of plasma interleukin-18 concentration is correlated with disease activity in systemic lupus erythematosus. *Rheumatology (Oxford)* 39 (2000) 1078-1081
115. Xu D, Chan WL, Leung BP, Hunter D, Schulz K, Carter RW, McInnes IB, Robinson JH, Liew FY: Selective expression and functions of interleukin 18 receptor on T helper (Th) type 1 but not Th2 cells. *J Exp Med* 188 (1998) 1485-1492
116. Xu Q, Tin SK, Paramalingam SS, Thumboo J, Koh DR, Fong KY: Interleukin-18 promotor gene polymorphisms in Chinese patients with systemic lupus erythematosus: association with CC genotype at position -607. *Ann Acad Med Singapore* 36 (2007) 91-95

117. Yoshimoto T, Takeda K, Tanaka T, Ohkusu K, Kashiwamura S, Okamura H, Akira S, Nakanishi K: IL-12 up-regulates IL-18 receptor expression on T cells, Th1 cells, and B cells: synergism with IL-18 for IFN-gamma production. *J Immunol* 161 (1998) 3400-3407