

9 Literaturverzeichnis

- F. H. ALLEN, J. E. DAVIES, J. J. GALLOY, O. JOHNSON, O. KENNARD, C. F. MACRAE, E. M. MITCHELL, G. F. MITCHELL, J. M. SMITH, D. G. WATSON. The development of version 3 and 4 of the Cambridge Structural Database System. *J. Chem. Inf. Comp. Sci.*, **31**, 187-204 (1991)
- G. D. ANDRETTI, F. UGOZZOLI, A. CASNATI, E. GHIDINI, A. POCHINI, R. UNGARO. Molecular inclusion in functional macromolecules. XVII. Crystal and molecular structure of p-tert-butylcalix[6]arene 1:1 tetrachloroethylene clathrate. *Gazz. Chim. Ital*, **119**, 47-50 (1989)
- G. D. ANDRETTI, G. CALESTANI, F. UGOZZOLI, A. ARDUINI, E. GHIDINI, A. POCHINI, R. UNGARO. Solid state studies on p-tert-butylcalix[6]arene derivatives *J. Inclusion Phenom.*, **5**, 123-26 (1987)
- G. D. ANDRETTI, R. UNGARO, A. POCHINI. Crystal and molecular structure of cyclo{quater[(5-t-butyl-2-hydroxy-1,3-phenylene)methylene]} toluene (1:1) clathrate. *J. Chem. Soc., Chem. Commun.*, 1005-07 (1979)
- K. ARAKI, K. AKAO, H. OTSUKA, K. NAKASHIMA, F. INOKUCHI, S. SHINKAI. Immobilization of the ring inversion motion in calix[6]arene by a cap with C₃-symmetry. *Chem. Lett.*, 1251-54 (1994)
- L. ARCE, A. S. CARRETERO, A. RÍOS, C. CRUCES, A. FERNÁNDEZ, M. VALCÁRCEL. Use of calixarene compounds as selectivity modifiers in capillary electrophoresis separations. *J. Chromatogr. A*, **816**, 243-49 (1998)
- A. ARDUINI, A. POCHINI, S. REVERBERI, R. UNGARO, G. D. ANDRETTI, F. UGOZZOLI. The preparation of a new lipophilic sodium selective ether ester ligand derived from p-*t*-butylcalix[4]arene. *Tetrahedron*, **42**, 2098-2100 (1986)
- A. ARDUINI, D. BOLDRINI, A. POCHINI, A. SECCHI, R. UNGARO. Selective calix[4]arene-based sensors for neutral organic analytes. *Conf. Proc.-Ital. Phys. Soc.*, **54**, 131-37 (1997)
- G. ARENA, A. CASNATI, A. CONTINO, L. MIRONE, D. SCIOTTO, R. UNGARO. Synthesis of new calixcrowns and their anchoring to silica gel for selective separation of Cs⁺ and K⁺. *J. Chem. Soc., Chem. Commun.*, 2277-78 (1996)
- D. W. ARMSTRONG and W. DeMOND. Cyclodextrin bonded phases for the liquid chromatographic separation of optical, geometrical, and structural isomers. *J. Chromatogr. Sci.*, **22**, 411-15 (1984)
- D. W. ARMSTRONG, W. DeMOND, A. ALAK, W. L. HINZE, T. E. RIEHL, K. H. BUI. Liquid Chromatographic separation of diastereomers and structural isomers on Cyclodextrin-Bonded Phases. *Anal. Chem.*, **57**, 234-37 (1985)
- F. ARNAUD-NEU, E. M. COLLINS, M. DEASY, G. FERGUSON, S. J. HARRIS, B. KAITNER, A. J. LOUGH, M. A. McKERVEY, E. MARQUES, B. L. RUHL, M. J. SCHWING-WEILL, E. M. SEWARD. Synthesis, X-ray crystal structures, and cation-binding properties of alkyl calixaryl esters and ketones, a new family of macrocyclic molecular receptors. *J. Am. Chem. Soc.*, **111**, 8681-91 (1989)
- S. ASFARI and J. VICENS. The chemistry of calix[5]arenes. *Acros Org. Acta*, **1**, 18-21

(1995)

J. L. ATWOOD, D. L. CLARK, R. K. JUNEJA, G. W. ORR, K. D. ROBINSON, R. L. VINCENT. Double partial cone conformation for $\text{Na}_8\{\text{calix}[6]\text{arene sulfonate}\} \times 20,5 \text{ H}_2\text{O}$ and its parent acid. *J. Am. Chem. Soc.*, **114**, 7558-59 (1992)

J. L. ATWOOD, G. A. KOUTSANTONIS, C. L. RASTON, I. B. N. SUDRIA.. C_{60} - und C_{70} -Verbände in den zangenartigen Hohlräumen von Calix[6]aren. *Angew. Chemie*, **110**, 1029-31 (1998)

J. L. ATWOOD, G. A. KOUTSANTONIS, C. L. RASTON. Purification of C_{60} and C_{70} by selective complexation with Calixarenes. *Nature*, **368**, 229-31 (1994)

K. BÄCHMANN, A. BAZZANELLA, I. HAAG, K.-Y. HAN, R. ARNECKE, V. BÖHMER, W. VOGT. Resorcarenes as pseudostationary phases with selectivity for electrokinetic chromatography. *Anal.Lett.*, **67**, 1722-26 (1995)

E. P. BARRETT, L. G. JOYNER, P. P. HALENDA. The determination of pore volume and area distributions in porous substances. I. Computations from nitrogen isotherms. *J. Am. Chem. Soc.*, **73**, 373-80 (1951)

G. BARRETT, M. A. MCKERVEY, J. F. MALONE, A. WALKER, F. ARNAUD-NEU, M.-J. SCHWING-WEILL, C. D. GUTSCHE, D. R. STEWART. Cation complexation by chemically modified calixarenes. Part 6. Alkali and silver(I) cation complexation by *p*-*tert*-butylcalix[5]arene derivatives and X-ray structure determination of a pentaester. *J. Chem. Soc., Perkin Trans. 2*, 1475-78 (1993)

S. K. BATRA and J. SAUMANDE. High performance liquid chromatographic separation of estradiol- 17α and - 17β in biological fluids; application to plasma, milk and urine of cows. *J. Liq. Chromatogr.*, **7**(12), 2431-46 (1984)

A. BAZZANELLA, H. MÖRBEL, K. BÄCHMANN, R. MILBRADT, V. BÖHMER, W. VOGT. Highly efficient separation of amines by electrokinetic chromatography using resorcarene-octa-carboxylic acids as pseudostationary phases. *Biomed. Chromatogr.*, **12**, 111-12 (1998)

S. E. J. BELL, J. K. BROWNE, V. MCKEE, M. A. MCKERVEY, J. F. MALONE, M. O'LEARY, A. WALKER. Cation complexation by chemically modified calixarenes. 11. Complexation and extraction of alkali cations by calix[5]- and -[6]arene ketones. Crystal and molecular structures of calix[5]arene ketones and Na^+ and Rb^+ complexes. *J. Org. Chem.*, **63**, 489-501 (1998)

C. BOCCHI, M. CARERI, A. CASNATI, G. MORI. Selectivity of calix[4]arene-crown-6 for cesium ion in ISE: Effect of the conformation. *Anal. Chem.*, **67**, 4234-38 (1995)

V. BÖHMER. Calixarene - Makromoleküle mit (fast) unbegrenzten Möglichkeiten. *Angew. Chem.*, **107**, 785-818 (1995)

B. BOTTA, G. D. MONACHE, M. C. De ROSA, C. SERI, E. BENEDETTI, R. IACOVINO, M. BOTTA, F. CORELLI, V. MASIGNANI, A. TAFI, E. GACS-BAITZ, A. SANTINI, D. MISITI. Synthesis of C-alkylcalix[4]arenes. 5. Design, synthesis, computational studies, and homodimerization of polymethylene-bridged resorc[4]arenes. *J. Org. Chem.*, **62**, 1788-94 (1997)

B. BOTTA, G. D. MONACHE, P. SALVATORE, F. GASPARRINI, C. VILLANI, M. BOTTA, F. CORELLI, A. TAFI, E. GACS-BAITZ, A. SANTINI, C. F. CARVALHO, D. MISITI. Synthesis of C-alkylcalix[4]arenes. 4. Design, synthesis, computational

- studies of novel chiral amido[4]resorcinarenes. *J. Org. Chem.*, **62**, 932-38 (1997)
- W. BRANDT, O. LUDWIG, I. THONDORF, A. BARTH. A new mechanism in serine proteases catalysis exhibited by dipeptidyl peptidase IV (DPIV). Results of PM3 semiempirical thermodynamic studies supported by experimental results. *Europ. J. Biochem.*, **236**, 109-14 (1996)
- J. F. BRANDTS, H. R. HALVORSON, M. BRENNAN. Consideration of the possibility that slow step in protein denaturation is due to *cis-trans* isomerism of proline residues. *Biochemistry*, **14**, 4953-63 (1975)
- R. BRINDLE, K. ALBERT, S. J. HARRIS, C. TRÖLTZSCH, E. HORNE, J. D. GLENNON. Silica-bonded calixarenes in chromatography I. Synthesis and characterization by solid-state NMR spectroscopy. *J. Chromatogr. A*, **731**, 41-46 (1996)
- S. BRUNAUER, P. H. EMMETT, E. TELLER. Adsorption of gases in multimolecular layers. *J. Am. Chem. Soc.*, **60**, 309-19 (1938)
- A. CADOGAN, Z. GAO, A. LEWENSTAM, A. IVASKA, D. DIAMOND. All-solid-state sodium-selective electrode based on calixarene ionophore in a poly(vinyl chloride) membrane with a polypyrrole solid contact. *Anal. Chem.*, **64**, 2496-501 (1992)
- A. CASNATI, P. MINARI, A. POCHINI, R. UNGARO, W. F. NIJENHUIS, F. de JONG, D. N. REINHOUDT. Selective complexation and membrane transport of guanidinium salts by calix[6]arene amides. *J. Isr. Chem. Soc.*, **32**, 79-87 (1992)
- A. CASNATI, P. MINARI, A. POCHINI, R. UNGARO. Conformational freezing of *p*-*tert*-butylcalix[6]arene in the cone structure by selective functionalization at the lower rim: Synthesis of new preorganized ligands. *J. Chem. Soc., Chem. Commun.*, 1413-14 (1991)
- A. CASNATI. Calixarenes: From chemical curiosity to a rich source for molecular receptors. *Gazz. Chim. Ital.*, **127**, 637-49 (1997)
- S.-K. CHANG, H.-S. HWANG, H. SON, J. YOUK, Y. S. KANG. Selective transport of amino acid esters through a chloroform liquid membrane by a calix[6]arene-based ester carrier. *J. Chem. Soc., Chem. Commun.*, 217-18 (1991)
- S.-K. CHANG, I. CHO. New metal cation-selective ionophores derived from calixarenes: their syntheses and ion-binding properties. *J. Chem. Soc., Perkin Trans. 1*, 211-14 (1986)
- M. CORUZZI, G. D. ANDRETTI, V. BOCCHI, A. POCHINI, R. UNGARO. Molecular inclusion in functionalized macrocycles. Part 5: The crystal and molecular structure of 25,26,27,28,29-pentahydroxycalix[5]arene-acetone (1:2) clathrate. *J. Chem. Soc., Perkin Trans. 2*, 1133-38 (1982)
- J. COSTANTE-CRASSOUS, T. J. MARRONE, J. M. BRIGGS, J. A. McCAMMON, A. COLLET. Absolute configuration of bromochlorofluoromethane from molecular dynamics simulation of its enantioselective complexation by cryptophane-C. *J. Am. Chem. Soc.*, **119**, 3818-23 (1997)
- M. CZUGLER, S. TISZA, G. SPEIER. Versatility in inclusion hosts: unusual conformation in the crystal structure of the *p*-*tert*-butylcalix[8]arene:pyridine (1:8) clathrate. *J. Inclusion Phenom. Mol. Recognit. Chem.*, **11**, 323-31 (1991)

- E. DALCANALE, S. LEVI, S. RÖSLER, J. AUGÉ, J. HARTMANN, B. HENNING. Macromolecules for new chemical sensor applications. *GIT Lab. J.*, **2**, 118-21 (1998)
- G. DENG, T. SAKAKI, Y. KAWAHARA, S. SHINKAI. Tunable chemical sensors: light-switched ion selective electrodes on the basis of a photoresponsive calix[4]arene. *Supramolecular Chemistry*, **2**, 71-76 (1993)
- D. DIAMOND and M. A. McKERVEY. Calixarene-based sensing agents. *Chem. Soc. Rev.*, 15-24 (1996)
- H. ENGELHARDT and L. ROHRSCHEIDER. Nomenclature for Chromatography. *Pure und Appl. Chem.*, **65**, 819-72 (1993)
- C. A. EVANS, D. L. RABENSTEIN. Nuclear magnetic resonance studies of the acid-base chemistry of amino acids and peptides. II. Dependence of the acidity of the C-terminal carboxyl group on the conformation of the C-terminal peptide bond. *J. Am. Chem. Soc.*, **96**, 7312-16 (1974)
- E. FISCHER. Einfluss der Configuration auf die Wirkung der Enzyme. *Ber. Dtsch. Chem. Ges.*, **27**, 2984-93 (1894)
- G. FISCHER. Faltung der Proteine: Der Weg vom Baustein zur Funktion. in Verhandlungen der Gesellschaft Deutscher Naturforscher und Ärzte, 117. Versammlung, Wissenschaftliche Verlagsgesellschaft mbH, Aachen, 1992, 245-60
- G. FISCHER. Über Peptidyl-Prolyl-*cis/trans*-Isomerasen und ihre Effektoren. *Angew. Chem.*, **106**, 1479-1501 (1994)
- S. FRIEBE, B. HARTRODT, K. NEUBERT, G.-J. KRAUSS. HPLC separation of *cis-trans* isomers of proline-containing peptides. II. Fractionation of oligopeptides in different cyclodextrin systems. *J. Chromatogr.* **661**, 7-12 (1994)
- S. FRIEBE, G.-J. KRAUSS, H. NITSCHKE. HPLC separation of *cis-trans* isomers of proline-containing peptides. I. Separation on cyclodextrin bonded silica. *J. Chromatogr.* **598**, 139-42 (1992)
- S. FRIEBE, S. GEBAUER, G.-J. KRAUSS, G. GÖRMAR, J. KRÜGER. HPLC on calixarene bonded silica gels. I. Characterization and applications of *p-tert*-butylcalix[4]arene bonded materials. *J. Chromatogr. Sci.*, **33**, 281-84 (1995)
- C. A. FYFE. Solid state NMR for chemists. CFC-Press, Guelph (1983)
- P. A. GALE, Z. CHEN, M. G. B. DREW, J. A. HEATH, P. D. BEER. Lower-rim ferrocenyl substituted calixarenes: new electrochemical sensors for anions. *Polyhedron*, **17**, 405-12 (1998)
- J. GASTEIGER and M. MARSILI. Iterative partial equalization of orbital electronegativity - a rapid access to charges. *Tetrahedron*, **36**, 3219- 28 (1980)
- S. GEBAUER, Diplomarbeit: Synthese von Calix[4]aren-Derivaten zum Einsatz als selektive Trennmedien, MLU Halle-Wittenberg, 1993
- S. GEBAUER, S. FRIEBE, G. GÜBITZ, G.-J. KRAUSS. HPLC on calixarene bonded silica gels. II. Separation of regio- and stereoisomers on *p-tert*-butylcalix[n]arene phases. *J. Chromatogr. Sci.*, **36**, 383-87 (1998a)

- S. GEBAUER, S. FRIEBE, G. SCHERER, G. GÜBITZ, G.-J. KRAUSS. HPLC on calixarene bonded silica gels. III. Separation of *cis/trans* isomers of proline containing peptides. *J. Chromatogr. Sci.*, **36**, 388-94 (1998b)
- T. GILDENAST and J. LASCH. Isolation of ceramide fraction from human stratum corneum lipid extracts by high-performance liquid chromatography. *Biochimica et Biophysica Acta*, **1346**, 69-74 (1997)
- R. K. GILPIN, J. A. KORPL, C. A. JANICKI. In-situ chemically bonded stationary phases for high pressure liquid chromatography. *Anal. Chem.*, **47**, 1498-1502 (1975)
- J. D. GLENNON, B. LYNCH, K. HALL, S. J. HARRIS, P. O'SULLIVAN. Water-eluent based ion chromatography on silica bonded molecular baskets. *Spec. Publ. - R. Soc. Chem.*, 196(Progress in Ion Exchange), 153-59 (1997)
- J. D. GLENNON, E. HORNE, D. COCKER, A. KUHN, S. J. HARRIS, M. A. McKERVEY. Silica-bonded calixarenes in chromatography II. Chromatographic retention of metal ions and amino acid ester hydrochlorides. *J. Chromatogr. A*, **731**, 47-55 (1996)
- J. D. GLENNON, E. HORNE, K. O'CONNOR, G. KEARNEY, S. J. HARRIS, M. A. McKERVEY. Chromatographic selectivity for amino acid esters and alkali metal ions on a silica bonded calix[4]arene tetraester stationary phase. *Anal. Proc. Includ. Analy. Commun.*, **31**, 33-35 (1994)
- J. D. GLENNON, K. O'CONNOR, S. SRIJARANAI, K. MANLEY, S. J. HARRIS, M. A. McKERVEY. Enhanced chromatographic selectivity for Na⁺ ions on a calixarene-bonded silica phase. *Anal. Lett.*, **26**, 153-62 (1993)
- J. D. GLENNON, S. HUTCHINSON, S. J. HARRIS, A. WALKER, M. A. McKERVEY, C. C. McSWEENEY. Molecular baskets in supercritical CO₂. *Anal. Chem.*, **69**, 2207-12 (1997)
- G. GÖRMAR, K. SEIFFARTH, M. SCHULZ. Verfahren zur Herstellung von p-Alkylcalix[4]aren. Pat.-Nr.: DD-292904 vom 22.03.1990
- G. GRATHWOHL and K. WÜTHRICH. The X-Pro peptide bond as an NMR probe for conformational studies of flexible linear peptides. *Biopolymers*, **15**, 2025-41 (1976)
- S. J. GREGG and K. S. W. SING. Adsorption, surface area and porosity. (Second edition), Academic Press Inc., London, 1991
- C. D. GUTSCHE and L. J. BAUER. Calixarenes. 5. Dynamic and NMR characteristics of *p-tert*-butylcalix[4]arenes and *p-tert*-butylcalix[8]arenes. *Tetrahedron Lett.*, **22**, 4763-66 (1981b)
- C. D. GUTSCHE, A. E. GUTSCHE, A. I. KARAULOV. Calixarenes. 11. Crystal and molecular structure of *p-tert*-butylcalix[8]arene. *J. Inclusion Phenom.*, **3**, 447-51 (1985)
- C. D. GUTSCHE, B. DHAWAN, K. H. NO, and R. MUTHUKRISHNAN. Calixarenes. 4. The synthesis, characterization, and properties of the calixarenes from *p-tert*-butylphenol. *J. Am. Chem. Soc.*, **103**, 3782-92 (1981a)
- C. D. GUTSCHE. *Calixarenes - Revisited*, Publisher: Royal Soc. Chem., Letchworth, UK, 1998

- C. D. GUTSCHE. Calixarenes. *Acc. Chem. Res.*, **16**, 161-70 (1983)
- C. D. GUTSCHE. *Calixarenes; Monographs in Supramolecular Chemistry*, ed. J. F. STODDARD, The Royal Society of Chemistry, Cambridge, 1989
- C. D. GUTSCHE: Calixarenes. *Aldrichimica Acta*, **28**, 3-9 (1995)
- T. HAINO, M. YANASE, Y. FUKAZAWA. Crystalline complexes of C₆₀ with calix[5]arenes. *Tetrahedron Lett.*, **38**, 3739-42 (1997)
- M. HALIT, D. OEHLER, M. PERRIN, A. THOZET, R. PERRIN, J. VIECENS, M. BOURAKHOUDAR. Crystal and molecular structure of two calix[6]arenes; *p*-isopropyl-calix[6]arene and *p*-*tert*-butylcalix[6]arene-benzene (1:3) complex. *J. Inclusion Phenom.*, **6**, 613-23 (1988)
- T. HARADA and S. SHINKAI. Combined NMR spectroscopy and molecular mechanics studies on the stable structure of calix[n]arenes. *J. Chem. Soc., Perkin Trans. 2*, 2231-42 (1995)
- L. O. HEALY, M. M. McENERY, D. G. McCARTHY, S. J. HARRIS, J. D. GLENNON. Silica-bonded calixarenes in chromatography: enantioseparations on molecular basket phases for rapid chiral LC. *Anal. Lett.*, **31** (9), 1543-51 (1998)
- D. E. HENDERSON and C. HORVÁTH. Low temperature high-performance liquid chromatography of *cis-trans* proline dipeptides. *J. Chromatogr.*, **368**, 203-13 (1986)
- A. HIRSCH. Die Chemie der Fullerene. *Chemie in unserer Zeit*, **2**, 79-87 (1994)
- C. HORVÁTH, W. MELANDER, I. MOLNÁR. Solvophobic interaction in liquid chromatography with nonpolar stationary phases. *J. Chromatogr.*, **125**, 129-156 (1976)
- D. HÜBNER, T. DRAKENBERG, S. FORSEN, G. FISCHER. Peptidyl-prolyl *cis-trans* isomerase activity as studied by dynamic proton NMR spectroscopy. *FEBS Lett.*, **284**, 79-81 (1991)
- A. IKEDA, M. YOSHIMURA, S. SHINKAI. Solution complexes formed by C₆₀ and calixarenes. On the Importance of the preorganized structure for cooperative interaction. *Tetrahedron Lett.*, **38**, 2107-10 (1997)
- N. IKI, F. NARUMI, T. SUZUKI, A. SUGAWARA, S. MIYANO. A new chiral stationary phase for gas chromatography by use of a chiral thiacalix[4]arene derivative. *Chem. Lett.*, 1065-66 (1998)
- N. S. ISAACS, P. J. NICHOLS, C. L. RASTON, C. A. SANDOVA, D. J. YOUNG. Solution volume studies of a deep cavity inclusion complex of C₆₀:[*p*-benzylcalix[5]arene ⊂ C₆₀]. *J. Chem. Soc., Chem. Commun.*, 1839-40 (1997)
- K. IWAMOTO, H. SHIMIZU, K. ARAKI, S. SHINKAI. Syntheses and optical resolution of calix[4]arenes with molecular asymmetry. Systematic classification of all possible chiral isomers derivable from calix[4]arene. *J. Am. Chem. Soc.*, 3997-4006 (1993)
- K. IWATA, H. KIMIZUKA, H. SUZUKI. Adsorptionsträger für Chromatographie. *Japan Patent Hei 6-58920* (1994)
- K. IWATA, S. MORIGUCHI. Adsorptionsträger für Chromatographie. *Japan Patent*

Hei 5-264531 (1993)

R. G. JANSSEN, J. P. M. van DUYNHOVEN, W. VERBOOM, G. J. van HUMMEL, S. HARKEMA, D. N. REINHOUDT. Studies on the dynamics of phosphorylated *p*-tert-butylcalix[6]arenes by using 2D NMR spectroscopy. *J. Am. Chem. Soc.*, **118**, 3666-75 (1996)

K. JINNO, K. YAMAMOTO, T. UEDA, H. NAGASHIMA, K. ITOH, J. C. FETZER, W. R. BIGGS. Liquid chromatographic separation of all-carbon molecules C₆₀ and C₇₀ with multi-legged phenyl group bonded silica phases. *J. Chromatogr.*, **594**, 105-09 (1992)

K. JINNO, T. UEMURA, H. OHTA, H. NAGASHIMA, K. ITOH. Separation and identification of higher molecular weight fullerenes by high-performance liquid chromatography with monomeric and polymeric octadecylsilica bonded phases. *Anal. Chem.*, **65**, 2650-54 (1993 b)

K. JINNO, Y. SAITO, Y.-L. CHEN, G. LUEHR, J. ARCHER, J. C. FETZER, W. R. BIGGS. Separation of C₆₀ and C₇₀ fullerenes on methoxyphenylpropyl bonded stationary phases in microcolumn liquid chromatography. *J. Microcolumn Sep.*, **5**, 135-40 (1993 a)

R. K. JUNEJA, K. D. ROBINSON, G. W. ORR, R. H. DUBOIS, K. A. BELMORE, J. L. ATWOOD, J. A. RIPMEESTER, C. I. RATCLIFFE. Inclusion of multi-ring compounds by *p*-tert-butylcalix[5]arene. *J. Inclusion Phenom. Mol. Recognit. Chem.*, **13**, 93-96 (1992)

A. KÁLMÁN, F. THUNECKE, R. SCHMIDT, P. W. SCHILLER, and C. HORVÁTH. Isolation and identification of peptide conformers by RP-HPLC and NMR at low temperature. *J. Chromatogr. A*, **729**, 155-71 (1996)

S. KANAMATHAREDDY and C. D. GUTSCHE. Synthesis and conformational properties of calix[6]arenes bridged on the lower rim: self-anchored rotaxanes. *J. Am. Chem. Soc.*, **115**, 6572-79 (1993)

Y. R. KANG, H. J. OH, K. M. LEE, G. S. CHA, H. NAM, K. PAEK, H. IHM. Potentiometric characteristics of ion-selective electrodes based on upper-rim calix[4]crown neutral carrier. *Bull. Korean Chem. Soc.*, **19**, 207-11 (1998)

T. KATSU. Determination of the pH difference across a cell membrane using a methylammonium-selective membrane electrode. *Analyst*, **123**, 1369-72 (1998)

H. KAWAURA, T. SAKAMOTO, T. BABA, Y. OCHIAI, J. FUJITA, S. MATSUI, J. SONE. Transistor operation of 30-nm gate-length EJ-MOSFETs. *IEEE Electron Device Lett.*, **19**, 74-76 (1998)

W. KEMIN, H. XIQIN, C. W. HOUNG, L. A. W. MING. Development of a dopamine-selective optrode membrane based on a lipophilic hexaester of calix[6]arene. *J. of Hunan University*, **22**, 55-60 (1995)

M. KRAMER and G. FISCHER. FKBP-like catalysis of peptidyl-prolyl bond isomerization by micelles and membranes. *Biopolymers*, **42**, 49-60 (1997)

G.-J. KRAUSS und G. KRAUSS. SERVA Praxishefte. Band 2 Hochleistungsflüssigchromatographie, Heidelberg, New York, 1986

- G.-J. KRAUSS. Dissertation für die Promotion B: Beiträge zur hochleistungs-chromatographischen Analytik von Nukleobasen, Nukleosiden und Nukleotiden in biologischen Systemen, MLU Halle-Wittenberg, 1987
- H. KUMAGAI, S. MIYANARI, S. MIYANO. Preparation of sulfonylcalixarene and sulfanylcalixarene derivatives as ion sensors. *PCT Int. Appl.*, 24 pp. (1996)
- M. A. LAMPE, A. L. BURLINGAME, J. WHITNEY, M. L. WILLIAMS, B. E. BROWN, E. ROITMAN, P. M. ELIAS. Human stratum corneum lipids: characterization and regional variations. *J. Lipid Res.*, **24**, 120-30 (1983)
- B. T. T. LAN, K. TOTH. Characterization of chromogenic calix[4]arene derivative based ion-selective optical sensors. *Anal. Sci.*, **14**, 191-97 (1998)
- F. LARA-OCHOA, J. A. COGORDAN, I. SILAGHI-DUMITRESCU. Interaction between calix[8]arenes and fullerenes. Molecular dynamics and molecular mechanics simulation of the 1:1 and 1:2 complexes. *Fullerene Science and Technology*, **4**(5), 887-96 (1996)
- H. J. LEE, H. J. OH, G. CUI, G. S. CHA, H. NAM. All-solid-state sodium-selective electrodes based on room temperature vulcanizing-type silicone rubber matrix . *Anal. Sci.*, **13**, 289-94 (1997)
- Y. L. LEE, Y. K. RYU, J. W. RYU, B. E. KIM, J. H. PARK. Reversed-phase liquid chromatography of some positional isomers on calix[6]arene-*p*-sulfonate-bonded silica. *Chromatographia*, **46**(9/10), 507-10 (1997)
- H. J. LIM, H. S. LEE, I. W. KIM, S. H. CHANG, S. C. MOON, B. E. KIM, J. H. PARK. Separation of positional isomers on polymeric calix[4]arene-siloxane stationary phases in capillary GC. *Chromatographia*, **48**(5/6), 422-26 (1998)
- L. LIN, C. Y. WU, Z. Q. YAN, X. Q. YAN, X. L. SU, H. M. HAN. Calix[4]arene derivative as stationary phases for capillary gas chromatography. *Chromatographia*, **47** (11/12), 689-94 (1998a)
- L. LIN, Z. YAN, X. SU, C. WU. Gas chromatographic study on the inclusion properties of calix[4]arene derivatives. *Sepu*, **16**(3), 208-10 (1998b)
- K. B. LIPKOWITZ, G. PEARL, B. CONER, M. A. PETERSON. Explanation of where and how enantioselective binding takes place on permethylated β -cyclodextrin, a chiral stationary phase used in gas chromatography. *J. Am. Chem. Soc.*, **119**, 600-10 (1997)
- K. B. LIPKOWITZ, R. CONER, M. A. PETERSON. Locating of maximum chiral discrimination: A computational study of enantioselection on a popular chiral stationary phase used in chromatography. *J. Am. Chem. Soc.*, **119**, 11269-76 (1997)
- S. A. LONG, P.W. WERTZ, J. S. STRAUSS, D. T. DOWNING. Human stratum corneum polar lipids and desquamation. *Arch. Derm. Res.*, **277**, 284-87 (1985)
- U. LÜNING, H. ROSS, I. THONDORF. Concave reagents. Part 27. Steric requirements of intraannular substituents in A,D-bridged calix[6]arenes. *J. Chem. Soc., Perkin Trans. 2*, 1313-17 (1998)
- S. MA, F. KÁLMÁN, A. KÁLMÁN, F. THUNECKE, C. HORVÁTH. Capillary zone electrophoresis at subzero temperatures I. Separation of the *cis* and *trans* conformers of small peptides. *J. Chromatogr. A*, **716**, 167-82 (1995)

- G. E. MACIEL and D. W. SINDORF. Silicon-29 nuclear magnetic resonance study of the surface of silica gel by cross polarization and magic-angle spinning. *J. Am. Chem. Soc.*, **102**, 7606-7 (1980)
- J. O. MAGRANS, A. M. RINCÓN, F. CUEVAS, J. LÓPEZ-PRADOS, P. M. NIETO, M. PONS, P. PRADOS, J. de MENDOZA. Interplay for steric hindrance and hydrogen bonding to restrict mono-*O*-substituted *p*-*tert*-butylcalix[6]arenes in cone conformation. *J. Org. Chem.*, **63**, 1079-85 (1998)
- A. MANGIA, A. POCHINI, R. UNGARO, G. D. ANDREETTI. 4-*tert*-butylcalix[8]-arene as a stationary phase in gas-solid chromatography. *Anal. Chem.*, **16**, 1027-36 (1983)
- D. E. MARTIRE and R. BOEHM. Unified theory of retention and selectivity in liquid chromatography. 2. Reversed-phase liquid chromatography with chemically bonded phases. *J. Phys. Chem.*, **87**, 1045-62 (1983)
- M. A. MCKERVEY, E. M. SEWARD, G. FERGUSON, B. L. RUHL, S. J. HARRIS. Synthesis. X-ray crystal structures, and cation transfer properties of alkyl calixaryl acetates, a new series of molecular receptors. *J. Chem. Soc., Chem. Commun.*, 388-90 (1985)
- W. R. MELANDER, J. JACOBSON, C. HORVÁTH. Effect of molecular structure and conformational change of proline-containing dipeptides in reversed-phase chromatography. *J. Chromatogr.*, **234**, 269-76 (1982)
- U. MENYES, U. ROTH, C. TRÖLTZSCH. Process for modifying supports containing silanol groups with calixarene for chromatography. *Eur. Pat. Appl.*, 1996
- S. MEYER, A. JABS, M. SCHUTKOWSKY, G. FISCHER. Separation of *cis/trans* isomers of a prolyl peptide bond by capillary zone electrophoresis. *Electrophoresis*, **15**, 1151-57 (1994)
- V. MEYER, *Praxis der Hochleistungsflüssigchromatographie, Reihe: Laborbücher Chemie*, Salle und Sauerländer, Frankfurt a. M., Aarau, 1990
- J. S. MILLERSHIP, M. A. MCKERVEY, J. A. RUSSELL. Complexation and chromatographic studies involving water-soluble calixarenes and nitrophenols. *Chromatographia*, **48**(5/6), 402-06 (1998)
- S. MIYAMOTO and P. A. KOLLMAN. Molecular dynamics studies of calixspherand complexes with alkali metal cations: Calculation of the absolute and relative free energies of binding of cations to a calixspherand. *J. Am. Chem. Soc.*, **114**, 3668-74 (1992)
- R. MLIKA, H. B. OUADA, N. JAFFREZIC-RENAULT, I. DUMAZET, R. LAMARTINE, M. GAMOUDI, G. GUILLAND. Study of ion-selective evaporated calixarene film used as sensitive layer on ISFET. *Sens. Actuators, B*, **B47**, 43-47 (1998)
- R. MLIKA, H. B. OUADA, R. BEN CHAABANE, M. GAMOUDI, G. GUILLAUD, N. JAFFREZIC-RENAULT, R. LAMARTINE. Calixarene membranes on semiconductor substrates for EIS chemical sensors. *Electrochim. Acta*, **43**, 842-47 (1997b)

- R. MLIKA, I. DUMAZET, M. GAMOUDI, R. LAMARTINE, H. BEN OUADA, N. JAFFREZIC-RENAULT, G. GUILLAUD. Membranes containing new large size calixarenes on semiconductor substrates for chemical microsensors. *Anal. Chim. Acta*, **354**, 283-89 (1997a)
- P. MNUK and L. FELTL. Gas chromatographic study of the inclusion properties of calixarenes I. *p*-*tert*-butylcalix[4]arene in a micropacked column. *J. Chromatogr. A*, **696**, 101-12 (1995)
- P. MNUK, L. FELTL, V. SCHURIG. Gas chromatographic study of the inclusion properties of calixarenes. II. Selective properties of cyclic tetra to octamers derived from phenol, and some problems associates with the use of calixarenes in capillary gas chromatography. *J. Chromatogr. A*, **732**, 63-74 (1996)
- M. A. MOLINS, P. M. NIETO, C. SÁNCHEZ, P. PRADOS, J. de MENDOZA, M. PONS. Solution structure and conformational equilibria of a symmetrical calix[6]arene. Complete sequential and cyclostereospecific assignment of the low-temperature NMR spectra of a cyclosymmetric molecule. *J. Org. Chem.*, **57**, 6924-31 (1992)
- N. MUZET, G. WIPFF, A. CASNATI, L. DOMIANO, R. UNGARO, F. UGOZZOLI. Alkaline earth and uranyl cation complexes of a calix[4]arene-tetraamide: MD and FEP simulations in aqueous and acetonitrile solutions and X-ray structure of its Sr(Picrate)₂ complex. *J. Chem. Soc., Perkin Trans. 2*, 1065-75 (1996)
- M. NELSON, W. HUMPHREY, A. GURSOY, A. DALKE, L. KALE, R. SKEEL, K. SCHULTEN, R. KUFRIN. MDScope – a visual computing environment for structural biology. *Comp. Phys. Comm.*, **91**, 111-33 (1995)
- M. NEMILOVA, N. V. SHVEDENE, O. V. KOCHULINA, I. A. PRONINA, I. V. PLETNEY. PVC-plasticized ion-selective electrodes based on neutral carriers for determining novocaine. *Vestn. Mosk. Univ., Ser. 2: Khim.*, **38**, 99-103 (1998)
- P. NERI, E. BATTOCOLO, F. CUNSOLO, C. GERACI, M. PIATTELLI. Study of *p*-*tert*-butylcalix[8]arene. Partially *O*-alkylated calix[8]arenes. *J. Org. Chem.*, **59**, 3880-89 (1994)
- K. M. O'CONNER, W. HENDERSON, E. O'NEILL, W. M. D. ARRIGAN, S. J. HARRIS, M. A. McKERVEY, G. SVEHLA. Calix[4]arenes in the partial cone conformation as ionophores in silver ion-selective electrodes. *Electroanalysis*, **9**, 311-15 (1997)
- K. OHTO, Y. TANAKA, K. INOUE. Adsorption of lead and zinc ions by novel type of calix[4]arene carboxylate resin immobilized with polyallylamine. *Chem. Lett.*, 647-48 (1997)
- S. PAPPALARDO and M. F. PARISI. Selective *endo*-calix complexation of linear alkylammonium cations by functionalized (1,3)-*p*-*tert*-butylcalix[5]crown ethers. *J. Org. Chem.*, **61**, 8724-25 (1996)
- C. PARINI, S. COLOMBO, A. CASNATI. Solid state interaction of steroids with calixarenes. I. A preliminary FTIR and DSC study on 4-en-3-keto-steroids. *J. Inclusion Phenom. Mol. Recognit. Chem.*, **18**, 341-51 (1994)
- J. H. PARK, Y. K. LEE, N. Y. CHEONG, M. D. JANG. Reversed phase liquid chromatographic separation of some mono-substituted phenols with calix[6]arene-*p*-sulfonate-modified eluents. *Chromatographia*, **37**, 221- 23 (1993)

- R. PATHAK and G. N. RAO. Synthesis and metal sorption studies of p-tert-butylcalix[8]arene chemically bound to polymeric support. *Anal. Chim. Acta*, **335**, 283-90 (1996)
- D. A. PEARLMAN, D. A. CASE, J. W. CALDWELL, W. S. ROSS, T. E. CHEATHAM III, D. M. FERGUSON, G. L. SEIBEL, U. C. SINGH, P. K. WEINER and P. A. KOLLMAN, AMBER 4.1, University of California, San Francisco, (1995)
- M. S. PENA, Y. ZHANG, I. M. WARNER. Enantiomeric separation by use of calixarene electrokinetic chromatography. *Anal. Chem.*, **69**, 3239-42 (1997)
- M. S. PENA, Y. ZHANG, S. THIBODEAUX, M. L. McLAUGHLIN, A. M. PENA, I. M. WARNER. Synthesis of a water-soluble chiral *N*-acylcalix[4]arene amino acid derivative. *Tetrahedron Lett.*, **37**, 5841-44 (1996)
- M. PERRIN and S. LECOCQ. Crystal and molecular structure of p-(1,1,3,3-tetramethylbutyl)calix[5]arene and its 1:1 complex with toluene. *J. Inclusion Phenom. Mol. Recognit. Chem.*, **11**, 171-83 (1991)
- R. J. PIETERS, J. CUNTZE, M. BONNET, F. DIEDERICH. Enantioselective recognition with C₃-symmetric cage-like receptors in solution and on a stationary phase. *J. Chem. Soc., Perkin Trans. 2*, 1891-1900 (1997)
- A. PINES, M. G. GIBBY, J. S. WAUGH. Proton-enhanced NMR of dilute spins in solids. *J. Phys. Chem.*, **59**, 569-90 (1973)
- A. POCHINI and R. UNGARO. Calixarenes and related hosts. In: *Comprehensive Supramolecular Chemistry*, eds. J. L. ATWOOD, J. E. D. DAVIES, M. ACNICOL, F. VÖGTLE, Elsevier Science Ltd., Oxford, Vol. **2**, 143ff (1996)
- B. PULLMAN (Hrsg.), *Intermolecular Forces*, Reidel, Dordrecht, 1991
- I. RODRIGUEZ, H. K. LEE, S. F. Y. LI. Capillary electrophoresis separation of *p*-sulfonated calix[*n*]arenes, *n*=4,6,8. *TALANTA*, **45**, 683-691 (1998)
- M. K. ROSEN, R. F. STANDAERT, A. GALAT, M. NAKATSUKA, S. L. SCHREIBER. Inhibition of FKBP rotamase activity by immunosuppressant FK506: twisted amide surrogate. *Science*, **248**, 863- 66 (1990)
- S. RÖSLER, R. LUCKLUM, R. BORNGRABER, J. HARTMANN, P. HAUPTMANN. Sensor system for the detection of organic pollutants in water by thickness shear mode resonators. *Sens. Actuators, B*, **B48**, 415-24 (1998)
- J. P. RYCKAERT, G. CICCOTTI, H. J. C. BERENDSEN. Numerical integration of the Cartesian equations of motion of a system with constraints: molecular dynamics of *n*-alkanes. *J. Comput. Phys.*, **23**(3), 327-41, (1977)
- T. SAIKI, K. GOTO, N. TOKITOH, M. GOTO, R. OKAZAKI. Structure of bridged calix[6]arenes bearing a bromide functionality in the cavity. *Tetrahedron Lett.*, **37**, 4039-42 (1996)
- K. D. SCHIERBAUM and W. GÖPEL. Functional polymers and supramolecular compounds for chemical sensors. *Synthetic Metals*, **61**, 37-45 (1993)
- H.-J. SCHNEIDER, R. KRAMER, I. THEIS, M.-Q. ZHOU. Large extra-cavity contributions to hydrophobic binding of lipophilic substrates in synthetic receptor

models. *J. Chem. Soc., Chem. Commun.*, 276-78 (1990b)

H.-J. SCHNEIDER, T. BLATTER, P. ZIMMERMANN. Wirt-Gast-Komplexe mit geschlossenen, halboffenen und gestreckten Rezeptoren: hydrophobe Effekte und induzierte Pol-Dipol-Wechselwirkungen. *Angew. Chemie*, **102**, 1194-95 (1990a)

H.-J. SCHNEIDER. Mechanismen in der molekularen Erkennung- Untersuchungen an organischen Wirt-Gast-Komplexen. *Angew. Chemie*, **103**, 1419-548 (1991)

A. B. SCHOLTEN, J. W. de HAAN, H. A. CLAESSENS, L. J. M. van de VEN, C. A. CRAMERS. 29-Silicon NMR evidence for the improved chromatographic siloxane bond stability of bulky alkylsilane ligands on a silica gel surface. *J. Chromatogr. A*, **688**, 25-29 (1994)

M.-J. SCHWING-WEILL and F. ARNAUD-NEU. Calixarenes for radioactive waste management. *Gazz. Chim. Ital.*, **127**, 687-92 (1998)

T. A. SERGEYEVA, N. V. LAVRIK, A. E. RACHKOV, Z. I. KAZANTSEVA, A. V. ELSKYA. An approach to conductometric immunosensor based on phthalocyanine thin film. *Biosens. Bioelectron.*, **13**, 359-69 (1998)

S. SHINKAI, H. KAWABATA, T. MATSUDA, H. KAWAGUCHI, O. MANABE. Synthesis and inclusion properties of „neutral“ water-soluble calixarenes. *Bull. Chem. Soc. Jpn.*, **63**, 1272-74 (1990)

S. SHINKAI, K. ARAKI, O. MANABE. NMR determination of association constants for calixarenes. Evidence for the formation of 1:2 complex with calix[8]arene. *J. Am. Chem. Soc.*, **110**, 7214-15 (1988)

D. SHOHAT and E. GRUSHKA. Use of calixarenes to modify selectivities in capillary electrophoresis. *Anal. Chem.*, **66**, 747-50 (1994)

R. C. SIMPSON, P. R. BROWN. High performance liquid chromatographic profiling of nucleic acid components in physiological samples. *J. Chromatogr., Biomed. Appl.*, **379**, 269-311 (1986)

D. W. SINDORF and G. E. MACIEL. ¹³C CP/MAS NMR study of molecular motion in *n*-alkylsilanes bonded to the silica surface. *J. Am. Chem. Soc.*, **105**, 1848-51 (1983a)

D. W. SINDORF and MACIEL G. E. Solid-state NMR studies of the reactions of silica surfaces with polyfunctional chlorosilanes and ethoxymethylsilanes. *J. Am. Chem. Soc.*, **105**, 3767-76 (1983b)

D. SLOTFELDT-ELLINGSEN and H. A. RESING. Motions of phenyl groups chemically bond to silica gel. Carbon-13 Nuclear Magnetic Resonance. *J. Phys. Chem.*, **84**, 2204-09 (1980)

D. R. STEWART, M. KRAWIEC, R. P. KASHYAP, W. H. WATSON, C. D. GUTSCHE. Conformational characteristics of ethers and esters of *p*-tert-butylcalix[5]arene. *J. Am. Chem. Soc.*, **117**, 586-601 (1995)

S. SUN, M. J. SEPANIAK, J.-S. WANG, C. D. GUTSCHE. Capillary electrokinetic chromatography employing *p*-(carboxyethyl) calix[*n*]arenes as running buffer additives. *Anal. Chem.*, **69**, 344-48 (1997)

T. SUZUKI, K. NAKASHIMA, S. SHINKAI. Influence of *para*-substituents and solvents on selective precipitation of fullerenes by inclusion in calix[8]arenes.

Tetrahedron Lett., **36**, 249-52 (1995)

T. SUZUKI, K. NAKASHIMA, S. SHINKAI. Very Convenient and efficient purification method for fullerene (C₆₀) with 5,11,17,23,35,41,47-octa-tert-butylcalix-[8]arene-49,50,51,52,53,54,55,56-octol. *Chem. Lett.*, 699-702 (1994)

B. I. SWANSON, D. LI, J.-X. SHI, S. JOHNSON, X. YANG. Smart-film sensors hydrocarbons for halogenated VOCs. *Proc. Int. Conf. Incineration Therm. Treat. Technol.*, 393-97 (1996)

Y. TANAKA, Y. KISHIMOTO, K. OTSUKA, S. TERABE. Strategy for separation solutions in capillary electrophoresis-mass spectrometry. *J. Chromatogr. A*, **817**, 49-57 (1998)

I. THONDORF and J. BRENN. Conformations of calix[5]arenes - a molecular mechanics study. *J. Chem. Soc., Perkin Trans. 2*, 2293-99 (1997)

I. THONDORF, G. HILLIG, W. BRANDT, J. BRENN, A. BARTH, V. BÖHMER. Molecular modelling study of a dissymmetric calix[4]arene and its methyl ethers. *J. Chem. Soc. Perkin Trans. 2*, 2259-67 (1994)

P. THUÉRY, M. NIERLICH, J. C. BRYAN, V. LAMARE, J.-F. DOZOL, Z. ASFARI, J. VICENS. Crown ether conformations in 1,3-calix[4]arene bis(crown ethers): crystal structures of a caesium complex and solvent adducts and molecular dynamics simulations. *J. Chem. Soc., Dalton Trans.*, 4191-4202 (1997)

F. THUNECKE, A. KÁLMÁN, F. KÁLMÁN, S. MA, A. S. RATHORE, C. HORVÁTH. Kinetic study on the *cis-trans* isomerization of peptidyl-proline dipeptides. *J. Chromatogr. A*, **744**, 259-72 (1996)

Y. S. TSANTRIZOS, W. CHEW, L. D. COLEBROOK, F. SAURIOL. Synthesis of a novel 1,4-bridged calix[8]arene „host“ cavity. *Tetrahedron Lett.*, **38**, 5411-14 (1997)

K. K. UNGER (Editor). Packings and stationary phases in chromatographic techniques. in: *Chromatographic Science Series*, Vol. 47, Marcel Dekker, New York und Basel, 1990

K. K. UNGER, N. BECKER, P. ROUMELIOTIS. Recent developments in the evaluation of chemically bonded silica packings for liquid chromatography. *J. Chromatogr.*, **125**, 115-27 (1976)

K. K. UNGER. *Handbuch der HPLC, Teil 1 - Leitfaden für Anfänger und Praktiker*. GIT Verlag, Darmstadt, 1989

K. K. UNGER. Hohlraumstruktur poröser Adsorbentien. *Angew. Chem.*, **8**, 331-43 (1972)

K. UNGER. Porous silica. Its properties and use as support in column liquid chromatography. *Journal of Chromatography Library*, Vol. 16, Elsevier Amsterdam (1975)

W. F. van GUNSTEREN und H. J. C. BERENDSEN. Moleküldynamik-Computer-simulationen; Methodik, Anwendungen und Perspektiven in der Chemie. *Angew. Chem.*, **102**, 1020-55 (1990)

W. P. van HOORN, F. C. J. M. van VEGGEL, D. N. REINHOUDT. Conformation of

hexahydroxycalix[6]arene. *J. Org. Chem.*, **61**, 7180-84 (1996)

H. A. M. VERHULST, L. J. M. van de VEN, J. W. de HAAN, H. A. CLAESSENS, F. EISENBEISS, C. A. CRAMERS. Patching in reversed-phase high-performance liquid chromatographic materials studied by solid-state NMR spectroscopy. *J. Chromatogr. A*, **687**, 213-21 (1994)

J. VICENS and V. BÖHMER. Calixarenes: A Versatile Class of Macrocyclic Compounds, Kluwer Academic Publishers, Dordrecht, 1991

C. VICOSO, M. J. F. REBELO, M. F. CAMOES. Studies with a sodium ion selective electrode using a calixarene as ionophore. *Port. Electrochim. Acta*, **15**, 297-301 (1997)

A. VINAS, S. GOMEZ, J. BERTRAN, F. TEIXIDOR, J.-F. DOZOL, H. ROUQUETTE. Cobalt bis(dicarbollide) derivatives as extractants for europium from nuclear wastes. *Chem. Commun. (Cambridge)*, **2**, 191-92 (1998)

R. VÖLKEL. High-resolution solid-state ¹³C-NMR spectroscopy of polymers. *Angew. Chem. Int. Ed. Engl.*, **27**, 1655-72 (1988)

J. WANG and J. LIU. Calixarene-coated amperometric detectors. *Anal. Chim. Acta*, **294**, 201-06 (1994)

P. K. WEINER and P. A. KOLLMAN. AMBER: Assisted model building with energy refinement. A general program for modeling molecules and their interactions. *J. Comput. Chem.*, **2**(3), 287-303 (1981)

P. K. WEINER, P. A. KOLLMAN, D. A. CASE, U. C. SINGH, C. GHIO, G. ALAGONA, S. PROFETA, P. WEINER. A new force field for molecular mechanical simulation of nucleic acids and proteins. *J. Am. Chem. Soc.*, **106**, 765-84 (1984)

C. J. WELCH, W. H. PIRKLE. Progress in the design of selectors for buckminsterfullerene. *J. Chromatogr.*, **609**, 89-101 (1992)

R. M. WILLIAMS and J. W. VERHOEVEN. Supramolecular encapsulation of fullerene c60 in a water-soluble calixarene: a core-shell charge-transfer complex. *Recl. Trav. Chim. Pays-Bas*, **111**, 531-32 (1992)

A. WUTTE, G. GÜBITZ, S. FRIEBE, G.-J. KRAUSS. HPLC separation of *cis-trans* isomers of proline-containing peptides. III. comparative studies with different stationary phases. *J. Chromatogr.* **677**, 186-91 (1994)

D.-Q. XIAO, Y. LING, Y.-X. WEN, R.-N. FU, J.-L. GU, R.-J. DAI, A.-Q. LUO. Synergistic effect of resorcarene and cyclodextrin mixed stationary phase in gas chromatography. *Chromatographia*, **46**, 177-82 (1997)

W. XU, J.-S. LI, Y.-Q. FENG, S.-L. DA, Y.-Y. CHEN, X.-Z. XIAO. Preparation and characterization of *p-tert*-butyl-calix[6]arene-bonded silica gel stationary phase for high-performance liquid chromatography. *Chromatographia*, **48** (3/4), 245-50 (1998)

P. K. ZARZYCKI, M. WIERZBOWSKA, H. LAMPARCZYK. The influence of temperature on the multiple separation of estrogenic steroids using mobile phases modified with β -cyclodextrin in high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, **15**, 1281-87 (1997)

S. ZELLMER and J. LASCH. Individual variation of human plantar stratum corneum lipids, determination by automated multiple development of high-performance thin-

- layer chromatography plates. *J. Chromatogr. B*, **691**, 321-29 (1997)
- Z. ZENG, J. WANG, X. TANG, S. TANG, X. LU. Two kinds of calix[4]arene derivatives as capillary gas chromatography stationary phases. *Fenxi Huaxue*, **26** (9), 1060-64 (1998)
- H. ZHANG, Y. LING, R. DAI, Y. WEN, R. FU, J. GU. Two alkyl resorcarenens as stationary phases for isomer separation by GC. *Chem. Lett.*, 225-26 (1997)
- S. ZHANG, W. ZHANG, C. WU, Z. ZHENLIN. Gas chromatography of calix[4]arene derivatives. *Fenxi Kexue Xuebao*, **14**, 14-18 (1998)
- Y. ZHANG and I. M. WARNER. Separation of water-soluble *p*-sulfonated calixarenes 4, 6 and 8 and 4-hydroxybenzene sulfonate by use of capillary zone electrophoresis. *J. Chromatogr. A*, **688**, 293-300 (1994)
- W. ZHANG, C. WU, J. WANG, S. ZHANG. A new calixcrown polysiloxane stationary phase for capillary gas chromatography. *Chinese J. Chromatogr.*, **15**, 204-5, (1997)
- T. ZHAO, X. HU, J. CHENG, X. LU. Use of calix[4]arene to separate positional isomers in capillary electrophoresis. *Anal. Chim. Acta*, **358**, 263-68 (1998)
- Z.-L. ZHONG, C.-P. TANG, C.-Y. WU, Y.-Y. CHEN. Synthesis and properties of calixcrown telomers. *J. Chem. Soc., Chem. Commun.*, 1737-38 (1995)
- P. L. ZHU. On the chromatography mechanism of reversed-phase liquid chromatography. *Chromatographia*, **21**(4), 229-33 (1986)