

## LITERATURVERZEICHNIS

- [1] Nomenclature committee of the international union of biochemistry and molecular biology (NC-IUBMB), Enzyme Supplement 5 (1999). Eur.J.Biochem. 264(2):610-650, 1999
- [2] Abramic M, Vitale L: New chloride-activated aminopeptidase from human erythrocytes. FEBS Letters 253:79-82, 1989
- [3] Abramic M, Vitale L: Basic Amino Acids Preferring Broad Specificity Aminopeptidase from Human Erythrocytes. Biol.Chem.Hoppe-Seyler 373:375-380, 1992
- [4] Aitken A, Learmonth M: Carboxymethylation of Cysteine using Iodacetamide/Iodacetic Acid, chap. 51, in The Protein Protocols Handbook, edited by John M.Walker 1996, pp 339-340
- [5] Amoscato AA, Alexander JW, Babcock W: Surface aminopeptidase activity of human lymphocytes. Biochemical and biologic properties of intact cells. J.Immunolog. 142(4):1245-1252, 1989
- [6] Ansorge S, Bühling F, Kähne T, Lendeckel U, Reinhold D, Träger M, Wrenger S: CD 26/dipeptidyl peptidase IV in lymphocyte growth regulation. Adv.Exp.Med.Biol. 421:127-140, 1997
- [7] Ansorge S, Langner J: *Cellular Peptidases in Immune Functions and Diseases* (vol 421). Plenum Press, New York, 1997
- [8] Anton LC, Snyder HL, Bennink JR, Vinitsky A, Orłowski M, Porgador A, Yewdell JW: Dissociation of proteasomal degradation of biosynthesized viral proteins from generation of MHC class I-associated antigenic peptides. J Immunol. 160:4859-4868, 1998
- [9] Aoyagi T, Yoshida S, Matsuda N, Ikeda T, Hamada M, Takeuchi T: Leuhistin, a new inhibitor of Aminopeptidase M, produced by *Bacillus laterosporus* BM156-14F1. The Journal of Antibiotics (Tokyo) 44 No.6:573-578, 1991

- 
- [10] Aoyagi T, Yoshida S, Nakamura Y, Shigihara Y, Hamada M, Takeuchi T: Probestin, a new inhibitor of aminopeptidase M, produced by *Streptomyces azureus* MH663-2F6. I. Taxonomy, production, isolation, physico-chemical properties and biological activities. *J Antibiot.(Tokyo)* 43:143-148, 1990
- [11] Aratani H, Kawata S, Tsuruyama S, Yoshida N, Makisumi S: Purification and Characterization of an Aminopeptidase from Bovine Leukocytes. *J.Biochem.* 96:107-115, 1984
- [12] Arfin SM, Bradshaw RA: Cotranslational processing and protein turnover in eukaryotic cells. *Biochemistry* 27:7979-7984, 1988
- [13] Arfin SM, Kendall RL, Hall L, Weaver LH, Stewart AE, Matthews BW, Bradshaw RA: Eukariotic methionyl aminopeptidases: Two classes of cobalt-dependent enzymes. *Biochemistry* 92:7714-7718, 1995
- [14] Arndt M, Lendeckel U, Spiess A, Faust J, Neubert K, Reinhold D, Ansorge S: Dipeptidyl peptidase IV (DP IV/CD26) mRNA expression in PWM-stimulated T-cells is suppressed by specific DP IV inhibition, an effect mediated by TGF-beta(1). *Biochem.Biophys.Res.Comm.* 274:410-414, 2000
- [15] Arndt M, Reinhold D, Lendeckel U, Spiess A, Faust J, Neubert K, Ansorge S: Specific inhibitors of dipeptidyl peptidase IV suppress mRNA expression of DP IV/CD26 and cytokines. *Adv.Exp.Med.Biol.* 477:139-143, 2000
- [16] Ashmun RA, Look TA: Metalloprotease Activity of CD13/Aminopeptidase N on the Surface of Human Myeloid Cells. *Blood* 75 No.2:462-469, 1990
- [17] Bank U, Kruger S, Langner J, Roessner A: Review: peptidases and peptidase inhibitors in the pathogenesis of diseases. Disturbances in the ubiquitin-mediated proteolytic system. Protease-antiprotease imbalance in inflammatory reactions. Role of cathepsins in tumour progression. *Adv.Exp.Med.Biol.* 477:349-378, 2000
- [18] Barrett AJ, Rawlings ND, Woessner JF: *Handbook of Proteolytic Enzymes.* (vol 343), Academic Press. ed. 1998
- [19] Bauvois B: Murine thymocytes possess specific cell surface-associated exoaminopeptidase activities: preferential expression by immature CD4-CD8-subpopulation. *Eur.J.Immunol.* 20:459-468, 1990
- [20] Belhacene N, Mari B, Rossi B: Characterization and purification of T lymphocyte aminopeptidase B: a putative marker of T cell activation. *Eur.J.Immunol.* 23:1948-1955, 1993

- 
- [21] Boes B, Hengel H, Ruppert T, Multhaup G, Koszinowski UH, Kloetzel PM: Interferon gamma stimulation modulates the proteolytic activity and cleavage site preference of 20S mouse proteasomes. *J.Exp.Med.* 179(3):901-909, 1994
- [22] Bongiorno L, Marini M, Urbani A, Ausiello CM, Roda LG: Effect of stimulation on soluble proteolytic enzymes released by peripheral blood mononuclear cells. *Int J Immunopharmacol* 15(5):621-629, 1993
- [23] Boyum A: Isolation of mononuclear cells and granulocytes from human blood. *J.Clin.Invest.* 22:923-930, 1968
- [24] Brou C, Logeat F, Gupta N, Bessia C, LeBail O, Doedens JR, Cumano A, Roux P, Black RA, Israel A: A novel proteolytic cleavage involved in Notch signaling: the role of the disintegrin-metalloprotease TACE. *Mol.Cell* 5:207-216, 2000
- [25] Brownlees J, Williams CH: Peptidases, peptides, and the mammalian blood-brain barrier. *J Neurochem.* 60:793-803, 1993
- [26] Chen H, Roques BP, Fournie-Zaluski MC: Design of the first highly potent and selective aminopeptidase N (EC 3.4.11.2) inhibitor. *Bioorg. Med.Chem.Lett.* 9:1511-1516, 1999
- [27] Chen W, Norbury CC, Cho Y, Yewdell JW, Bennink JR: Immunoproteasomes shape immunodominance hierarchies of antiviral CD8(+) T cells at the levels of T cell repertoire and presentation of viral antigens. *J Exp.Med.* 193:1319-1326, 2001
- [28] Constam DB, Rensing-Ehl A, Kemler K, Hersh LB, Fontana A: Puromycin-sensitive Aminopeptidase. *The Journal of Biological Chemistry* 270, No.45:26931-26939, 1995
- [29] Constam DB, Tobler AR, Schmitt-Gräff A, Malipiero U, Schlapbach R, Fontana A: Cloning of the Puromycin-Sensitive Aminopeptidase and Evidence for Expression in Neurons. *J.of Neurochemistry* 68:889-897, 1997
- [30] Costerousse O, Allegrini J, Lopez M, Alhenc-Gelas F: Angiotensin I - converting enzyme in human circulating mononuclear cells: genetic polymorphism of expression in T-lymphocytes. *Biochem.J* 290 ( Pt 1):33-40, 1993
- [31] Cotrell GS, Hooper NM, Turner AJ: Cloning, expression, and characterization of human cytosolic aminopeptidase P: a single Manganase(II)-dependent enzyme. *Biochemistry* 39 (49):15121-15128, 2000
- [32] Coux O, Tanaka K, Goldberg AL: Structure and functions of the 20S and 26S proteasomes. *Annu.Rev.Biochem.* 65:801-847, 1996

- 
- [33] Davidoff AN, Mendelov BV: Unexpected cytokinetic effects by puromycin include a G2-arrest, a metaphase-mitotic-arrest, and apoptosis. *Leukemia Res.* 16:1077-1085, 1992
- [34] Davidoff AN, Mendelov BV: Substituted purines elicit differential cytokinetic, molecular and phenotypic effects in HL-60 cells. *Exp.Hematol.* 21:456-460, 1993
- [35] de Gandarias JM, Irazusta J, Gil J, Fernandez D, Casis L: Brain soluble and membrane-bound Tyr-aminopeptidase activities during the stages of estrous and proestrous in the female rat. *Brain Res.* 620(1):146-148, 1993
- [36] de Souza ANC, Bruno JA, Carvalho KM: An enkephalin degrading Aminopeptidase of human brain preserved during the vertebrate phylogeny. *Comp.Biochem.Physiol.* 99 C:363-367, 1991
- [37] Desser L, Holomanova D, Zavadova E, Pavelka K, Mohr T, Herbacek I: Oral therapy with proteolytic enzymes decreases excessive TGF-beta levels in human blood. *Cancer Chemother.Pharmacol.* 47 Suppl:S10-S15, 2001
- [38] Doedens JR, Black RA: Stimulation-induced down-regulation of tumor necrosis factor-alpha converting enzyme. *J Biol.Chem.* 275:14598-14607, 2000
- [39] Dyer SH, Slaughter CA, Moomaw CR, Hersh LB: Comparison of the Soluble and Membrane-Bound Forms of the Puromycin-Sensitive Enkephalin-Degrading Aminopeptidases from Rat. *J.of Neurochemistry* 54:547-554, 1990
- [40] Engelbrecht S, Macartey HW, Pieper S, Rautenberg W, Tschesche H, Wenzel HR: Separation of the human Leucocyte Enzymes. *Hoppe-Seyler's Z.Physiol.Chem.* 363:305-315, 1982
- [41] Erbeznik H, Hersh LB: A Novel Mammalian High-Molecular-Weight Aminopeptidase. *Archives of Biochemistry and Biophysics* 344 No.1:228-234, 1997
- [42] Flores M, Aristoy MC, Toldra F: HPLC Purification and Characterization of Soluble Alanyl Aminopeptidase from Porcine Skeletal Muscle. *J.of Agricultural and Food Chemistry* 44, No.9:2578-2583, 1996
- [43] Gaczynska M, Rock KL, Goldberg AL: Role of proteasomes in antigen presentation. *Enzyme Protein* 47:354-369, 1993
- [44] Gibson AM, McDermott JR, Lauffart B, Mantle D: Specificity of action of human brain alanyl aminopeptidase on leu-enkephalin and dynorphin-related peptides. *Neuropeptides* 13:259-262, 1989

- 
- [45] Glickman MH: Getting in and out of the proteasome. *Semin.Cell Dev.Biol.* 11(3):149-158, 2000
- [46] Goding JW: Ecto-enzymes: physiology meets pathology. *J Leukoc.Biol.* 67:285-311, 2000
- [47] Grdisa M, Vitale L: Types and Localisation of Aminopeptidases in Different Human Blood Cells. *Int.J.Biochem.* 23 No.3:339-345, 1991
- [48] Griffin TA, Nandi D, Cruz M, Fehling HJ, Kaer LV, Monaco JJ, Colbert RA: Immunoproteasome assembly: cooperative incorporation of interferon gamma (IFN-gamma)-inducible subunits. *J Exp.Med.* 187:97-104, 1998
- [49] Grisk O, Küster U, Ansorge S: The activity of  $\gamma$ -glutamyl transpeptidase in population of mononuclear cells from human peripheral blood. *Biol.Chem.Hoppe-Seyler* 374:287-290, 1993
- [50] Harris ELV, Angal S: *Protein purification methods.* 1989.
- [51] Heemels MT, Ploegh H: Generation, translocation, and presentation of MHC class I-restricted peptides. *Annu.Rev.Biochem.* 64:463-491, 1995
- [52] Hendriks D, De Meester I, Umiel T, Vanhoof G: Aminopeptidase P and dipeptidyl peptidase IV activity in human leukocytes and stimulated lymphocytes. *Clinica Chimica Acta* 196:96, 1991
- [53] Hersh LB: Solubilization and Characterization of Two Rat Brain Membrane-Bound Aminopeptidases Active on Met-Enkephalin. *Biochemistry* 20:2345-2350, 1981
- [54] Hersh LB: Characterization of Membrane-Bound Aminopeptidases from Rat Brain: Identification of the Enkephalin-Degrading Aminopeptidase. *J.of Neurochemistry* 44:1427-1435, 1985
- [55] Hersh LB, McKelvy JF: An Aminopeptidase from Bovine Brain which catalyzes the Hydrolysis of Enkephalin. *J.of Neurochemistry* 36:171-178, 1981
- [56] Hiroi Y, Endo Y, Natori Y: Purification and properties of an aminopeptidase from rat-liver cytosol. *Arch.Biochem.Biophys.* 294:440-445, 1992
- [57] Hooper N: Families of zinc metalloproteases. *FEBS Letters* 354:1-6, 1994
- [58] Hooper NM, Turner AJ: Protein processing mechanisms: from angiotensin-converting enzyme to Alzheimer's disease. *Biochem.Soc.Trans.* 28:441-446, 2000

- 
- [59] Huang K, Takahara S, Kinouchi T, Takeyama M, Ishida T, Ueyama H, Nishi K, Ohkubo I: Alanyl aminopeptidase from human seminal plasma: purification, characterization, and immunohistochemical localization in the male genital tract. *J Biochem.(Tokyo)* 122:779-787, 1997
- [60] Hui KS, Hui MP, Lajtha A: Major Rat Brain Membrane-Associated and Cytosolic Enkephalin-degrading Aminopeptidases: Comparison studies. *Journal of Neuroscience Research* 20:231-240, 1988
- [61] Hui KS, Hui MP, Lajtha A, Saito M: Cellular Localization of Puromycin-Sensitive Aminopeptidase Isozymes. *Neurochemical Research* 15 No.12:1147-1151, 1990
- [62] Hui KS, Wang Y, Lajtha A: Purification and Characterization of an Enkephalin Aminopeptidase from Rat Brain Membranes. *Biochemistry* 22 No.5:1062-1067, 1983
- [63] Hui M, Budai ED, Lajtha A, Palkovits M, Hui KS: Changes in puromycin-sensitive aminopeptidases in post mortem schizophrenic brain regions. *Neurochem.Int.* 4/5:433-441, 1995
- [64] Janeway CA, Travers P: *Immunologie*. Spektrum Akademischer Verlag, Heidelberg, Berlin, Oxford, 1997
- [65] Jocelyn PC: *Biochemistry of the SH Group*, Academic Press, London, New York ed. 1972, 77-100
- [66] Julius MH, Simpson E, Herzenberg LA: A rapid method for the isolation of functional thymus-derived murine lymphocytes. *Eur.J.Immunol.* 3:645-649, 1973
- [67] Kahne T, Lendeckel U, Wrenger S, Neubert K, Ansorge S, Reinhold D: Dipeptidyl peptidase IV: a cell surface peptidase involved in regulating T cell growth (review). *Int.J Mol.Med.* 4:3-15, 1999
- [68] Kakuta H, Koiso Y, Takahashi H, Nagasawa K, Hashimoto Y: Novel specific puromycin-sensitive aminopeptidase inhibitors:3-(2,6-Diethylphenyl)-2,4 (1*H*,3*H*)-Quinazolin-2-one and N-(2,6-Diethylphenyl)-2-Amino-4*H*-3,1-Benzoxazin-4-One. *Heterocycles* 55, No.8:1433-1438, 2001
- [69] Kehlen A, Gohring B, Langner J, Riemann D: Regulation of the expression of aminopeptidase A, aminopeptidase N/CD13 and dipeptidylpeptidase IV/CD26 in renal carcinoma cells and renal tubular epithelial cells by cytokines and cAMP-increasing mediators. *Clin.Exp.Immunol.* 111:435-441, 1998

- [70] Kenny AJ, Turner A, Murray H: The aminopeptidase activity in the human T-cell lymphoma line (Jurkat) is not at the cell surface and is not aminopeptidase N (CD13). *Biochem.J* 298:353-360, 1994
- [71] Kleber HP, Schlee D, Schöpp W: *Biochemisches Praktikum*. VEB Gustav Fischer Verlag Jena, 1988
- [72] Kleining H, Sitte M: *Zellbiologie*. Gustav Fischer Verlag, Stuttgart, Jena, Lübeck, Ulm., 1999
- [73] Kohno H, Kanno T: Intracellular localization and properties of aminopeptidases in human peripheral lymphocytes. *Int.J.Biochem.* 17:1143-1148, 1985
- [74] Kohno H, Kanno T: Properties and activities of aminopeptidases in normal and mitogen-stimulated human lymphocytes. *Biochem.J.* 226:59-65, 1985
- [75] Komoda M, Kakuta H, Takahashi H, Fujimoto Y, Kadoya S, Kato F, Hashimoto Y: Specific inhibitor of puromycin-sensitive aminopeptidase with a homophthalimide skeleton: identification of the target molecule and a structure-activity relationship study. *Bioorganic & Medicinal Chemistry* 9(1):121-131, 2001
- [76] Kunz D, Bühling F, Hutter HJ, Aoagi T, Ansorge S: Aminopeptidase N (CD13, EC 3.4.11.2) occurs on the surface of resting and concavalin A-stimulated lymphocytes. *Biol.Chem.Hoppe-Seyler* 374(4):291-296, 1993
- [77] Langner J, Ansorge S: *Cellular Peptidases in Immune Functions and Diseases 2*. Kluwer Academic/ Plenum Publishers, New York, 2000
- [78] Lasch J, Moschner S, Sann H, Zellmer S, Koelsch R: Aminopeptidase P a cell-surface antigen of endothelial and lymphoid cells: catalytic and immunohistotopical evidences. *Biol.Chem.* 379:705-709, 1998
- [79] Lauffart B, Mantle D: Rationalization of aminopeptidase activities in human skeletal soluble extract. *Biochimica et Biophysica Acta* 956:300-306, 1988
- [80] Lendeckel U, Arndt M, Frank K, Wex T, Ansorge S: Role of alanyl aminopeptidase in growth and function of human T cells (review). *Int J Mol.Med.* 4:17-27, 1999
- [81] Lendeckel U, Kahne T, Arndt M, Frank K, Ansorge S: Inhibition of alanyl aminopeptidase induces MAP-kinase p42/ERK2 in the human T cell line KARPAS-299. *Biochem.Biophys.Res.Commun.* 252:5-9, 1998
- [82] Lendeckel U, Kähne T, Riemann D, Neubert K, Arndt M, Reinhold D: Review: The Role of Membrane peptidases in Immune Functions, in Cellular

---

Peptidases in Immune Functions and Diseases 2, Kluwer Academic/Plenum Publishers, New York ed., edited by Langner and Ansorge 2000, pp 1-24

- [83] Lendeckel U, Scholz B, Arnd M, Frank K, Spiess A, Chen A, Roques B, Ansorge S: Inhibition of Alanyl-Amino-peptidase Suppress the Activation-Dependent Induktion of Glykogen Synthase Kinase-3 $\beta$  in Human T cells. *Biochemical and Biophysical Research Communications* 273:62-65, 2000
- [84] Lendeckel U, Wex T, Ittenson A, Arndt M, Frank K, Mayboroda O, Schubert W, Ansorge S: Rapid mitogen-induced amino-peptidase N surface expression in human T cells is dominated by mechanism independent of de novo protein biosynthesis. *Immunobiology* 197:55-69, 1997
- [85] Lendeckel U, Wex T, Kähne T, Frank K, Reinhold D, Ansorge S: Expression of the Amino-peptidase N (CD13) Gene in the Human T Cell Lines HuT78 and H9 Cellular Immunology 153:214-226, 1994
- [86] Lendeckel U, Wex T, Reinhold D, Arndt M, Ittenson A, Frank K, Ansorge S: Activation-dependent induction of T cell alanyl amino-peptidase and its possible involvement in T cell growth. *Adv.Exp.Med.Biol.* 421:59-66, 1997
- [87] Lendeckel U, Wex T, Reinhold D, Kähne T, Frank K, Faust J, Neubert K, Ansorge S: Induction of the membrane alanyl amino-peptidase gene and surface expression in human T-cells by mitogenic activation. *Biochem.J* 319:817-821, 1996
- [88] Letarte M, Vera S, Tran R, Addis JB, Onizuka RJ, Quackenbush EJ, Jongeneel CV, McInnes R: Common acute lymphocytic leukemia antigen is identical to neutral endopeptidase. *J.Exp.Med.* 168(4):1247-1253, 1988
- [89] Li X, Chang YH: Evidence that the human homologue of a rat initiation factor-2 associated protein (p<sup>67</sup>) is a methionine amino-peptidase. *Biochemical and Biophysical Research Communications* 227(1):152-159, 1996
- [90] Look TA, Ashmun RA, Shapiro LH, Peiper SC: Human Myeloid Plasma Membrane Glycoprotein CD13 is identical to Amino-peptidase N. *J.Clin.Invest.* 83:1299-1307, 1989
- [91] Lowther WT, Matthews BW: Structure and function of the methionine amino-peptidases. *BBA* 1477(1-2):157-167, 2000
- [92] Luckey CJ, King GM, Marto JA, Venketeswaran S, Maier BF, Crotzer V, Colella TA, Shabanowitz J, Hunt DF, Engelhard VH: Proteasomes can either generate or destroy MHC class I epitopes: evidence for nonproteasomal epitope generation in the cytosol. *J.Immunolog.* 161(1):112-121, 1998



- 
- [93] Lundgren E, Roos G, Tärnik A: Subcellular Distribution of Amino Acid Naphthylamidase Isozymes in human Leucocytes. *Acta path. microbiol. scand. Sect.A* 83:176-182, 1975
- [94] Mantle D: Comparison of soluble aminopeptidases in human cerebral cortex, skeletal muscle and kidney tissues. *Clinica Chimica Acta* 207:107-118, 1992
- [95] Mantle D, Hardy MF, Lauffart B, McDermott JR, Smith AI, Pennington RJT: Purification and characterization of the major aminopeptidase from human skeletal muscle. *Biochem.J* 211:567-573, 1983
- [96] Mantle D, Lauffart B, Gibson A: Purification and characterization of leucyl aminopeptidase and pyroglutamyl aminopeptidase from human skeletal muscle. *Clinica Chimica Acta* 197(1):35-45, 1991
- [97] Mantle D, Lauffart B, McDermott J, Gibson A: Characterization of aminopeptidases in human kidney soluble fraction. *Clinica Chimica Acta* 187:105-114, 1990
- [98] Mantle D, Lauffart B, McDermott JR, Kidd AM, Pennington RJ: Purification and charakterisation of two Cl-activated aminopeptidases hydrolysing basic termini from human skeletal muscle. *J.Biochem.* 147:307-312, 1985
- [99] Mäkinen KK, Mäkinen PL: Purification and Characterization of Two Human Erythrocyte Arylamidases preferentially Hydrolysing N-Terminal Arginine or Lysine Residues. *Biochem.J* 175:1051-1068, 1978
- [100] McDermott JR, Mantle D, Lauffart B, Kidd AM: Purification and Characterization of a Neuropeptide-Degrading Aminopeptidase from human Brain. *J.of Neurochemistry* 45:752-759, 1985
- [101] McDonald JK: Exopeptidases, in *Mammalian Proteases* London Academic Press, 1986,
- [102] McLellan S, Dyer S, Rodriguez G, Hersh L: Studies of the Tissue Distribution of the Puromycin-Sensitive Enkephalin-Degrading Aminopeptidases. *J.of Neurochemistry* 51:1552-1559, 1988
- [103] Murray H, McClaren H, Turner A, Kenny AJ: Immunocytochemical Studies of an Antigen in a Human T-cell Lymphoma Line (Jurkat) Recognized by Certain Monoclonal Antibodies to CD-13 ( Aminopeptidase-N). *Leukemia and Lymphoma* 17:501-508, 1995
- [104] Nagai M, Kojima F, Naganawa H, Hamada M, Aoyagi T, Takeuchi T: Phebestin, a new inhibitor of aminopeptidase N, produced by *Streptomyces* sp. MJ716-m3. *J.Antibiot.(Tokyo)* 50:82-84, 1997

- 
- [105] Nagaoka I, Yamashita T: Leucine Aminopeptidase as an ecto-Enzyme of polymorphonuclear neutrophils. *Biochimica et Biophysica Acta* 598(1):169-172, 1980
- [106] Nagaoka I, Yamashita T: Inactivation during phagocytosis of leucine aminopeptidase, an ecto-enzyme of polymorphonuclear neutrophils. *BBA* 678:342-351, 1981
- [107] Nagaoka IYT: Inactivation of phagocytosis-stimulating activity of Tuftsin by polymorphonuclear neutrophils. *BBA* 675:85-93, 81 A.D.
- [108] Nagata Y, Mizutani S, Nomura S, Kurauchi O, Kasugai M, Tomoda Y: Purification and properties of human placental aminopeptidase B. *Enzyme* 45(4):165-173, 1991
- [109] Nano R, De Piceis Polver P, Gerzeli G: Leucine Aminopeptidase activity in blood and hemopoietic organs of different Vertebrates. *Acta histochem.* 82:83-88, 1987
- [110] Naquet P, Pierres M: Cell-surface enzymes and lymphocyte functions. *Current Opinion in Immunology* 3:326-329, 1991
- [111] Ohkubo K, Baraniuk JN, Hohman R, Merida M, Hersh LB, Kaliner MA: Aminopeptidase activity in human nasal mucosa. *J Allergy Clin Immunol* 102(5):741-750, 1998
- [112] Osada M, Ikegami S, Takiguchi-Hayashi K, Yamazaki Y, Katoh-Fukui Y, Higashina-Kagawa T, Sakaki Y, Takeuchi T: Increased anxiety and impaired pain response in puromycin-sensitive aminopeptidase gene-deficient mice obtained by a mouse gene-trap method. *Journal of Neuroscience Research* 19(14):6068-6078, 1999
- [113] Osada M, Watanabe G, Kondo S, Toyoda M, Sakaki Y, Takeuchi T: Male Reproductive Defects Caused by Puromycin-Sensitive Aminopeptidase Deficiency in Mice. *Mol.Endocrinol.* 15(6):960-971, 2001
- [114] Osada M, Watanabe G, Sakaki Y, Takeuchi T: Puromycin-sensitive Aminopeptidase Is Essential for the Maternal Recognition of Pregnancy in Mice. *Mol.Endocrinol.* 15(6):882-893, 2001
- [115] Patterson EK, Hsiao SH, Keppel A: Studies on Dipeptidases and Aminopeptidases. *The Journal of Biological Chemistry* 238 No.11:3611-3620, 1963
- [116] Proost P, Struyf S, Schols D, Durinx C, Wuyts A, Lenaerts JP, De Clercq E, De Meester I, Van Damme J: Processing by CD26/dipeptidyl-peptidase IV

- reduces the chemotactic and anti-HIV-1 activity of stromal-cell-derived factor-1alpha. *FEBS Lett.* 432:73-76, 1998
- [117] Rautenberg W, Tschesche H: Detection and Partial Purification of Four Proteases from Human Leucocytes. *Biol.Chem.Hoppe-Seyler* 362:230, 1981
- [118] Rautenberg W, Tschesche H: Aminopeptidases from Human Leucocytes. *Hoppe-Seyler's Z.Physiol.Chem.* 365:49-58, 1984
- [119] Rawlings ND, Barrett AJ: Evolutionary families of peptidases. *Biochem.J* 290 ( Pt 1):205-218, 1993
- [120] Rehli M, Krause SW, Kreutz M, Andreesen R: Carboxypeptidase M is identical to the MAX.1 antigen and its expression is associated with monocyte to macrophage differentiation. *J Biol.Chem.* 270:15644-15649, 1995
- [121] Reinhold D, Bank U, Buhling F, Lendeckel U, Ulmer AJ, Flad HD, Ansorge S: Transforming growth factor-beta 1 (TGF-beta 1) inhibits DNA synthesis of PWM-stimulated PBMC via suppression of IL-2 and IL-6 production. *Cytokine* 6:382-388, 1994
- [122] Reinhold D, Bank U, Böhling F, Täger M, Born I, Neubert K, Faust J, Ansorge S: Inhibitors of dipeptidyl peptidase IV (DPIV,CD 26) induces secretion of transforming growth factor-beta 1 (TGF-beta 1) in stimulated mouse splenocytes and thymocytes. *Immunol.Lett.* 58(1):29-35, 1997
- [123] Reinhold D, Hemmer B, Gran B, Steinbrecher A, Brocke S, Kahne T, Wrenger S, Born I, Faust J, Neubert K, Martin R, Ansorge S: Dipeptidyl peptidase IV (CD26): role in T cell activation and autoimmune disease. *Adv.Exp.Med.Biol.* 477:155-160, 2000
- [124] Riemann D, Kehlen A, Langner J: Stimulation of the expression and the enzyme activity of aminopeptidase N/CD13 and dipeptidylpeptidase IV/CD26 on human renal cell carcinoma cells and renal tubular epithelial cells by T cell-derived cytokines, such as IL-4 and IL-13. *Clin.Exp.Immunol.* 100:277-283, 1995
- [125] Rivett AJ, Bose S, Brooks P, Broadfoot KI: Regulation of proteasome complexes by gamma-interferon and phosphorylation. *Biochimie.* 83:363-366, 2001
- [126] Rusu I, Yaron A: Aminopeptidase P from human leukocytes. *Eur.J Biochem.* 210:93-100, 1992

- 
- [127] Safavi A, Herrsh LB: Degradation of Dynorphin-Related Peptides by the Puromycin-Sensitive Aminopeptidase and Aminopeptidase M. *J. of Neurochemistry* 65:389-395, 1995
- [128] Saito M, Takegoshi K, Aoyagi T, Umezawa H, Hagai Y: Stimulatory Effect of Bestatin, A New specific Inhibitor of Aminopeptidases, on the Blastogenesis of Guinea Pig Lymphocytes. *Cellular Immunology* 40:247-262, 1978
- [129] Sandernik GJ, Artur Y, Siest G: Human Aminopeptidases: A Review of the literature. *J.Clin.Chem.Clin.Biochem.* 26 No.12:765-807, 1988
- [130] Schellenberger A: *Enzymkatalyse* (vol 2). VEB Gustav Fischer Verlag Jena, 1989
- [131] Schnebli HP, Phillipps MA, Bardey RK: Isolation and characterization of an enkephalin-degrading aminopeptidase from rat brain. *Biochimica et Biophysica Acta* 569:89-98, 1979
- [132] Scornik OA, Botbol V: Bestatin as an experimental tool in mammals. *Curr Drug Metab* 2:67-85, 2001
- [133] Shah T, Webster DB, Peters T: Enzyme Analysis and subcellular fractionation of human peripheral blood lymphocytes with special reference to the localisation of putative plasma membrane enzymes. *Cell Biochemistry and Function* 1:117-124, 1983
- [134] Sharma KK, Elsner NJ, Kester K: Comparison of leucine aminopeptidase and aminopeptidase III activities in lens. *Curr Eye Res* 15(7):774-781, 1996
- [135] Sharma KK, Ortwerth BJ: Aminopeptidase III activity in normal and cataractous lenses. *Curr Eye Res* 5(5):373-380, 1986
- [136] Sharma KK, Ortwerth BJ: Isolation and Characterization of a New Aminopeptidase from Bovine Lens. *The Journal of Biological Chemistry* 261 No.9:4295-4301, 1986
- [137] Shipp AM, Look AT: Hämatopoietic Differentiation Antigens That Are Membrane-Associated Enzymes: Cuuting is the key? *Blood* 82(4):1052-1070, 1993
- [138] Smith GP, MacGregor RR, Peters TJ: Subcellular localisation of leucine aminopeptidase in human polymorphonuclear leukocytes. *Biochimica et Biophysica Acta* 728:222-227, 1983

- [139] Söderling E: Substrate Specificities of Cl<sup>-</sup>-Activated Arginine Aminopeptidases from Human and Rat Origin. *Archives of Biochemistry and Biophysics* 220:1-10, 1983
- [140] Sprinkle TJ, Caldwell C, Ryan JW: Cloning, chromosomal sublocalization of the human soluble aminopeptidase P gene (XPNPEP1) to 10q25.3 and conservation of nthe putative proton shuttle and metal ligand binding sites with XPNPEP2. *Arch.Biochem.Biophys.* 378(1):51-56, 2000
- [141] Steinbrecher A, Reinhold D, Quigley L, Gado A, Tresser N, Izikson L, Born I, Faust J, Neubert K, Martin R, Ansorge S, Brocke S: Dipeptidyl peptidase IV in inflammatory CNS disease. *Adv.Exp.Med.Biol.*; 477:145-153, 2000
- [142] Stoltze L, Schirle M, Schwarz M, Schroter C, Thompson MW, Hersh LB, Kalbacher H, Stevanovic S, Rammensee HG, Sxchild H: Two new proteases in the MHC class I processing pathway. *Nat.Immunol.* 1(5):413-418, 2000
- [143] Suda H, Aoyagi T, Takeuchi T, Umezawa H: Inhibition of Aminopeptidase B and Leucine Aminopeptidase by Bestatin and Its stereoisomer. *Archives of Biochemistry and Biophysics* 177:196-200, 1976
- [144] Swanson AA, Davis RM, Albers-Jackson B, McDonald JK: Lens exopeptidases. *Curr Eye Res* 32 (2):163-173, 1981
- [145] Takahashi H, Komoda M, Kakuta H, Hashimoto Y: [Preparation of novel specific aminopeptidase inhibitors with a cyclic imide skeleton]. *Yakugaku Zasshi* 120:909-921, 2000
- [146] Tanaka K: Role of proteasomes modified by interferon-gamma in antigen processing. *J Leukoc.Biol.* 56:571-575, 1994
- [147] Taylor A: Aminopeptidases: structure and function. *FASEB* 7:290-298, 1993
- [148] Taylor A: Aminopeptidases: towards a mechanism of action. *TIBS* 18:167-172, 1993
- [149] Thompson MW, Tobler A, Fontana A, Hersh LB: Cloning and Analysis of the Gene for the Human Puromycin-Sensitive Aminopeptidase. *Biochemical and Biophysical Research Communications* 258:234-240, 1999
- [150] Ulmer AJ, Mattern T, Feller AC, Heymann E, Flad HD: CD26 antigen is a surface dipeptidyl peptidase IV (DPIV) as characterized by monoclonal antibodies clone TII-19-4-7 and 4EL1C7. *Scand.J.Immunol.* 31(4):429-435, 1990

- 
- [151] Umezawa H, Aoyagi T, Tanaka T, Suda H, Okuyama A, Naganawa H, Hamada M, Takeuchi T: Production of actinonin, an inhibitor of aminopeptidase M, by actinomycetes. *J Antibiot.(Tokyo)* 38:1629-1630, 1985
- [152] Van den Eynde BJ, Morel S: Differential processing of class-I-restricted epitopes by the standard proteasome and the immunoproteasome. *Curr Opin.Immunol.*; 13:147-153, 2001
- [153] van Hal PT, Hopstaken-Broos JP, Prins A, Favalaro EJ, Huijbens RJ, Hilvering C, Figdor CG, Hoogsteden HC: Potential indirect anti-inflammatory effects of IL-4. Stimulation of human monocytes, macrophages, and endothelial cells by IL-4 increases aminopeptidase-N activity (CD13; EC 3.4.11.2). *J Immunol.* 153:2718-2728, 1994
- [154] Vanhoof G, De M, I, Goossens F, Hendriks D, Scharpe S, Yaron A: Kininase activity in human platelets: cleavage of the Arg1-Pro2 bond of bradykinin by aminopeptidase P. *Biochem.Pharmacol.* 44:479-487, 1992
- [155] Vinitzky A, Anton LC, Snyder HL, Orłowski M, Bennink JR, Yewdell J: The generation of MHC class I-associated peptides is only partially inhibited by proteasome inhibitors: involvement of nonproteasomal cytosolic proteases in antigen processing? *J.Immunolog.* 159(2):554-564, 1997
- [156] Vitale L, Grdisa M, Wrisher M: Human Polymorphonuclear Leukocytes Aminopeptidases. *Folia Histochemica et Cytobiologica* 24 No.2:139-148, 1986
- [157] Watanabe Y, Kumagai Y, Shimamori Y, Fujimoto Y: Purification and Characterization of Thiol Aminopeptidase from the Cytosolic Fraction of Human Placenta. *Biochemical Medicine and Metabolic Biology* 37:235-245, 1987
- [158] Welch PA: Regulation of B cell precursor proliferation by aminopeptidase A. *Int Immunol.* 7:737-746, 1995
- [159] Wex T, Lendeckel U, Kähne T, Ittenson A, Frank K, Ansorge S: The main neutral aminopeptidase activity of human lymphoid tumor cell lines does not originate from the aminopeptidase N-( APN; CD 13) gene. *Biochimica et Biophysica Acta* 1355:147-154, 1997
- [160] Wilk S, Wilk E, Magnusson RP: Purification, characterization, and cloning of a cytosolic aspartyl aminopeptidase. *J.Biol.Chem.* 273(26):15961-15970, 1998
- [161] Wu Q, Li L, Cooper MD, Pierres M, Gorvel JP: Aminopeptidase A activity of the murine B-lymphocyte differentiation antigen BP-1/6C3. *Proc. Natl.Acad.Sci.U.S.A* 88:676-680, 1991

- 
- [162] Xu Y, Lai LT, Gabrilowe JL, Scheinberg DA: Antitumor activity of actinonin in vitro and in vivo. *Clin.Cancer Res* 4(1):171-176, 1998
- [163] Yamamoto S, Higuchi Y, Yoshiyama K, Shimizu E, Kataoka M, Hijiya N, Matsuura K: ADAM family proteins in the immune system. *Immunol.Today* 20:278-284, 1999
- [164] Yamamoto Y, Li YH, Ushiyama I, Nishimura A, Ohkubo I, Nishi K: Puromycin-sensitive alanyl aminopeptidase from human liver cytosol: purification and characterization. *Forensic Sci Int* 113(1-3):143-146, 2000
- [165] Yamamoto Y, Yao-Hua L, Huang K, Ohkubo I, Nishi M: Isolation and Characterization of an Alanyl Aminopeptidase from Rat Liver Cytosol as a Puromycin-Sensitive Enkephalin-Degrading Aminopeptidase. *Biol.Chem.* 379:711-719, 1998
- [166] Yoshida S, Naganawa H, Aoyagi T, Takeuchi T, Takeuchi Y, Kodama Y: Leuhistin, a new inhibitor of aminopeptidase M, produced by *Bacillus laterosporus* BMI156-14F1. II. Structure determination of leuhistin. *J.Antibiot.(Tokyo)* 44:579-581, 1991
- [167] Yoshida S, Nakamura Y, Naganawa H, Aoyagi T, Takeuchi T: Probestin, a new inhibitor of aminopeptidase M, produced by *Streptomyces azureus* MH663-2F6. II. Structure determination of probestin. *J Antibiot.(Tokyo)* 43:149-153, 1990
- [168] Ziaber J, Baj Z, Pasnik J, Chmielewski H, Tchorzewski H: Increased expression of neutral endopeptidase (NEP) and aminopeptidase N (APN) on peripheral blood mononuclear cells in patients with multiple sclerosis. *Immunol.Lett.* 71 (2):127-129, 2000