

6 Literaturverzeichnis

- Ahmad, S., L. Wang, und P. E. Ward. 1992. Dipeptidyl(amino)peptidase IV and aminopeptidase M metabolize circulating substance P *in vivo*. *J. Pharmacol. Exp. Ther.* 260:1257-1261.
- Ansorge, S. und E. Schön. 1987. Dipeptidyl peptidase IV (DP IV), a functional marker of the T lymphocyte system. *Acta Histochem.* 82:41-46.
- Aoyagi, T., M. Nagai, K. Ogawa, F. Kojima, M. Okada, T. Ikeda, M. Hamada, und T. Takeuchi. 1991. Poststatin, a new inhibitor of prolyl endopeptidase, produced by *Streptomyces viridochromogenes* MH534-30F3. I. Taxonomy, production, isolation, physico-chemical properties and biological activities. *J. Antibiot. (Tokyo)* 44:949-955.
- Aoyagi, T., T. Wada, M. Nagai, F. Kojima, S. Harada, T. Takeuchi, H. Takahashi, K. Hirokawa, und T. Tsumita. 1990. Increased gamma-aminobutyrate aminotransferase activity in brain of patients with Alzheimer's disease. *Chem. Pharm. Bull. (Tokyo)* 38:1748-1749.
- Atack, J. R., N. Suman-Chauhan, G. Dawson, und J. J. Kulagowski. 1991. *In vitro* and *in vivo* inhibition of prolyl endopeptidase. *Eur. J. Pharmacol.* 205:157-163.
- Augustinack, J. C., A. Schneider, E. M. Mundelkow, und B. T. Hyman. 2002. Specific tau phosphorylation sites correlate with severity of neuronal cytopathology in Alzheimer's disease. *Acta Neuropathol. (Berl)* 103:26-35.
- Barelli, H., A. Petit, E. Hirsch, S. Wilk, G. De Nanteuil, P. Morain, und F. Checler. 1999. S 17092-1, a highly potent, specific and cell permeant inhibitor of human proline endopeptidase. *Biochem. Biophys. Res. Commun.* 257:657-661.
- Berridge, M. J. 1993. Cell signalling. A tale of two messengers. *Nature* 365:388-389.
- Berridge, M. J. 1998. Neuronal calcium signaling. *Neuron* 21:13-26.
- Berridge, M. J., C. P. Downes, und M. R. Hanley. 1989. Neural and developmental actions of lithium: a unifying hypothesis. *Cell* 59:411-419.
- Bezchlibnyk, Y. und L. T. Young. 2002. The neurobiology of bipolar disorder: focus on signal transduction pathways and the regulation of gene expression. *Can. J. Psychiatry* 47:135-148.
- Bhat, N. R. 1995. Signal transduction mechanisms in glial cells. *Dev. Neurosci.* 17:267-284.
- Birney, Y. A. und B. F. O'Connor. 2001. Purification and characterization of a Z-pro-prolinal-insensitive Z-Gly-Pro-7-amino-4-methyl coumarin-hydrolyzing peptidase from bovine serum - a new proline-specific peptidase. *Protein Expr. Purif.* 22:286-298.
- Blackburn, R. V., S. S. Galoforo, C. M. Berns, N. M. Motwani, P. M. Corry, und Y. J. Lee. 1998. Differential induction of cell death in human glioma cell lines by sodium nitroprusside. *Cancer* 82:1137-1145.
- Bouras, M., J. F. Huneau, C. Luengo, C. Erlanson-Albertsson, und D. Tome. 1995. Metabolism of enterostatin in rat intestine, brain membranes, and serum: differential involvement of proline-specific peptidases. *Peptides* 16:399-405.
- Bradford, M.M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the prinziple of protein-dye binding. *Anal. Biochem.* 72:248-254.
- Brion, J. P. 1998. Neurofibrillary tangles and Alzheimer's disease. *Eur. Neurol.* 40:130-140.
- Caffrey, J. J., K. Hidaka, M. Matsuda, M. Hirata, und S. B. Shears. 1999. The human and rat forms of multiple inositol polyphosphate phosphatase: functional homology with a histidine acid phosphatase up-regulated during endochondral ossification. *FEBS Lett.* 442:99-104.

- Camargo, A. C., M. L. Almeida, und P. C. Emson. 1984. Involvement of endo-oligopeptidases A and B in the degradation of neurotensin by rabbit brain. *J. Neurochem.* 42:1758-1761.
- Cohen, S. N., A. C. Chang, und L. Hsu. 1972. Nonchromosomal antibiotic resistance in bacteria: genetic transformation of *Escherichia coli* by R-factor DNA. *Proc. Natl. Acad. Sci. U.S.A* 69:2110-2114.
- Crawford, J. L., W. N. Lipscomb, und C. G. Schellman. 1973. The reverse turn as a polypeptide conformation in globular proteins. *Proc. Natl. Acad. Sci. U.S.A* 70:538-542.
- Creighton, E. T. 1978. Possible implications of many proline residues for the kinetics of protein unfolding and refolding. *J. Mol. Biol.* 125:401-406.
- Crumpton, M. J. und D. Snary. 1974. Preparation and properties of lymphocyte plasma membrane. *Contemp. Top. Mol. Immunol.* 3:27-56.
- Cruzblanca, H., D. S. Koh, und B. Hille. 1998. Bradykinin inhibits M current via phospholipase C and Ca²⁺ release from IP₃-sensitive Ca²⁺ stores in rat sympathetic neurons. *Proc. Natl. Acad. Sci. U.S.A* 95:7151-7156.
- Cunningham, C. C., R. Vegners, R. Bucki, M. Funaki, N. Korde, J. H. Hartwig, T. P. Stossel, und P. A. Janmey. 2001. Cell permeant polyphosphoinositide-binding peptides that block cell motility and actin assembly. *J. Biol. Chem.* 276:43390-43399.
- Cunningham, D. F. und B. O'Connor. 1997. Proline specific peptidases. *Biochim. Biophys. Acta* 1343:160-186.
- De Wied, D. 1997. Neuropeptides in learning and memory processes. *Behav. Brain Res.* 83:83-90.
- Defea, K., F. Schmidlin, O. Dery, E. F. Grady, und N. W. Bunnett. 2000. Mechanisms of initiation and termination of signalling by neuropeptide receptors: a comparison with the proteinase-activated receptors. *Biochem. Soc. Trans.* 28:419-426.
- Demuth, H.-U., D. Schlenzig, A. Schierhorn, G. Grosche, M. P. Chapot-Chartier, und J. C. Gripon. 1993. Design of (omega-N-(O-acyl)hydroxy amid) aminodicarboxylic acid pyrrolidides as potent inhibitors of proline-specific peptidases. *FEBS Lett.* 320:23-27.
- Diefenthal, T., H. Dargatz, V. Witte, G. Reipen, und I. Svendsen. 1993. Cloning of proline-specific endopeptidase gene from *Flavobacterium meningosepticum*: expression in *Escherichia coli* and purification of the heterologous protein. *Appl. Microbiol. Biotechnol.* 40:90-97.
- Dragunow, M. 1996. A role for immediate-early transcription factors in learning and memory. *Behav. Genet.* 26:293-299.
- Dresdner, K., L. A. Barker, M. Orłowski, und S. Wilk. 1982. Subcellular distribution of prolyl endopeptidase and cation-sensitive neutral endopeptidase in rabbit brain. *J. Neurochem.* 38:1151-1154.
- Eistetter, H. R., A. Mills, R. Brewster, S. Alouani, C. Rambosson, und E. Kawashima. 1992. Functional characterization of neurokinin-1 receptors on human U373MG astrocytoma cells. *Glia* 6:89-95.
- Ekman, R., P. Juhasz, M. Heilig, H. Agren, und C. E. Costello. 1996. Novel neuropeptide Y processing in human cerebrospinal fluid from depressed patients. *Peptides* 17:1107-1111.
- Feany, M. B. 1996. Neuropeptide modulation of learning and memory processes. *Rev. Neurosci.* 7:151-164.
- Feldkamp, M. M., N. Lau, und A. Guha. 1997. Signal transduction pathways and their relevance in human astrocytomas. *J. Neurooncol.* 35:223-248.
- Fiedorowicz, A., I. Figiel, B. Kaminska, M. Zaremba, S. Wilk, und B. Oderfeld-Nowak. 2001. Dentate granule neuron apoptosis and glia activation in murine hippocampus induced by trimethyltin exposure. *Brain Res.* 912:116-127.
- Fukuda, M. und K. Mikoshiba. 1997. The function of inositol high polyphosphate binding proteins. *Bioessays* 19:593-603.

- Fukuda, M. und K. Mikoshiba. 2000. Expression of synaptotagmin I or II promotes neurite outgrowth in PC12 cells. *Neurosci. Lett.* 295:33-36.
- Fulop, V., Z. Bocskei, und L. Polgar. 1998. Prolyl oligopeptidase: an unusual beta-propeller domain regulates proteolysis. *Cell* 94:161-170.
- Fulop, V., Z. Szeltner, und L. Polgar. 2000. Catalysis of serine oligopeptidases is controlled by a gating filter mechanism. *EMBO Rep.* 1:277-281.
- Gallegos, M. E., M. M. Zannatha, E. G. Osornio, A. S. Sanchez, und F. A. Posadas del rio. 1999. The activities of six exo-and endopeptidases in the substantia nigra, neostriatum, and cortex of the rat brain. *Neurochem. Res.* 24:1557-1561.
- Gee, N. S., C. I. Ragan, K. J. Watling, S. Aspley, R. G. Jackson, G. G. Reid, D. Gani, und J. K. Shute. 1988a. The purification and properties of myo-inositol monophosphatase from bovine brain. *Biochem. J.* 249:883-889.
- Gee, N. S., G. G. Reid, R. G. Jackson, R. J. Barnaby, und C. I. Ragan. 1988b. Purification and properties of inositol-1,4-bisphosphatase from bovine brain. *Biochem. J.* 253:777-782.
- Goossens, F., M. De, I. G. Vanhoof, und S. Scharpe. 1992. A sensitive method for the assay of serum prolyl endopeptidase. *Eur. J. Clin. Chem. Clin. Biochem.* 30:235-238.
- Goossens, F., M. De, I. G. Vanhoof, und S. Scharpe. 1996. Distribution of prolyl oligopeptidase in human peripheral tissues and body fluids. *Eur. J. Clin. Chem. Clin. Biochem.* 34:17-22.
- Goossens, F., G. Vanhoof, M. De, I. K. Augustyns, M. Borloo, D. Tourwe, A. Haemers, und S. Scharpe. 1997. Development and evaluation of peptide-based prolyl oligopeptidase inhibitors--introduction of N-benzyloxycarbonyl-prolyl-3-fluoropyrrolidine as a lead in inhibitor design. *Eur. J. Biochem.* 250:177-183.
- Greene, L. J., A. C. Spadaro, A. R. Martins, W. D. Perussi De Jesus, und A. C. Camargo. 1982. Brain endo-oligopeptidase B: a post-proline cleaving enzyme that inactivates angiotensin I and II. *Hypertension* 4:178-184.
- Griffiths, E. C., J. R. McDermott, und A. I. Smith. 1982. Inactivation of thyrotropin-releasing hormone (TRH) and (3Me-His) TRH by brain peptidases studied by high-performance liquid chromatography. *Neurosci. Lett.* 28:61-65.
- Hagihara, M. und T. Nagatsu. 1987. Post-proline cleaving enzyme in human cerebrospinal fluid from control patients and parkinsonian patients. *Biochem. Med. Metab. Biol.* 38:387-391.
- Hasebe, T., J. Hua, A. Someya, P. Morain, F. Checler, und I. Nagaoka. 2001. Involvement of cytosolic prolyl endopeptidase in degradation of p40-phox splice variant protein in myeloid cells. *J. Leukoc. Biol.* 69:963-968.
- Holloschi, A. und M. Hafner. 2002. A new green fluorescent protein reporter cell line to measure the bioactivity of calcitonin. *Eur. J. Cell Biol.* 79:50.
- Hou, X. E. und A. Dahlstrom. 2000. Synaptic vesicle proteins and neuronal plasticity in adrenergic neurons. *Neurochem. Res.* 25:1275-1300.
- Inagaki, N., H. Fukui, S. Ito, und H. Wada. 1991. Type-2 astrocytes show intracellular Ca²⁺ elevation in response to various neuroactive substances. *Neurosci. Lett.* 128:257-260.
- Irazusta, J., G. Larrinaga, J. Gonzalez-Maeso, J. Gil, J. J. Meana, und L. Casis. 2002. Distribution of prolyl endopeptidase activities in rat and human brain. *Neurochem. Int.* 40:337-345.
- Ishino, T., S. Ohtsuki, K. Homma, und S. Natori. 1998. cDNA cloning of mouse prolyl endopeptidase and its involvement in DNA synthesis by Swiss 3T3 cells. *J. Biochem.(Tokyo)* 123:540-545.
- Jalonen, T. O., R. R. Margraf, D. B. Wielt, C. J. Charniga, M. L. Linne, und H. K. Kimelberg. 1997. Serotonin induces inward potassium and calcium currents in rat cortical astrocytes. *Brain Res.* 758:69-82.

- Johnston, J. A., M. Jensen, L. Lannfelt, B. Walker, und C. H. Williams. 1999. Inhibition of prolylendopeptidase does not affect gamma-secretase processing of amyloid precursor protein in a human neuroblastoma cell line. *Neurosci. Lett.* 277:33-36.
- Kabayama, H., K. Takei, M. Fukuda, K. Ibata, und K. Mikoshiba. 1999. Functional involvement of synaptotagmin I/II C2A domain in neurite outgrowth of chick dorsal root ganglion neuron. *Neuroscience* 88:999-1003.
- Kamei, H., T. Ueki, Y. Obi, Y. Fukagawa, und T. Oki. 1992. Protective effect of eurystatins A and B, new prolyl endopeptidase inhibitors, on scopolamine-induced amnesia in rats. *Jpn. J. Pharmacol.* 60:377-380.
- Kameoka, J., T. Tanaka, Y. Nojima, S. F. Schlossman, und C. Morimoto. 1993. Direct association of adenosine deaminase with a T cell activation antigen, CD26. *Science* 261:466-469.
- Kanematsu, T., I. S. Jang, T. Yamaguchi, H. Nagahama, K. Yoshimura, K. Hidaka, M. Matsuda, H. Takeuchi, Y. Misumi, K. Nakayama, T. Yamamoto, N. Akaike, M. Hirata, und K. Nakayama. 2002. Role of the PLC-related, catalytically inactive protein p130 in GABA(A) receptor function. *EMBO J.* 21:1004-1011.
- Kanematsu, T., Y. Misumi, Y. Watanabe, S. Ozaki, T. Koga, S. Iwanaga, Y. Ikehara, und M. Hirata. 1996. A new inositol 1,4,5-trisphosphate binding protein similar to phospholipase C-delta 1. *Biochem. J.* 313:319-325.
- Kato, A., A. Fukunari, Y. Sakai, und T. Nakajima. 1997. Prevention of amyloid-like deposition by a selective prolyl endopeptidase inhibitor, Y-29794, in senescence-accelerated mouse. *J. Pharmacol. Exp. Ther.* 283:328-335.
- Kato, T., M. Okada, und T. Nagatsu. 1980. Distribution of post-proline cleaving enzyme in human brain and the peripheral tissues. *Mol. Cell Biochem.* 20:117-121.
- Katsube, N., K. Sunaga, H. Aishita, D. M. Chuang, und R. Ishitani. 1999. ONO-1603, a potential antidementia drug, delays age-induced apoptosis and suppresses overexpression of glyceraldehyde-3-phosphate dehydrogenase in cultured central nervous system neurons. *J. Pharmacol. Exp. Ther.* 288:6-13.
- Katsube, N., K. Sunaga, D. M. Chuang, und R. Ishitani. 1996. ONO-1603, a potential antidementia drug, shows neuroprotective effects and increases m3-muscarinic receptor mRNA levels in differentiating rat cerebellar granule neurons. *Neurosci. Lett.* 214:151-154.
- Kovacs, G. L. und D. De Wied. 1994. Peptidergic modulation of learning and memory processes. *Pharmacol. Rev.* 46:269-291.
- Kyhse-Andersen, J. 1884. Electrophoretic transfer of multiple gels: a simple apparatus without buffer tank for rapid transfer of proteins from polyacrylamide to nitrocellulose. *J. Biochem. Biophys. Methods.* 10:203-209.
- Leammler, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature.* 227:680-685
- Laitinen, K. S., T. van Groen, H. Tanila, J. Venalainen, P. T. Mannisto, und I. Alafuzoff. 2001. Brain prolyl oligopeptidase activity is associated with neuronal damage rather than beta-amyloid accumulation. *Neuroreport* 12:3309-3312.
- Loster, K., K. Zeilinger, D. Schuppan, und W. Reutter. 1995. The cysteine-rich region of dipeptidyl peptidase IV (CD 26) is the collagen-binding site. *Biochem. Biophys. Res. Commun.* 217:341-348.
- Luo, W., T. R. Sharif, P. J. Houghton, und M. Sharif. 1997. CGP 41251 and tamoxifen selectively inhibit mitogen-activated protein kinase activation and c-Fos phosphoprotein induction by substance P in human astrocytoma cells. *Cell Growth Differ.* 8:1225-1240.
- Luo, W., T. R. Sharif, und M. Sharif. 1996. Substance P-induced mitogenesis in human astrocytoma cells correlates with activation of the mitogen-activated protein kinase signaling pathway. *Cancer Res.* 56:4983-4991.

- Machleidt W., D.K. Nagler, I. Assfalg-Machleidt, M.T. Stubbs, H. Fritz und E.A. Auerswald. 1995. Temporary inhibition of papain by hairpin loop mutants of chicken cystatin. Distorted binding of the loops results in cleavage of the Gly(9)-Ala10 bond. *FEBS Lett.* 361:185-190.
- Maes, M., F. Goossens, S. Scharpe, J. Calabrese, R. Desnyder, und H. Y. Meltzer. 1995. Alterations in plasma prolyl endopeptidase activity in depression, mania, and schizophrenia: effects of antidepressants, mood stabilizers, and antipsychotic drugs. *Psychiatry Res.* 58:217-225.
- Maes, M., F. Goossens, S. Scharpe, H. Y. Meltzer, P. D'Hondt, und P. Cosyns. 1994. Lower serum prolyl endopeptidase enzyme activity in major depression: further evidence that peptidases play a role in the pathophysiology of depression. *Biol. Psychiatry* 35:545-552.
- Manna, T., T. Sarkar, A. Poddar, M. Roychowdhury, K. P. Das, und B. Bhattacharyya. 2001. Chaperone-like activity of tubulin. binding and reactivation of unfolded substrate enzymes. *J. Biol. Chem.* 276:39742-39747.
- Mantle, D., G. Falkous, S. Ishiura, P. J. Blanchard, und E. K. Perry. 1996. Comparison of proline endopeptidase activity in brain tissue from normal cases and cases with Alzheimer's disease, Lewy body dementia, Parkinson's disease and Huntington's disease. *Clin. Chim. Acta* 249:129-139.
- Marighetto, A., K. Touzani, N. Etchamendy, C. C. Torrea, G. De Nanteuil, D. Guez, R. Jaffard, und P. Morain. 2000. Further evidence for a dissociation between different forms of mnemonic expressions in a mouse model of age-related cognitive decline: effects of tacrine and S 17092, a novel prolyl endopeptidase inhibitor. *Learn. Mem.* 7:159-169.
- Marqueze, B., F. Berton, und M. Seagar. 2000. Synaptotagmins in membrane traffic: which vesicles do the tagmins tag? *Biochimie* 82:409-420.
- Marriott, D. R., G. P. Wilkin, und J. N. Wood. 1991. Substance P-induced release of prostaglandins from astrocytes: regional specialisation and correlation with phosphoinositol metabolism. *J. Neurochem.* 56:259-265.
- Meckelein, B., D. C. Marshall, K. J. Conn, M. Pietropaolo, W. Van Nostrund, und C. R. Abraham. 1998. Identification of a novel serine protease-like molecule in human brain. *Brain Res. Mol. Brain Res.* 55:181-197.
- Mendez, M., C. Cruz, P. Joseph-Bravo, S. Wilk, und J. L. Charli. 1990. Evaluation of the role of prolyl endopeptidase and pyroglutamyl peptidase I in the metabolism of LHRH and TRH in brain. *Neuropeptides* 17:55-62.
- Mentlein, R. 1988. Proline residues in the maturation and degradation of peptide hormones and neuropeptides. *FEBS Lett.* 234:251-256.
- Mentlein, R., P. Dahms, D. Grundt, und R. Kruger. 1993. Proteolytic processing of neuropeptide Y and peptide YY by dipeptidyl peptidase IV. *Regul. Pept.* 49:133-144.
- Mentlein, R. und T. Roos. 1996. Proteases involved in the metabolism of angiotensin II, bradykinin, calcitonin gene-related peptide (CGRP), and neuropeptide Y by vascular smooth muscle cells. *Peptides* 17:709-720.
- Mentlein, R., M. von Kolszynski, R. Sprang, und R. Lucius. 1990. Proline-specific proteases in cultivated neuronal and glial cells. *Brain Res.* 527:159-162.
- Mikoshiba, K., M. Fukuda, K. Ibata, H. Kabayama, und A. Mizutani. 1999. Role of synaptotagmin, a Ca²⁺ and inositol polyphosphate binding protein, in neurotransmitter release and neurite outgrowth. *Chem. Phys. Lipids* 98:59-67.
- Misumi, Y., Y. Hayashi, F. Arakawa, und Y. Ikehara. 1992. Molecular cloning and sequence analysis of human dipeptidyl peptidase IV, a serine proteinase on the cell surface. *Biochim. Biophys. Acta* 1131:333-336.
- Miura, N., S. Shibata, und S. Watanabe. 1995. Increase in the septal vasopressin content by prolyl endopeptidase inhibitors in rats. *Neurosci. Lett.* 196:128-130.

- Mochizuki, Y. und T. Takenawa. 1999. Novel inositol polyphosphate 5-phosphatase localizes at membrane ruffles. *J. Biol. Chem.* 274:36790-36795.
- Morain, P., J. L. Robin, G. De Nanteuil, R. Jochemsen, V. Heidet, und D. Guez. 2000. Pharmacodynamic and pharmacokinetic profile of S 17092, a new orally active prolyl endopeptidase inhibitor, in elderly healthy volunteers. A phase I study. *Br. J. Clin. Pharmacol.* 50:350-359.
- Moriyama, A., M. Nakanishi, und M. Sasaki. 1988. Porcine muscle prolyl endopeptidase and its endogenous substrates. *J. Biochem.(Tokyo)* 104:112-117.
- Nagiec, E. W., K. E. Sampson, und I. Abraham. 2001. Mutated tau binds less avidly to microtubules than wildtype tau in living cells. *J. Neurosci. Res.* 63:268-275.
- Nakajima, T., Y. Ono, A. Kato, J. Maeda, und T. Ohe. 1992. Y-29794--a non-peptide prolyl endopeptidase inhibitor that can penetrate into the brain. *Neurosci. Lett.* 141:156-160.
- Nanri, M. und H. Kaneto. 1987. Anti-amnesic effect of prolyl endopeptidase inhibitors in mice. *Nippon Yakurigaku Zasshi* 89:323-329.
- Nogimori, K., P. J. Hughes, M. C. Glennon, M. E. Hodgson, J. W. Putney, Jr., und S. B. Shears. 1991. Purification of an inositol (1,3,4,5)-tetrakisphosphate 3-phosphatase activity from rat liver and the evaluation of its substrate specificity. *J. Biol. Chem.* 266:16499-16506.
- O'Cuinn, G. 1998. Peptide metabolism in cytoplasm of brain cells. *Biochem. Soc. Trans.* 26:279-292.
- O'Leary, R. M., S. P. Gallagher, und B. O'Connor. 1996. Purification and characterization of a novel membrane-bound form of prolyl endopeptidase from bovine brain. *Int. J. Biochem. Cell Biol.* 28:441-449.
- O'Leary, R. M. und B. O'Connor. 1995. Identification and localisation of a synaptosomal membrane prolyl endopeptidase from bovine brain. *Eur. J. Biochem.* 227:277-283.
- Ohmori, T., T. Nakagami, H. Tanaka, und S. Maruyama. 1994. Isolation of prolylendopeptidase-inhibiting peptides from bovine brain. *Biochem. Biophys. Res. Commun.* 202:809-815.
- Ohtsuki, S., K. Homma, S. Kurata, und S. Natori. 1997. Molecular cloning of cDNA for Sarcophaga prolyl endopeptidase and characterization of the recombinant enzyme produced by an E. coli expression system. *Insect Biochem. Mol. Biol.* 27:337-343.
- Ohtsuki, S., K. Homma, S. Kurata, und S. Natori. 1997. Nuclear localization and involvement in DNA synthesis of Sarcophaga prolyl endopeptidase. *J. Biochem.(Tokyo)* 121:1176-1181.
- Oliver, K. G., J. W. Putney, Jr., J. F. Obie, und S. B. Shears. 1992. The interconversion of inositol 1,3,4,5,6-pentakisphosphate and inositol tetrakisphosphates in AR4-2J cells. *J. Biol. Chem.* 267:21528-21534.
- Piazza, G. A., H. M. Callanan, J. Mowery, und D. C. Hixson. 1989. Evidence for a role of dipeptidyl peptidase IV in fibronectin-mediated interactions of hepatocytes with extracellular matrix. *Biochem. J.* 262:327-334.
- Pittaway, K. M., G. P. Reynolds, und P. C. Emson. 1984. Decreased proline endopeptidase activity in the basal ganglia in Huntington's disease. *J. Neurochem.* 43:878-880.
- Polgar, L. 1991. pH-dependent mechanism in the catalysis of prolyl endopeptidase from pig muscle. *Eur. J. Biochem.* 197:441-447.
- Polgar, L. 1992. Structural relationship between lipases and peptidases of the prolyl oligopeptidase family. *FEBS Lett.* 311:281-284.
- Polgar, L., E. Kollt, und M. Hollosi. 1993. Prolyl oligopeptidase catalysis. Reactions with thiono substrates reveal substrate-induced conformational change to be the rate-limiting step. *FEBS Lett.* 322:227-230.
- Rawlings, N. D., L. Polgar, und A. J. Barrett. 1991. A new family of serine-type peptidases related to prolyl oligopeptidase. *Biochem. J.* 279:907-908.

- Rees, D. C., M. Lewis, R. B. Honzatko, W. N. Lipscomb, und K. D. Hardman. 1981. Zinc environment and cis peptide bonds in carboxypeptidase A at 1.75-Å resolution. *Proc. Natl. Acad. Sci. U.S.A* 78:3408-3412.
- Rennex, D., B. A. Hemmings, J. Hofsteenge, und S. R. Stone. 1991. cDNA cloning of porcine brain prolyl endopeptidase and identification of the active-site seryl residue. *Biochemistry* 30:2195-2203.
- Renzetti, A. R., R. M. Catalioto, C. Carloni, M. Criscuoli, P. Cucchi, A. Giolitti, S. Zappitelli, L. Rotondaro, und C. A. Maggi. 1999. Defects of tyrosine289phenylalanine mutation on binding and functional properties of the human tachykinin NK2 receptor stably expressed in chinese hamster ovary cells. *Biochem. Pharmacol.* 57:899-906.
- Scala-Guenot, D., D. Mouginot, und M. T. Strosser. 1994. Increase of intracellular calcium induced by oxytocin in hypothalamic cultured astrocytes. *Glia* 11:269-276.
- Scanlan, M. J., B. K. Raj, B. Calvo, P. Garin-Chesa, M. P. Sanz-Moncasi, J. H. Healey, L. J. Old, und W. J. Rettig. 1994. Molecular cloning of fibroblast activation protein alpha, a member of the serine protease family selectively expressed in stromal fibroblasts of epithelial cancers. *Proc. Natl. Acad. Sci. U.S.A* 91:5657-5661.
- Schiavo, G., Q. M. Gu, G. D. Prestwich, T. H. Sollner, und J. E. Rothman. 1996. Calcium-dependent switching of the specificity of phosphoinositide binding to synaptotagmin. *Proc. Natl. Acad. Sci. U.S.A* 93:13327-13332.
- Schneider, J. S., M. Giardiniere, und P. Morain. 2002. Effects of the prolyl endopeptidase inhibitor S 17092 on cognitive deficits in chronic low dose MPTP-treated monkeys. *Neuropsychopharmacology* 26:176-182.
- Selcher, J. C., E. J. Weeber, A. W. Varga, J. D. Sweatt, und M. Swank. 2002. Protein kinase signal transduction cascades in mammalian associative conditioning. *Neuroscientist*. 8:122-131.
- Shao, Y. und K. D. McCarthy. 1995. Receptor-mediated calcium signals in astroglia: multiple receptors, common stores and all-or-nothing responses. *Cell Calcium* 17:187-196.
- Shears, S. B. 1998. The versatility of inositol phosphates as cellular signals. *Biochim. Biophys. Acta* 1436:49-67.
- Shinoda, M., A. Miyazaki, und K. Toide. 1999. Effect of a novel prolyl endopeptidase inhibitor, JTP-4819, on spatial memory and on cholinergic and peptidergic neurons in rats with ibotenate-induced lesions of the nucleus basalis magnocellularis. *Behav. Brain Res.* 99:17-25.
- Shinoda, M., K. Toide, I. Ohsawa, und S. Kohsaka. 1997. Specific inhibitor for prolyl endopeptidase suppresses the generation of amyloid beta protein in NG108-15 cells. *Biochem. Biophys. Res. Commun.* 235:641-645.
- Shishido, Y., M. Furushiro, S. Tanabe, S. Shibata, S. Hashimoto, und T. Yokokura. 1999. Effects of prolyl endopeptidase inhibitors and neuropeptides on delayed neuronal death in rats. *Eur. J. Pharmacol.* 372:135-142.
- Spitznagel, H., J. Baulmann, A. Blume, T. Unger, und J. Culman. 2001. C-FOS expression in the rat brain in response to substance P and neurokinin B. *Brain Res.* 16:11-21.
- Takeuchi, H., M. Oike, H. F. Paterson, V. Allen, T. Kanematsu, Y. Ito, C. Erneux, M. Katan, und M. Hirata. 2000. Inhibition of Ca(2+) signalling by p130, a phospholipase-C-related catalytically inactive protein: critical role of the p130 pleckstrin homology domain. *Biochem. J.* 349:357-368.
- Taylor, W. L., P. C. Andrews, C. K. Henrikson, und J. E. Dixon. 1980. New fluorogenic substrates for a rat brain proline endopeptidase. *Anal. Biochem.* 105:58-64.
- Tezuka, Y., W. Fan, R. Kasimu, und S. Kadota. 1999. Screening of crude drug extracts for prolyl endopeptidase inhibitory activity. *Phytomedicine.* 6:197-203.
- Tischmeyer, W. und R. Grimm. 1999. Activation of immediate early genes and memory formation. *Cell Mol. Life Sci.* 55:564-574.

- Toide, K., M. Shinoda, und A. Miyazaki. 1998. A novel prolyl endopeptidase inhibitor, JTP-4819--its behavioral and neurochemical properties for the treatment of Alzheimer's disease. *Rev. Neurosci.* 9:17-29.
- Torrens, Y., M. C. Dagué De Montety, M. el Etr, J. C. Beaujouan, und J. Glowinski. 1989. Tachykinin receptors of the NK1 type (substance P) coupled positively to phospholipase C on cortical astrocytes from the newborn mouse in primary culture. *J. Neurochem.* 52:1913-1918.
- Trotter, E. W., L. Berenfeld, S. A. Krause, G. A. Petsko, und J. V. Gray. 2001. Protein misfolding and temperature up-shift cause G1 arrest via a common mechanism dependent on heat shock factor in *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. U.S.A.* 98:7313-7318.
- Umemura, K., K. Kondo, Y. Ikeda, T. Kobayashi, Y. Urata, und M. Nakashima. 1997. Pharmacokinetics and safety of JTP-4819, a novel specific orally active prolyl endopeptidase inhibitor, in healthy male volunteers. *Br. J. Clin. Pharmacol.* 43:613-618.
- Umemura, K., K. Kondo, Y. Ikeda, M. Nishimoto, Y. Hiraga, Y. Yoshida, und M. Nakashima. 1999. Pharmacokinetics and safety of Z-321, a novel specific orally active prolyl endopeptidase inhibitor, in healthy male volunteers. *J. Clin. Pharmacol.* 39:462-470.
- van Dijken, P., J. C. Bergsma, und P. J. van Haastert. 1997. Phospholipase-C-independent inositol 1,4,5-trisphosphate formation in *Dictyostelium* cells. Activation of a plasma-membrane-bound phosphatase by receptor-stimulated Ca²⁺ influx. *Eur. J. Biochem.* 244:113-119.
- van Dijken, P., J. R. de Haas, A. Craxton, C. Erneux, S. B. Shears, und P. J. van Haastert. 1995. A novel, phospholipase C-independent pathway of inositol 1,4,5-trisphosphate formation in *Dictyostelium* and rat liver. *J. Biol. Chem.* 270:29724-29731.
- Vanhoof, G., F. Goossens, M. De, I. D. Hendriks, und S. Scharpé. 1995. Proline motifs in peptides and their biological processing. *FASEB J.* 9:736-744.
- Walter, R., H. Shlank, J. D. Glass, I. L. Schwartz, und T. D. Kerenyi. 1971. Leucylglycinamide released from oxytocin by human uterine enzyme. *Science* 173:827-829.
- Welches, W. R., K. B. Brosnihan, und C. M. Ferrario. 1993. A comparison of the properties and enzymatic activities of three angiotensin processing enzymes: angiotensin converting enzyme, prolyl endopeptidase and neutral endopeptidase 24.11. *Life Sci.* 52:1461-1480.
- Welihinda, A. A., W. Tirasophon, und R. J. Kaufman. 1999. The cellular response to protein misfolding in the endoplasmic reticulum. *Gene Expr.* 7:293-300.
- Wetzel, R. 1988. Structure, function, nomenclature. *Protein Eng.* 2:1-3.
- Wilk, S. 1983. Prolyl endopeptidase. *Life Sci.* 33:2149-2157.
- Williams, R. S., L. Cheng, A. W. Mudge, und A. J. Harwood. 2002. A common mechanism of action for three mood-stabilizing drugs. *Nature* 417:292-295.
- Williams, R. S., M. Eames, W. J. Ryves, J. Viggars, und A. J. Harwood. 1999. Loss of a prolyl oligopeptidase confers resistance to lithium by elevation of inositol (1,4,5) trisphosphate. *EMBO J.* 18:2734-2745.
- Yang, X., M. Rudolf, M. A. Carew, M. Yoshida, V. Nerreter, A. M. Riley, S. K. Chung, K. S. Bruzik, B. V. Potter, C. Schultz, und S. B. Shears. 1999. Inositol 1,3,4-trisphosphate acts in vivo as a specific regulator of cellular signaling by inositol 3,4,5,6-tetrakisphosphate. *J. Biol. Chem.* 274:18973-18980.
- Yaron, A. 1987. The role of proline in the proteolytic regulation of biologically active peptides. *Biopolymers* 26 Suppl: S215-222.:S215-S222.
- Yaron, A. und D. Mlynar. 1968. Aminopeptidase-P. *Biochem. Biophys. Res. Commun.* 32:658-663.
- Yaron, A. und F. Naider. 1993. Proline-dependent structural and biological properties of peptides and proteins. *Crit Rev. Biochem. Mol. Biol.* 28:31-81.

- Yoshimoto, T., K. Kado, F. Matsubara, N. Koriyama, H. Kaneto, und D. Tsuru. 1987. Specific inhibitors for prolyl endopeptidase and their anti-amnesic effect. *J. Pharmacobiodyn.* 10:730-735.
- Yoshimoto, T., K. Miyazaki, N. Haraguchi, A. Kitazono, T. Kabashima, und K. Ito. 1997. Cloning and expression of the cDNA encoding prolyl oligopeptidase (prolyl endopeptidase) from bovine brain. *Biol. Pharm. Bull.* 20:1047-1050.
- Yoshimoto, T., T. Nishimura, T. Kita, und D. Tsuru. 1983. Post-proline cleaving enzyme (prolyl endopeptidase) from bovine brain. *J. Biochem.(Tokyo)* 94:1179-1190.
- Yoshimura, K., H. Takeuchi, O. Sato, K. Hidaka, N. Doira, M. Terunuma, K. Harada, Y. Ogawa, Y. Ito, T. Kanematsu, und M. Hirata. 2001. Interaction of p130 with, and consequent inhibition of, the catalytic subunit of protein phosphatase 1alpha. *J. Biol. Chem.* 276:17908-17913.
- Zhang, X., J. Rizo, und T. C. Sudhof. 1998. Mechanism of phospholipid binding by the C2A-domain of synaptotagmin I. *Biochemistry* 37:12395-12403.