

## 6 Literature

- (1) Ananiev, E. V., Phillips, R. L. and Rines, H. W. (1998) Chromosome-specific molecular organization of maize (*Zea mays* L.) centromeric regions. *Proc. Natl. Acad. Sci. USA* **95**, 13073-13078.
- (2) Aragón-Alcaide, L., Miller, T., Schwarzacher, T., Reader, S. and Moore, G. (1996) A cereal centromeric sequence. *Chromosoma* **105**, 261-268.
- (3) Armstrong, L. and Snyder, J. A. (1989) Selective reduction of anaphase B in quinacrine-treated PtK-1 cells. *Cell Motil Cytoskel* **14**: 220-229.
- (4) Baum, M., Ngan, V. K. and Clarke, L. (1994) The centromeric K-type repeat and the central core are together sufficient to establish a functional *Schizosaccharomyces pombe* centromere. *Mol. Biol. Cell* **5**: 747-761.
- (5) Belostotsky, D. A. and Ananiev, E. V. (1990) Characterization of relic DNA from barley genome. *Theor. Appl. Genet.* **80**, 374-380.
- (6) Berg, D. E. and Howe, M. M. (1989) Mobile DNA. *Am. Soc. Microbiol.*, Washington, DC.
- (7) Bernard, P., Maure, J.-F., Partridge, J. F., Genier, S., Javerzat, J.-P. and Allshire, R. C. (2001) Requirement of heterochromatin for cohesion at centromeres. *Science* **294**, 2539-2542.
- (8) Bernatzky, R. and Tanksley, S. D. (1986) Genetics of actin-related sequences in tomato. *Theor. Appl. Genet.* **72**, 314-324.
- (9) Boeke, J. (1989) Transposable elements in *Saccharomyces cerevisiae*, pp. 335-374 in Mobile DNA edited by Berg, D. E. and Howe, M. M. (1989)
- (10) Brandes, A., Heslop-Harrison, J. S., Kamm, A., Kubis, S., Doudrick, R. L. and Schmidt, T. (1997) Comparative analysis of the chromosomal and genomic

- organization of Ty1-*copia*-like retrotransposons in pteridophytes, gymnosperms and angiosperms. *Plant Mol. Biol.* **33**, 11-21.
- (11) Brown, M. T. (1995) Sequence similarities between the yeast chromosome segregation protein Mif2 and the mammalian centromere protein CENP-C. *Gene* **160**, 111-116.
- (12) Carmena, M. and Gonzales, C. (1995) Transposable elements map in a conserved pattern of distribution extending from beta-heterochromatin to centromeres in *Drosophila melanogaster*. *Chromosoma* **103**, 676-684.
- (13) Cheng, Z., Dong, F., Langdon, T., Ouyang, S., Buell, C. R., Gu, M., Blattner, F. R. and Jiang, J. (2002) The functional rice centromeres are marked by a satellite repeat and a centromere-specific retrotransposon. *Plant Cell* **14**, 1691-1704.
- (14) Choo, K. H. A., Vissel, B., Nagy, A., Earle, E. and Kalitsis, P. (1991) A survey of the genomic distribution of alpha satellite DNA on all the human chromosomes, and derivation of a new consensus sequence. *Nucleic Acids Res.* **19**, 1179-1182.
- (15) Choo, K. H. A. (1997) The centromere. *Oxford Univ. Press*, Oxford.
- (16) Clarke, L. (1990) Centromeres of budding and fission yeasts. *Trends Genet.* **6**, 150-154.
- (17) Clarke, L. and Carbon, J. (1985) Structure and function of yeast centromeres. *Annu. Rev. Genet.* **19**, 29-56.
- (18) Clarke, L., Baum, M., Marschall, L. G., Ngan, V. K. and Steiner, N. C. (1993) Structure and function of *Schizosaccharomyces pombe* centromeres. *Cold Spring Harb. Symp. Quant. Biol.* **58**, 687-695.

- (19) Dawe, R. K., Reed, L. M., Yu, H.-G., Muszynski, M. G. and Hiatt, E. N. (1999) A maize homolog of mammalian CENPC is a constitutive component of the inner kinetochore. *Plant Cell* **11**, 1227-1238.
- (20) Depinet, T. W., Zackowski, J. L., Earnshaw, W. C., Kaffe, S., Sekhon, G. S., Stallard, R., Sullivan, B. A., Vance, G. H., Dyke, D. L. V., Willard, H. F., Zinn, A. B. and Schwartz, S. (1997) Characterization of neo-centromeres in marker chromosomes lacking detectable alpha-satellite DNA. *Hum. Mol. Genet.* **6**, 1195-1204.
- (21) Dobie, K. W., Hari, K. L., Maggert, K. A. and Karpen, G. H. (1999) Centromere proteins and chromosome inheritance: a complex affair. *Curr. Opin. Genet. Dev.* **9**, 206-217.
- (22) Dong, F., Miller, J. T., Jackson, S. A., Wang, G.-L., Ronald, P. C. and Jiang, J. (1998) Rice (*Oryza sativa*) centromeric regions consist of complex DNA. *Proc. Natl. Acad. Sci. USA* **95**, 8135-8140.
- (23) du Sart, D., Cancilla, M. R., Earle, E., Mao, J.-I., Saffery, R., Tainton, K. M., Kalitsis, P., Martyn, J., Barry, A. E. and Choo, K. H. A. (1997) A functional neo-centromere formed through activation of a latent human centromere and consisting of non-alpha-satellite DNA. *Nature Genet.* **16**, 144-153.
- (24) Earnshaw, W. C. and Rothfield, N. (1985) Identification of a family of human centromere proteins using autoimmune sera from patients with scleroderma. *Chromosoma* **91**, 313-321.
- (25) Earnshaw, W. C. and Bernart, R. L. (1990) Chromosomal passengers: towards an integrated view of mitosis. *Chromosoma* **100**, 139-146.
- (26) Feinberg, A. P. and Vogelstein, B. (1983) A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Analyt Biochem* **132**, 6-13.

- (27) Flavell, A. J., Smith, D. B. and Kumar, A. (1992) Extreme heterogeneity of *Ty1-copia* group retrotransposons in plants. *Mol. Gen. Genet.* **231**, 233-242.
- (28) Fransz, P. F., Alonso-Blanco, C., Liharska, T. B., Peeters, A. J. M., Zabel, P. and de Jong, J. H. (1996) High-resolution physical mapping in *Arabidopsis thaliana* and tomato by fluorescence *in situ* hybridization to extended DNA fibres. *Plant J.* **9**, 421-430.
- (29) Fransz, P. F., Armstrong, S., Alonso-Blanco, C., Fischer, T. C., Torres-Ruiz, R. A. and Jones, G. (1998) Cytogenetics for the model system *Arabidopsis thaliana*. *Plant J.* **13**, 867-876.
- (30) Fransz, P. F., Armstrong, S., de Jong, J. H., Parnell, L. D., van Drunen, C., Dean, C., Zabel, P., Bisseling, T. and Jones, G. H. (2000) Integrated cytogenetic map of chromosome arm 4S of *A. thaliana*: structural organization of heterochromatic knob and centromere region. *Cell* **100**, 367-376.
- (31) Fukagawa, T., Mikami, Y., Nishihashi, A., Regnier, V., Haraguchi, T., Hiraoka, Y., Sugata, N., Todokoro, K., Brown, W. and Toshimichi, I. (2001) CENP-H, a constitutive centromere component, is required for centromere targeting of CENP-C in vertebrate cells. *EMBO J.* **20**, 4603-4617.
- (32) Fukui, K.-N., Suzuki, G., Lagudah, E. S., Rahman, S., Appels, R., Yamamoto, M. and Mukai, Y. (2001) Physical arrangement of retrotransposon-related repeats in centromeric regions of wheat. *Plant Cell Physiol.* **42**, 189-196.
- (33) Georgiev, S., Gecheff, K., Nicoloff, H., Künzel, G. and Rieger, R. (1985) Giemsa N-banding as a tool for identification of chromosome reconstruction in barley. *Biol. Zbl.* **104**: 29-34.
- (34) Gerstel, D. U. and Burns, J. A. (1966) Chromosomes of unusual length in hybrids between two species of *Nicotiana*. *Chromosomes today* **1**: 41-56.

- (35) Gerstel, D. U. and Burns, J. A. (1976) Enlarged euchromatic chromosomes (“megachromosomes”) in hybrids between *Nicotiana tabacum* and *N. plumbaginifolia*. *Genetica* **46**: 139-153.
- (36) Gindullis, F., Desel, C., Galasso, I. and Schmidt, T. (2001) The large-scale organization of the centromeric region in *Beta* species. *Genome Res.* **11**, 253-265.
- (37) Grandbastien, M-A., Lucas, H., Morel, J.-B., Mhiri, C., Vernhettes, S. and Casacuberta, J. M. (1997) The expression of the tobacco Tnt1 retrotransposon is linked to plant defense responses. *Genetica* **100**, 241-252.
- (38) Guldner, H. H., Lakomek, H. J. and Bautz, F. A. (1984) Human anti-centromere sera recognize a 19.5 kd nonhistone chromosomal protein from HeLa cells. *Clin. Exp. Immunol.* **58**, 13-20.
- (39) Harrison, G. E. and Heslop-Harrison, J. S. (1995) Centromeric repetitive DNA sequences in the genus *Brassica*. *Theor. Appl. Genet.* **90**, 157-165.
- (40) Heslop-Harrison, J. S., Brandes, A., Taketa, S., Schmidt, T. Vershinin, A. V. *et al.* (1997) The *Chromosomal* distribution of Ty1-*copia* group retrotransposable elements in higher plants and their implication for genome evolution. *Genetica* **100**, 197-204.
- (41) Heslop-Harrison, J. S., Murata, M., Ogura, Y., Schwarzacher, T. and Motoyoshi, F. (1999) Polymorphisms and genomic organization of repetitive DNA from centromeric regions of *Arabidopsis* chromosomes. *Plant Cell* **11**, 31-42.
- (42) Hieter, P., Pridmore, D., Hegemann, J., Thomas, M., Davis, R. and Phillipsen, P. (1985) Functional selection and analysis of yeast centromeric DNA. *Cell* **42**, 913-921.
- (43) Hoheisel, J. D., Maier, E., Mott, R., McCarthy, L., Grigoriev, A. V., Schalkwyk, L. C., Nizetic, D., Francis, F. and Lehrach, H. (1993) High resolution cosmid

and P1 maps spanning the 14 Mb genome of the fission yeast *S. pombe*. *Cell* **73**, 109-120.

- (44) Houben, A., Brandes, A., Pich, U., Manteuffel, R. and Schubert, I. (1996) Molecular-cytogenetic characterization of a higher plant centromere/kinetochore complex. *Theor. Appl. Genet.* **93**, 477-484.
- (45) Houben, A., Wako, T., Furushima-Shimogawara, R., Presting, G. G., Künzel, G., Schubert, I. and Fukui, K. (1999) The cell cycle dependent phosphorylation of histone H3 is correlated with the condensation of plant mitotic chromosomes. *Plant J.* **18**, 675-679.
- (46) Hoyt, M. A. and Geiser, J. R. (1996) Genetic analysis of the mitotic spindle. *Annu. Rev. Genet.* **30**: 7-33.
- (47) Ijdo, J. W., Wells, R. A., Baldini, A. and Reeders, S. T. (1991) Improved telomere detection using a telomere repeat probe (TTAGGG)<sub>n</sub> generated by PCR. *Nucleic Acids Res* **19**, 4780.
- (48) Jiang, J., Nasuda, S., Dong, F., Scherrer, C. W., Woo, S.-S., Wing, R. A., Gill, B. S. and Ward, D. C. (1996) A conserved repetitive DNA element located in the centromeres of cereal chromosomes. *Proc. Natl. Acad. Sci. USA* **93**, 14210-14213.
- (49) Karpen, G. H. and Allshire, R. C. (1997) The case for epigenetic effects on centromere identity and function. *Trends Genet.* **13**, 489-496.
- (50) Kipling, D., Ackford, H. E., Taylor, B. A. and Cooke, H. J. (1991) Mouse minor satellite DNA genetically maps to the centromere and is physically linked to the proximal telomere. *Genomics* **11**, 235-241.
- (51) Kipling, D., Wilson, H. E., Mitchell, A. R., Taylor, B. A. and Cooke, H. J. (1994) Mouse centromere mapping using oligonucleotide probes that detect variants of the minor satellite. *Chromosoma* **103**, 46-55.

- (52) Kishii, M., Nagaki, K. and Tsujimoto, H. (2001) A tandem repetitive sequence located in the centromeric region of common wheat (*Triticum aestivum*) chromosomes. *Chromosome Res.* **9**, 417-428.
- (53) Konieczny, A., Voytas, D. F., Cummings, M. P. and Ausubel, F. M. (1991) A superfamily of *Arabidopsis thaliana* retrotransposons. *Genetics* **127**, 801-809.
- (54) Kumar, A. and Bennetzen, J. L. (1999) Plant retrotransposons. *Annu. Rev. Genet.* **33**, 479-532.
- (55) Kumekawa, N., Ohtsubo, E. and Ohtsubo, H. (1999) Identification and phylogenetic analysis of *gypsy*-type retrotransposons in the plant kingdom. *Genes Genet. Syst.* **74**, 299-307.
- (56) Kumekawa, N., Ohtsubo, E., Ohtsubo, H., Ohmido, N. and Fukui, K. (2001) A new *gypsy*-type retrotransposon *RIRE7*: preferential insertion into the tandem repeat sequence TrsD in pericentromeric heterochromatin regions of rice chromosomes. *Mol. Genet. Genomics* **265**, 480-488.
- (57) Künzel, G., Korzun, L. and Meister, A. (2000) Cytologically integrated physical restriction fragment length polymorphism maps for the barley genome based on translocation breakpoints. *Genetics* **154**: 397-412.
- (58) Langdon, T., Seago, C., Mende, M., Leggett, M., Thomas, H., Forster, J. W., Thomas, H., Jones, R. N. and Jenkins, G. (2000) Retrotransposon evolution in diverse plant genomes. *Genetics* **156**, 313-325.
- (59) Levis, R., Dunsmuir, P. and Rubin, G.M. (1980) Terminal repeats of the *Drosophila* transposable element *copia*: nucleotide sequence and genomic organization. *Cell* **21**, 581-588.

- (60) Liao, H., Winkfein, R. J., Mack, G., Rattner, J. B. and Yen, T. J. (1995) CENP-F is a protein of the nuclear matrix that assembles onto kinetochores at late G2 and is rapidly degraded after mitosis. *J. Cell Biol.* **130**, 507-518.
- (61) Maluszynska, J. and Heslop-Harrison, J. S. (1991) Localization of tandemly repeated DNA sequences in *Arabidopsis thaliana*. *Plant J.* **1**, 159-166.
- (62) Maney, T., Ginkel, L. M., Hunter, A. W. and Wordeman, L. (1999) The kinetochore of higher eucaryotes: a molecular view. *Int. Rev. Cytol.* **194**, 67-131.
- (63) Manninen, I. and Schulman, A. H. (1992) Analysis of BARE-1, a retrotransposon in barley. *Physiol Plantarum* **85**, A6 33.
- (64) Manuelidis, L. (1978a) Complex and simple sequences in human repeated DNAs. *Chromosoma* **66**, 1-21.
- (65) Manuelidis, L. (1978b) Chromosomal location of complex and simple repeated human dans. *Chromosoma* **66**, 23-32.
- (66) Manzanero, S., Arana, P., Puertas, M. J. And Houben, A. (2000) The chromosomal distribution of phosphorylated histone H3 differs between plants and animals at meiosis. *Chromosoma* **109**, 308-317.
- (67) Martinez-Zapater, J. M., Estelle, M. A. and Somerville, C. R. (1986) A highly repeated DNA sequence in *Arabidopsis thaliana*. *Mol. Gen. Genet.* **204**, 417-423.
- (68) McNeilage, L. J., Whittingham, S., McHugh, N. and Barnett, A. J. (1986) A highly conserved 72 000 dalton centromeric antigen reactive with autoantibodies from patients with progressive systemic sclerosis. *J. Immunol.* **137**, 2541-2547.



- (69) Miller, J. T., Dong, F., Jackson, S. A., Song, J. and Jiang, J. (1998a) Retrotransposon-related DNA sequences in the centromeres of grass chromosomes. *Genetics* **150**, 1615-1623.
- (70) Miller, J. T., Jackson, S. A., Nasuda, S., Gill, B. S., Wing, R. A. and Jiang, J. (1998b) Cloning and characterization of a centromere-specific repetitive DNA element from *Sorghum bicolor*. *Theor. Appl. Genet.* **96**, 832-839.
- (71) Ming, Y. and Hong, M. (2001) Male meiotic spindle lengths in normal and mutant Arabidopsis cells. *Plant Physiology* **126**: 622-630.
- (72) Mitchell, A. R., Gosden, J. R. and Miller, D. A. (1985) A cloned sequence, p82H of the alphoid repeated DNA family found at the centromeres of all human chromosomes. *Chromosoma* **92**, 369-377.
- (73) Mount, S. M. and Rubin, G. M. (1985) Complete nucleotide sequence of the *Drosophila* transposable element *copia*: Homology between *copia* and retroviral proteins. *Mol. Cell. Biol.* **5**, 1630-1638.
- (74) Murphy, T. D. and Karpen, G. H. (1995) Localization of centromere function in a *Drosophila* minichromosome. *Cell* **82**, 599-609.
- (75) Nizetic, D., Drmanac, R. and Lehrach, H. (1991) An improved bacterial colony lysis procedure enables direct DNA hybridization using short (10, 11 bases) oligonucleotides to cosmids. *Nucleic Acids Res* **19**, 182.
- (76) Nonomura, K.-I. and Kurata, N. (1999) Organization of the 1.9-kb repeat unit RCE1 in the centromeric region of rice chromosomes. *Mol. Gen. Genet.* **261**, 1-10.
- (77) Nonomura, K.-I. and Kurata, N. (2001) The centromere composition of multiple repetitive sequences on rice chromosome 5. *Chromosoma* **110**, 284-291.
- O'Hare, K., Alley, M. R., Cullingford, T. E., Driver, A. and Sanderson, M. J. (1991) DNA sequence of the *Doc* retrotransposon in the white-one mutant

- of *Drosophila melanogaster* and of secondary insertion in the phenotypically altered derivatives white-honey and white-eosin. *Mol. Gen. Genet.* **225**, 17-24.
- (78) Pearce, S. R., Harrison, G., Li, D., Heslop-Harrison, J. S., Kumar, A. and Flavell, A. J. (1996) The Ty1-*copia* group retrotransposons in *Vicia* species: copy number, sequence heterogeneity and chromosomal localisation. *Mol. Gen. Genet.* **250**, 305-315.
- (79) Pélissier, T., Tutois, S., Deragon, J. M., Tourmente, S. and Picard, G. (1995) *Athila*, a new retroelement from *Arabidopsis thaliana*. *Plant Mol. Biol.* **29**, 441-452.
- (80) Pélissier, T., Tutois, S., Tourmente, S., Deragon, J. M. and Picard, G. (1996) DNA regions flanking the major *Arabidopsis thaliana* satellite are principally enriched in *Athila* retroelement sequences. *Genetica* **97**, 141-151.
- (81) Pich, U. and Schubert, I. (1998) Terminal heterochromatin and alternative telomeric sequences in *Allium cepa*. *Chromosome Res.* **6**, 315-321.
- (82) Pimpinelli, S., Berloco, M., Fanti, L., Dimitri, P., Bonaccorsi, S., Marchetti, E., Caizzi, R., Caggese, C. and Gatti, M. (1995) Transposable elements are stable structural components of *Drosophila melanogaster* heterochromatin. *Proc. Natl. Acad. Sci. USA* **92**, 3804-3808.
- (83) Presting, G. G., Malysheva, L., Fuchs, J. and Schubert, I. (1998) A TY3/GYPSY retrotransposon-like sequence localizes to the centromeric regions of cereal chromosomes. *Plant J* **16**, 721-728.
- (84) Ramage, R. T. (1971) Reports from coordinators: Translocations and balanced tertiary trisomics. *Barley Genetics Newsletter* **1**, 74-80.

- (85) Ramage, R. T. (1975) Report from coordinator: Translocations and balanced tertiary trisomics. *Barley Genetics Newsletter* **5**, 76-84.
- (86) Rattner, J. B. (1991) The structure of the mammalian centromere. *Bioessays* **13**, 51-56.
- (87) Rattner, J. B., Rao, A., Fritzler, M. J., Valencia, D. W. and Yen, T. J. (1993) CENP-F is a 400 kDa kinetochore protein that exhibits a cell-cycle dependent localization. *Cell Motil. Cytoskeleton* **26**, 214-226.
- (88) Rieder, C. L. and Salmon, E. D. (1998) The vertebrate cell kinetochore and its roles during mitosis. *Trends Cell Biol.* **8**, 310-318.
- (89) Rogowsky, P. M., Shepherd, K. W. and Langridge, P. (1992) Polymerase chain reaction based mapping of rye involving repeated DNA sequences. *Genome* **35**, 621-626.
- (90) Saffery, R., Irvine, D. V., Griffiths, B., Kalitsis, P., Wordeman, L. and Choo, K. H. A. (2000) Human centromeres and neocentromeres show identical distribution patterns of >20 functionally important kinetochore-associated proteins. *Hum. Mol. Genet.* **9**, 175-185.
- (91) Salamini, F., Özkan, H., Brandolini, A., Schäfer-Pregl, R. and Martin, W. (2002) Genetics and geography of wild cereal domestication in the Near East. *Nat. Rev. Genet.* **3**, 429-441.
- (92) SanMiguel, P., Gaut, B. S., Tikhonov, A., Nakajima, Y. and Bennetzen, J. L. (1998) The paleontology of intergene retrotransposons of maize. *Nature Genet.* **20**, 43-45.
- (93) Schmidt, T. and Metzloff, M. (1991) Cloning and characterization of a *Beta vulgaris* satellite DNA family. *Gene* **101**, 247-250.
- (94) Schubert, I. and Oud, J. L. (1997) There is an upper limit of chromosome size for normal development of an organism. *Cell* **88**: 515-520.

- (95) Schubert, I., Oud, J. L. and Pich, U. (1998a) Unscheduled apoptosis in meristematic plant cells is triggered via terminal deletions in artificially elongated chromosome arms. *Theor. Appl. Genet.* **96**: 1022-1026.
- (96) Schubert, I., Shi, F., Fuchs, J. and Endo, T. R. (1998b) An efficient screening for terminal deletions and translocations of barley chromosomes added to common wheat. *Plant J.* **14**: 489-495.
- (97) Schubert, I. (2001) Alteration of chromosome numbers by generation of minichromosomes – Is there a lower limit of chromosome size for stable segregation? *Cytogenet Cell Genet.* **93**: 175-181.
- (98) Simoens, C. R., Gielen, J., Montagu, M. V. and Inze, D. (1988) Characterization of highly repetitive sequences of *Arabidopsis thaliana*. *Nucleic Acids Res.* **16**, 6753-6766.
- (99) Smit, A. F. A (1996) The origin of interspersed repeats in the human genome. *Curr. Opin. Genet. Dev.* **6**, 743-748.
- (100) Smit, A. F. A. (2000) Interspersed repeats and other mementos of transposable elements in mammalian genomes. *Curr. Opin. Genet. Dev.* **9**, 657-663.
- (101) Starr, D. A., Williams, B. C., Li, Z., Etemad-Moghadam, B., Dawe, R. K. and Goldberg, M. L. (1997) Conservation of the centromere/kinetochore protein ZW10. *J. Cell Biol.* **138**, 1289-1301.
- (102) Sugata, N., Li, S., Earnshaw, W. C., Yen, T. J., Yoda, K., Masumoto, H., Munekata, E., Warburton, P. E. and Todokoro, K. (2000) Human CENP-H multimers colocalize with CENP-A and CENP-C at active centromere-kinetochore complexes. *Hum. Mol. Genet.* **9**, 2919-2926.
- (103) Sullivan, W., Daily, D. R., Fogarty, P., Yook, K. J. and Pimpinelli, S. (1993) Delays in anaphase initiation occur in individual nuclei of the syncytial *Drosophila* embryo. *Mol. Biol. Cell* **4**: 885-896.

- (104) Sun, X., Wahlstrom, J. and Karpen, G. (1997) Molecular structure of a functional *Drosophila* centromere. *Cell* **91**, 1007-1019.
- (105) Suonemi, A., Narvanto, A. and Schulman A. H. (1996) The *BARE-1* retrotransposon is transcribed in barley from an LTR promoter active in transient assays. *Plant Mol. Biol.* **31**, 295-306.
- (106) Suonemi, A., Schmidt, D. and Schulman, A. H. (1997) *Bare-1* insertion site preferences and evolutionary conservation of RNA and cDNA preprocessing sites. *Genetica* **100**, 219-230.
- (107) Takahashi, K., Murakami, S., Chikashige, Y., Funabiki, H., Niwa, O. and Yanagida, M. (1992) A low copy number central sequence with strict symmetry and unusual chromatin structure in fission yeast centromere. *Mol. Biol. Cell* **3**, 819-835.
- (108) Taylor, S. S. and McKeon, F. (1997) Kinetochores localization of murine Bub1 is required for normal mitotic timing and checkpoint response to spindle damage. *Cell* **89**, 727-735.
- (109) ten Hoopen, R., Manteuffel, R., Dolezel, J., Malysheva, L. and Schubert, I. (2000) Evolutionary conservation of kinetochore protein sequences in plants. *Chromosoma* **109**, 482-489.
- (110) Tuleen, N. A. and Gardenhire, J. H. (1974) Determination of breakpoint positions in barley translocations by karyotype analysis of permanent rings of six chromosomes. *Can. J. Genet. Cytol.* **16**: 539-548.
- (111) Valdivia, M. and Brinkley, B. R. (1985) Fractionation and initial characterization of the kinetochore from mammalian metaphase chromosomes. *J. Cell Biol.* **101**, 1124-1134.
- (112) Vig, B. K. (1994) Do specific nucleotide bases constitute the centromere? *Mutat. Res.* **309**, 1-10.

- (113) Voytas, D. F. (1996) Retroelements in genome organisation. *Science* **274**, 737-738.
- (114) White, M. J. D. (1973) *Animal cytology and evolution*. Cambridge: Cambridge University Press.
- (115) Willard, H. F. (1985) Chromosome-specific organization of human alpha satellite DNA. *Am. J. Hum. Genet.* **37**, 524-532.
- (116) Yao, X., Abrieu, A., Zheng, Y., Sullivan, K. F. and Cleveland, D. W. (2000) CENP-E forms a link between attachment of spindle microtubules to kinetochores and the mitotic checkpoint. *Nature Cell Biol.* **2**, 484-491.
- (117) Yen, T. J., Compton, D. A., Wise, D., Zinkowski, R. P., Brinkley, R. B., Earnshaw, W. C. *et al.* (1991) CENP-E a novel human centromere-associated protein required for progression from metaphase to anaphase. *EMBO J.* **10**, 1245-1254.
- (118) Yu, Y., Tomkins, J. P., Waugh, R., Frisch, D. A., Kudrna, D., Kleinhofs, A., Brueggeman, R. S., Muehlbauer, G. J., Wise, R. P. and Wing, R. A. (2000) A bacterial artificial chromosome library for barley (*Hordeum vulgare* L.) and the identification of clones containing putative resistance genes. *Theor. Appl. Genet.* **101**, 1093-1099.
- (119) Zhong, C. X., Marshall J. B., Topp, Ch., Mroczek, R., Kato, A., Nagaki, K., Birchler, J. A., Jiang, J. and Dawe, R. K. (2002) Centromeric retroelements and satellites interact with maize kinetochore protein CENH3. *Plant Cell* (in press).