

7. Literaturverzeichnis

- Ali, S., C. R. Müller and J. T. Epplen, 1986: DNA fingerprinting by oligonucleotide probes specific for simple repeats. *Hum. Genet.* 74, 239-243.
- Alonso-Blanco, C., A. J. M. Peeters, M. Kornef, C. Lister, C. Dean, N. van den Bosch, J. Pot and M. T. R. Kuiper, 1998: Development of an AFLP based linkage map of Ler, Col and Cvi *Arabidopsis thaliana* ecotypes and construction of a Ler/Cvi recombinant inbred line population. *Plant J.* 14, 259 - 271.
- Bachem, C. W. B., R. S. van der Hoeven, S. M. de Bruijn, D. Vreugdenhil, M. Zabeau and R. G. F. Visser, 1996: Visualization of differential gene expression using a novel method of RNA fingerprinting based on AFLP: analysis of gene expression during potato tuber development. *Plant J.* 9, 745-753.
- Bai, G., F. L. Kolb, G. Shaner and L. L. Domier, 1999: Amplified Fragment Length Polymorphism Markers Linked to a Major Quantitative Trait Locus Controlling Scab Resistance in Wheat. *Phytopathology* 89, 343-348.
- Barcaccia, G., E. Albertini, S. Tavoletti, M. Falcinelli and F. Veronesi, 1999: AFLP fingerprinting in *Medicago* spp.: its development and application in linkage mapping. *Plant Breeding* 118, 335-340.
- Bell, C. J. and J. R. Ecker, 1994: Assignment of 30 Microsatellite Loci to the Linkage Map of *Arabidopsis*. *Genomics* 19, 137-144.
- Bennett, M. D. and I. J. Leitch, 1995: Nuclear DNA Amounts in Angiosperms. *Annales of Botany* 76, 113-176.
- Bierwerth, S., G. Kahl, F. Weigand and K. Weising, 1992: Oligonucleotide fingerprinting of plant and fungal genomes: a comparison of radioactive, colorigenic and chemiluminescent detection methods. *Electrophoresis* 13, 115-122.
- Bilcescu, O., 1965: Beiträge zur Bestimmung des Fasergehaltes bei Massenuntersuchungen von Hanf bei der Züchtung auf Fasergehalt. *Z. für Pflanzenzüchtg.* 54, 206-224.
- Bócsa, I., 1958: Beiträge zur Züchtung eines ungarischen monözischen Hanfes und zur Kenntnis der Inzuchterscheinungen beim Hanf (*Cannabis sativa* L.). *Z. für Pflanzenzüchtg.* 39, 11-34.

- Bócsa, I., 1995: Die Hanfzüchtung in Ungarn; Zielsetzungen, Methoden und Ergebnisse. In: Tagungsband 1. Biorohstoff Hanf Symposium, Frankfurt/Main, 200-211.
- Bócsa, I. und M. Karus, 1997: Der Hanfanbau: Botanik, Sorten, Anbau und Ernte. 1. Auflage, C. F. Müller Verlag Heidelberg.
- Boerwinkle, E., W. J. Xiong, E. Fourest and L. Chan, 1989: Rapid typing of tandemly repeated hypervariable loci by the polymerase chain reaction: Application to the apolipoprotein B 3' hypervariable region. Proc. Natl. Acad. Sci. USA 86, 212-216.
- Botstein, D., R. L. White, M. Skolnick and R. W. Davies, 1980: Construction of a genetic linkage map in man using restriction fragment length polymorphisms. Am. J. Hum. Genet. 32, 314-331.
- Bracale, M., E. Caporali, M. G. Galli, C. Longo, G. Marziani-Longo, G. Rossi, A. Spada, C. Soave, A. Falavigna, F. Raffaldi, E. Maestri, F. M. Restivo and F. Tassi, 1991: Sex determination and differentiation I. *Asparagus officinalis* L.. Plant Science 80, 67-77.
- Brady, J. M., N. S. Scott and M. R. Thomas, 1996: DNA typing of hops (*Humulus lupulus*) through application of RAPD and microsatellite marker sequences converted to sequence tagged sites (STS). Euphytica 91, 277-284.
- Bredemann, G., 1922a: Die Bestimmung des Fasergehaltes in Bastfaserpflanzen bei züchterischen Untersuchungen. Faserforschung 2, 239-259.
- Bredemann, G., 1922b: Über Faserausbeutebestimmungen bei Hanfzüchtung. Angew. Botanik 4, 223-233.
- Bredemann, G., 1924: Beiträge zur Hanfzüchtung: Auslese faserreicher Männchen zur Befruchtung durch Faserbestimmung an der lebenden Pflanze vor der Blüte. Angew. Botanik 6, 348-360.
- Bredemann, G., 1927: Beiträge zur Hanfzüchtung. III. Weitere Versuche zur Züchtung auf Fasergehalt. Z. für Pflanzenzüchtg. 12, 259-268.
- Bredemann, G., 1940: Fasergehalt und Faseausbeute beim Hanf in verschiedener Stengelhöhe. Forschungsdienst 10, 57-67.
- Bredemann, G., 1942a: Die Bestimmung des Fasergehaltes bei Massenuntersuchungen von Hanf, Flachs, Fasernesseln und anderen Bastfaserpflanzen. Faserforschung 16, 14-39.

- Bredemann, G., 1942b: Züchtung auf Fasergehalt bei Hanf (*Cannabis sativa* L.). Der Züchter 14, 201-213.
- Bredemann, G., 1952: Weitere Beobachtungen bei Züchtung des Hanfes auf Fasergehalt. Der Züchter 22, 257-269.
- Bredemann, G., K. Garber, P. Harteck and K. A. Suhr, 1947: Die Temperaturabhängigkeit der Lebensdauer von Blütenpollen. Die Naturwissenschaften 9, 279-280.
- Bredemann, G., K. Garber, W. Huhnke, und R. von Sengbusch, 1961: Die Züchtung von monözischen und diözischen, faserertragsreichen Hanfsorten Fibrimon und Fibridia. Z. für Pflanzenzüch. 46, 235-245.
- Breslavetz, L., 1928: Polyploide Mitosen bei *Cannabis sativa* L.. I. Ber. dtsh. bot. Ges., 644-649.
- Breslavetz, L., 1932: Polyploide Mitosen bei *Cannabis sativa* L. II. Planta 17, 644-649.
- Bruford, M. W. and R. K. Wayne, 1993: Microsatellites and their application to population genetic studies. Curr. Biol. 3, 939-943.
- Caetano-Anollés, G., B. J. Bassam and P. M. Gresshoff, 1991: High resolution DNA amplification fingerprinting using very short arbitrary oligonucleotide primers. Biotechnology 9, 553-557.
- Charlesworth, D., 2002: Plant sex determination and sex chromosomes. Heredity 88, 94-101.
- Charmet, G., P. F. Bert and F. Balfourier, 2000: A computerised algorithm for selecting a subset of multiplex molecular markers and optimising linkage map construction Theor. Appl. Genet. 101, 90-94.
- Cipriani, G., G. Lot, W.-G. Huang, M. T. Marrazzo, E. Peterlunger and R. Testolin, 1999: AC/GT and AG/CT microsatellite repeats in peach [*Prunus persica* (L) Batsch]: isolation, characterisation and cross-species amplification in *Prunus*. Theor. Appl. Genet. 99, 65-72.
- Cnops, G., B. den Boer, A. Gerats, M. Van Montagu and M. Van Lijsebettens, 1996: Chromosome landing at the *Arabidopsis* *TORNADO1* locus using an AFLP-based strategy. Mol. Gen. Genet. 253, 32-41.
- Coen, E. S. and E. M. Meyerowitz, 1991: The war of the whorls: genetic interactions controlling flower development. Nature 353, 31-37.

- Collevatti, R. G., R. V. Brondani and D. Grattapaglia, 1999: Development and characterisation of microsatellite markers for genetic analysis of a Brazilian endangered species *Caryocar brasiliens*. *Heredity* 83, 748-756.
- Costa, P., D. Pot, C. Dubos, J. M. Frigerio, C. Pionneau, C. Bodénés, E. Bertocchi, M.-T. Cervera, D. L. Remington and C. Plomion, 2000: A genetic map of Maritime pine based on AFLP, RAPD and protein markers. *Theor. Appl. Genet.* 100, 39-48.
- Crescini, F., 1940: Formen von Hanf (*Cannabis sativa* L.). *Der Züchter* 12, 105-115.
- Debener, T. and L. Mattiesch, 1999: Construction of a genetic linkage map for roses using RAPD and AFLP markers. *Theor. Appl. Genet.* 99, 891-899.
- Debruyne, D., M. Moulin, M.-C. Bigot and R. Camsonne, 1981: Identification and differentiation of resinous *cannabis* and textile *cannabis*: combined use of HPLC and high-resolution GLC. *Bulletin on Narcotics* 2, 49-58.
- Decousset, L., S. Griffiths, R. P. Dunford, N. Pratchett and D. A. Laurie, 2000: Development of STS markers closely linked to the *Ppd-H1* photoperiod response gene of barley (*Hordeum vulgare* L.). *Theor. Appl. Genet.* 101, 1202-1206.
- Dellaporta, S. L. and A. Calderon-Urrea, 1993: Sex determination in flowering plants. *Plant Cell* 5, 1241-1251.
- de Meijer, E. P. M., H. J. van der Kamp and F. A. van Eeuwijk, 1992: Characterisation of *Cannabis* accessions with regard to cannabinoid content in relation to other plant characters. *Euphytica* 62, 187-200.
- Dettori, M. T., R. Quarta and I. Verde, 2001: A peach linkage map integrating RFLPs, SSRs, RAPDs, and morphological markers. *Genome* 44, 783-790.
- Di Stilio, V. S., R. V. Kesseli and D. L. Mulcahy, 1998: A Pseudautosomal Random Amplified Polymorphic DNA Marker for the Sex Chromosomes of *Silene dioica*. *Genetics* 149, 2057-2062.
- Don, R. H., P. T. Cox, B. J. Wainwright, K. Baker and J. S. Mattick, 1991: "Touchdown" PCR to circumvent spurious priming during gene amplification. *Nuc. Ac. Res.* 19, 4008.
- D'Ovidio, R., O. A. Tanzarella, A. Cenci, E. Iacono and E. Porceddu, 1994: RFLP analysis in wheat. Isolation and chromosomal assignment of digoxigenin-labelled clones. *J. Genet. Breed.* 48, 73-80.
- Dumas, V., S. Herder, A. Bebbia, C. Cadoux-Barnabé, C. Bellec and G. Cuny, 1998: Polymorphic microsatellites in *Simulium damnosum* s.l. and their use for differentiating two savannah populations: implications for epidemiological studies. *Genome* 41: 154-161.

- Dusabenyagasani, M., N. Lecours and R. C. Hamelin, 1998: Sequence-tagged sites (STS) for studies of molecular epidemiology of scleroderris canker of conifers. *Theor. Appl. Genet.* 97, 789-796.
- El-Ghany, M. E. A., 2001: Molekulargenetische Diversität einer monözischen und einer diözischen Hanfsorte und Analyse des Fasergehaltes von verschiedenen Hanfformen (*Cannabis sativa* L.). Dissertation, Martin-Luther-Universität Halle-Wittenberg.
- Faeti, V., G. Mandolino and P. Ranalli, 1996: Genetic diversity of *Cannabis sativa* germplasm based on RAPD markers. *Plant Breeding* 115, 367-370.
- Fischer, D. and K. Bachmann, 1998: Microsatellite Enrichment in Organisms with Large Genomes (*Allium cepa* L.). *BioTechniques* 24, 796-802.
- Flachowsky, H. 2001: Entwicklung und Kartierung geschlechtsspezifischer Marker bei Hanf (*Cannabis sativa* L.). *Vortr. Pflanzenzüchtung* 52: 121-124.
- Flachowsky, H., E. Schumann, W. E. Weber and A. Peil, 2001: Application of AFLP for the detection of sex specific markers in hemp (*Cannabis sativa* L.). *Plant Breeding* 120: 305-309.
- Fladung, M. and M. Rai Ahuja, 1995: Sandwich method for nonradioactive hybridization. *BioTechniques* 18, 254-256.
- Forapani, S., A. Carboni, C. Paoletti, V. M. C. Moliterni, P. Ranalli and G. Mandolino, 2001: Comparison of Hemp Varieties Using Random Amplified Polymorphic DNA Markers. *Crop Sci.* 41, 1682-1689.
- Fruwirth, C., 1922: Zur Hanfzüchtung. *Z. für Pflanzenzüchtg.* 8, 340-401.
- Gasser, R. B., X. Zhu and W. Woods, 1999: Genotyping *Taenia* tapeworms by single-strand confirmation polymorphism of mitochondrial DNA. *Electrophoresis* 20, 2834-2837.
- Gigliano, G. S., 1999: Preliminary Data on the usefulness of internal transcribed spacer I (ITS1) sequence in *Cannabis sativa* L. identification. *J. Forensic Sci.* 44, 475-477.
- Gu Cho, Y., M. W. Blair, O. Panaud and S. R. McCouch, 1996: Cloning and mapping of variety-specific rice genomic DNA sequences: amplified fragment length polymorphisms (AFLP) from silver-stained polyacrylamide gels. *Genome* 39, 373-378.
- Guillemaut, P. and L. Maréchal-Drouard, 1992: Plant DNA isolation: a fast, inexpensive and reliable method. *Plant Mol. Biol. Rep.* 10: 60-65.
- Guttman, D. S. and D. Charlesworth, 1998: An X-linked gene with a degenerate Y-linked homologue in a dioecious plant. *Nature* 393, 263-266.
- Handke, S., 1995: Die Züchtung von monözischen und diözischen faserreichen Hanfsorten. In: *Tagungsband 1. Biorohstoff Hanf Symposium, Frankfurt/Main, 188-1999.*

- Hanf, C.-H., 1997: Hanf. Teil 2: Wissenschaftliche Untersuchung zum Thema "Anbau, Ernte und Aufbereitung sowie Verwendung von Hanf". Landwirtschaftsverlag Münster.
- Hartl, L. and S. Seefelder, 1998: Diversity of selected hop cultivars detected by fluorescent AFLP's. *Theor. Appl. Genet.* 96, 112-116.
- Harvey, C. F., G. P. Gill, L. G. Fraser and M. A. McNeilage, 1997: Sex determination in *Actinidia*. 1. Sex-linked markers and progeny sex ratio in diploid *A. chinensis*. *Sexual Plant Reproduction* 10, 149-154.
- Harvey, S. C. and M. E. Viney, 2001: Sex determination in the parasitic nematode *Strongyloides ratti*. *Genetics* 158, 1527-1533.
- Hawthorne, D. J., 2001: AFLP-based genetic linkage map of the Colorado potato beetle *Leptinotarsa decemlineata* sex chromosomes and a pyrethroid-resistance candidate gene. *Genetics* 158, 695-700.
- Herich, R., 1961: Geschlechtsausprägung und Größe der Zellkerne sowie Polkörper bei monözischem Hanf verschiedener Formen. *Der Züchter* 31, 48-51.
- Heslop-Harrison, J., 1956: Auxin and Sexuality in *Cannabis sativa*. *Physiol. Plant.* 9, 588-597.
- Heuser, O. 1924: Der deutsche Hanf. Verlag Hirzel, Leipzig.
- Heuser, O., 1927: Hanf- und Hartfasern. In: Technologie der Textilfasern, Band 5 (Teil 2), Axel Springer-Verlag, Berlin.
- Hirata, K., 1929: Sex determination in hemp *Cannabis sativa* L. *J. Genet.* 19, 65- 69.
- Hitzemann, W., 1941: Untersuchungen auf "Haschisch" bei verschiedenen Hanfsorten eigenen Anbaus in Deutschland. *Arch. Pharmazie* 279, 353-387.
- Hoffmann, W., 1938: Das Geschlechtsproblem des Hanfes in der Züchtung. *Z. für Pflanzenzüchtg.* 22, 453-461.
- Hoffmann, W., 1941: Gleichzeitig reifender Hanf. *Der Züchter* 13, 277-283.
- Hoffmann, W., 1947a: Die Vererbung der Geschlechtsformen des Hanfes (*Cannabis sativa* L.). *Der Züchter* 17/18, 257-277.
- Hoffmann, W., 1947b: Helle Stengel - eine wertvolle Mutation des Hanfes (*Cannabis sativa* L.). *Der Züchter* 17/18, 56-59.
- Hoffmann, W. and E. Knapp, 1940: Röntgenstrahlungen beim Hanf. *Der Züchter* 12, 1-9.
- Hoffmann, W., A. Mudra, und W. Plarre, 1970: Lehrbuch der Züchtung Landwirtschaftlicher Kulturpflanzen, Band 2, Spezieller Teil, Paul Parey Verlag, Berlin/Hamburg, 415-430. Hoffmann, W., A. Mudra und W. Plarre, 1985: Lehrbuch der Züchtung Landwirtschaftlicher Kulturpflanzen, Band 2, Spezieller Teil, Paul Parey Verlag, Berlin/Hamburg, 344-349.

- Höppner, F. und U. Menge-Hartmann, 1996: Organspezifische Entwicklung der Δ^9 -Tetrahydrocannabinol (THC)- und Cannabidiol (CBD)-Konzentration während der Vegetationsperiode zweier Faserhanfsorten. *Landbauforschung Völkenrode* 2, 55-64.
- Hornig, J. M. and M. C. Huang, 2000: Male-Specific Band in Random Amplified Microsatellite Polymorphism Fingerprints of Holstein Cattle. *Proc. Natl. Acad. Sci. USA* 24, 41-46.
- Huhnke, W., Ch. Jordan, H. Neuer und R. von Sengbusch, 1951: Grundlagen für die Züchtung eines monözischen Hanfes. *Z. für Pflanzenzüchtg.* 29, 55-75.
- Imai G., 1938: Sex linked mutant characters in the hemp (*Cannabis sativa*). *Journal of Genetics* 35, 431-432.
- Jagdish, V., J. Robertson and A. Gibbs, 1996: RAPD analysis distinguishes *Cannabis sativa* samples from different sources. *Forensic Sci. Int.* 79, 113-121.
- Jakše, J. and B. Javornik, 2001: High Throughput Isolation of Microsatellite in Hop (*Humulus lupulus* L.). *Plant Mol. Biol. Rep.* 19, 217-226.
- Jakše, J., K. Kindlhofer and B. Javornik, 2001: Assessment of genetic variation and differentiation of hop genotypes by microsatellite and AFLP markers. *Genome* 44, 773-782.
- Jeffreys, A. J., V. Wilson and S. L. Thein, 1985: Hypervariable "minisatellite" regions in human DNA. *Nature* 314, 67-73.
- Jeffreys, A. J., A. MacLeod, K. Tamaki, D. L. Neil and D. G. Monckton, 1991: Minisatellite repeat coding as a digital approach to DNA typing. *Nature* 354, 204-209.
- Jeppsson, N., I. V. Bartish and H. A. Persson, 1999: DNA Analysis as a Tool in Sea Buckthorn Breeding. Reprinted from: *Perspectives on new crops and new uses*, J. Janick (ed.), ASHS Press, Alexandria, VA.
- Jiang, C. and K. C. Sink, 1997: RAPD and SCAR markers linked to the sex expression locus *M* in *asparagus*. *Euphytica* 94, 329-333.
- Jones, C. J., K. J. Edwards, S. Castaglione, M. O. Winfield, F. Sale, C. Van de Wiel, G. Bredemeijer, M. Buiatti, E. Maestri, A. Malcevshi, N. Marmioli, R. Aert, G. Volckaert, J. Rueda, R. Linacero, A. Vazquez and A. Karp, 1997: Reproducibility testing of RAPD, AFLP and SSR markers in plants by a network of European laboratories. *Mol. Breed.* 3, 381-390.
- Kaemmer, D., K. Weising, G. Kahl, B. Beyermann, T. Börner and J. T. Epplen, 1995: Oligonucleotide fingerprinting of tomato DNA. *Plant Breeding* 114, 12-17.

- Kaneko, T., H. Kotani, Y. Nakamura, S. Sato, E. Asamizu, N. Miyajima and S. Tabata, 1998: Structural analysis of *Arabidopsis thaliana* chromosome 5. V. Sequence features of the regions of 1,381,565 bp covered by twenty one physically assigned P1 and TAC clones. *DNA Res.* 5, 131-45.
- Karp, A., K. J. Edwards, M. Bruford, B. Vosman, M. Morgante, O. Seberg, A. Kremer, P. Boursot, P. Arctander, D. Tautz and G. Hewitt, 1997: Molecular technologies for biodiversity evaluation: Opportunities and challenges. *Nature Biotechnol.* 15, 625-628.
- Kater, M. M., J. Franken, K. J. Carney, L. Colombo and G. C. Angenent, 2001: Sex determination in the monoecious species cucumber is confined to specific floral whorls. *Plant Cell* 13, 481-493.
- Keijnovský, E., J. Vrána, S. Matsunaga, P. Souček, J. Šíroky, J. Doležel and B. Vyskot, 2001: Localization of Male-Specifically Expressed *MROS* Genes of *Silene latifolia* by PCR on Flow-Sorted Sex Chromosomes and Autosomes. *Genetics* 158, 1269-1277.
- Kim, E. S. and P. G. Mahlberg, 1997: Immunochemical localization of tetrahydrocannabinol (THC) in cryofixed glandular trichomes of *Cannabis* (*Cannabaceae*). *Amer. J. Bot.* 83, 336-342.
- Köhler, D., 1958: Zur Vererbung der Monözie beim Hanf. *Zeitschrift für Vererbungslehre* 89, 437-447.
- Köhler, D., 1961: Ein Beitrag zur Physiologie und Genetik der Geschlechtsausprägung von *Cannabis sativa*. *Planta* 56, 150-173.
- Konieczny, A. and F. M. Ausubel, 1993: A procedure for mapping *Arabidopsis* mutations using co-dominant ecotype-specific PCR-based markers. *Plant J.* 4, 403-410.
- Kotoda, N., M. Wada, S. Komori, S. Kidou, K. Abe, T. Masuda and J. Soejima, 2000: Expression pattern of homologues of floral meristem identity gene *LFY* and *API* during flower development in apple. *J. Amer. Soc. Hort. Sci.* 125, 398-403.
- Kovács, B., S. Egedi, R. Bártfai and L. Orbán, 2001: Male-specific DNA markers from African catfish (*Clarias gariepinus*). *Genetica* 110, 267-276.
- Kühl, H., S. Röttger, H. Heilbronner, H. Enders and W. Schempp, 2001: Loss of the Y chromosomal PAR2 region in four familial cases of satellited Y chromosomes Yqs. *Chromosome Research* 9, 215-222.
- Kumar, L. S. S. and S. Vishveshwaralah, 1952: Sex mechanism in *Coccinea indica*. *Nature* 170, 330-331.

- Lahaye, T., B. Rueger, S. Toepsch, J. Thalhammer and P. Schulze-Lefert, 1996: Detection of single-copy sequences with Digoxigenin-labeled probes in a complex plant genome after separation on pulsed-field gels. *BioTechniques* 21, 1067-1072.
- Lahn, B. T. and D. C. Page, 1999: Four Evolutionary Strata on the Human X Chromosome. *Science* 286, 964-967.
- Lahn, B. T., N. M. Pearson and K. Jegalian, 2001: The human Y chromosome, in the light of evolution. *Nature Genet. Reviews* 2, 207-216.
- Lander, E. S. and N. J. Schork, 1994: Genetic dissection of complex traits. *Science* 265, 2037-2048.
- Lashermes, P., M. C. Combes, N. S. Prakash, P. Trouslot, M. Lorieux and A. Charrier, 2001: Genetic linkage map of *Coffea canephora*: effect of segregation distortion and analysis of recombination rate in male and female meioses. *Genome* 44, 589-596.
- Lebel-Hardenack, S., E. Hauser, T. Law, J. Schmid and S. R. Grant, 2002: Mapping of Sex Determination Loci on the White Campion (*Silene latifolia*) Y Chromosome Using Amplified Fragment Length Polymorphism. *Genetics* 160: 717-725.
- Lemmetyinen, J., T. Pennanen, M. Lännenpää and T. Sopanen, 2001: Prevention of flower formation in dicotyledonous. *Mol. Breed.* 7, 341-350.
- Lespinasse, D., M. Rodier-Goud, L. Grivet, A. Leconte, H. Legnate and M. Seguin, 2000: A saturated genetic linkage map of rubber tree (*Hevea* spp.) based on RFLP, AFLP, microsatellite and isozyme markers. *Theor. Appl. Genet.* 100, 127-138.
- Liang, P. and A. B. Pardee, 1992: Differential display of eukaryotic messenger RNA by means of the polymerase chain reaction. *Science* 257, 967-971.
- Lin, T. Y., Y. Y. Kao, S. Lin, R. F. Lin, C. M. Chen, C. H. Huang, C. K. Wang, Y. Z. Lin and C. C. Chen, 2001: A genetic linkage map of *Nicotiana plumbaginifolia*/*Nicotiana longiflora* based on RFLP and RAPD markers. *Theor. Appl. Genet.* 103, 905-911.
- Linacre, A. and J. Thorpe, 1998: Detection and identification of *Cannabis* by DNA. *Forensic Sci. Int.* 91, 71-76.
- Linn, S. and S. Arber, 1968: A restriction enzyme from *Hemophilus influenzae*. I. Purification and general properties. *Proc. Natl. Acad. Sci. USA* 59, 1300.
- Linnert, G., 1991: Geschlechtschromosomen bei Pflanzen. *Lehrbuch der Allgemeinen Cytogenetik*. Paul Paray Verlag, Hamburg, 78-80.
- Litt, M. and J. A. Luty, 1989: A hypervariable microsatellite revealed by *in vitro* amplification of a dinucleotide repeat within the cardiac muscle actin gene. *Am. J. Hum. Genet.* 44, 397-401.

- Loch, L., 1995: Erfahrungen bei der Züchtung eines einhäusigen Hanfes. In: Tagungsband 1. Biorohstoff Hanf Symposium, Frankfurt/Main, 170-187.
- Lotti, C., S. Salvi, A. Pasqualone, R. Tuberosa and A. Blanco, 2000: Integration of AFLP markers into an RFLP-based map of durum wheat. *Plant Breeding* 119, 393-401.
- Loxdale, H. D. and G. Lushai, 1998: Molecular markers in entomology. *Bulletin of Entomol. Research* 88, 577-600.
- Lu, Z. X., K. Sossey-Alaoui, G. L. Reighard, W. V. Baird and A. G. Abbott, 1999: Development and characterization of a codominant marker linked to root-knot nematode resistance and its application to peach rootstock breeding. *Theor. Appl. Genet.* 99, 115-122.
- Mackay, E. L., 1939: Sex chromosomes of *Cannabis sativa*. *Amer. J. Bot.* 26, 707-708.
- Magome, H., N. Yoshikawa and T. Takahashi, 1999: Single-Strand Confirmation Polymorphism Analysis of Apple Stem Grooving Capillovirus Sequence Variants. *Phytopathology* 89, 136-140.
- Maheswaran, M., P. K. Subudhi, S. Nandi, J. C. Xu, A. Parco, D. C. Yang and N. Huang, 1997: Polymorphism, distribution, and segregation of AFLP markers in a doubled haploid rice population. *Theor. Appl. Genet.* 94, 39-45.
- Maliepaard, C., F. H. Alston, G. van Arkel, L. M. Brown, E. Chevreau, F. Dunemann, K. M. Evans, S. Gardiner, P. Guilford, A. W. van Heusden, J. Janse, F. Laurens, J. R. Lynn, A. G. Manganaris, A. P. M. den Nijs, N. Periam, E. Rikkerink, P. Roche, C. Ryder, S. Sansavini, H. Schmidt, S. Tartarini, J. J. Verhaegh, M. Vrielink-van Ginkel and G. J. King, 1998: Aligning male and female linkage maps of apple (*Malus pumila* Mill.) using multi-allelic markers. *Theor. Appl. Genet.* 97, 60-73.
- Mandel, M. A. and M. F. Yanofsky, 1995: A gene triggering flower formation in *Arabidopsis*. *Nature* 377, 522-524.
- Mandolino, G. and P. Ranalli, 2002: The Applications of Molecular Markers in Genetics and Breeding of Hemp. *Journal of Industrial Hemp* 7, 7-24.
- Mandolino, G., A. Carboni, S. Forapani, V. Faeti and P. Ranalli, 1999: Identification of DNA markers linked to the male sex in dioecious hemp (*Cannabis sativa* L.). *Theor. Appl. Genet.* 98, 86-92.
- Mansfeld, R., 1986: Verzeichnis landwirtschaftlicher und gärtnerischer Kulturpflanzen (ohne Zierpflanzen). 2 Aufl., Axel Springer Verlag, Berlin.
- Marec, F., A. Tothová, K. Sahara and W. Traut, 2001: Meiotic pairing of sex chromosome fragments and its relation to atypical transmission of a sex-linked marker in *Ephesia kuehniella* (Insecta, Lepidoptera). *Heredity* 87, 659-671.

- Matsunaga, S. and S. Kawano S., 2001: Sex determination by sex chromosomes in dioecious plants. *Plant Biology* 3, 1-8.
- Matsunaga, S., S. Kawano and T. Kuroiwa, 1997: A male stamen-specific gene in the dioecious campion *Silene latifolia* expressed in mature pollen. *Plant Cell Physiol.* 38, 499-502.
- Matsunaga, S., S. Kawano, H. Takano, H. Uchida and A. Sakai, 1996: Isolation and developmental expression of male reproductive organ-specific genes in a dioecious campion, *Melandrium album* (*Silene latifolia*). *Plant J.* 10, 679-689.
- Matthäus, B., L. Brühl, U. Krise, E. Schumann and A. Peil, 2001: Hanföl: Ein "Highlight" für die Küche? *Forschungsreport* 2, 22-25.
- Mc Phee, H., 1925: The genetics of sex in hemp. *J. Agric. Res.* 31, 935-943.
- Mekada, K., M. Harada, L. K. Lin, K. Koyasu, P. M. Borodin and S. I. Oda, 2001: Pattern of X-Y chromosome pairing in the Taiwan vole, *Microtus kikuchii*. *Genome* 44, 27-31.
- Meksem, K., D. Leister, J. Peleman, M. Zabeau, S. F. and C. Gebhardt, 1995: A high-resolution map of the vicinity of the R1 locus on chromosome V of potato based on RFLP and AFLP markers. *Mol. Gen. Genet.* 249, 74-81.
- Meksem, K., E. Ruben, D. Hyten, K. Triwitayakorn and A. D. Lightfoot, 2001: Conversion of AFLP bands to high-throughput DNA markers. *Mol. Gen. Genomics* 265, 207-214.
- Mendel, G., 1865: Versuche über Pflanzen-Hybriden. *Verhandlungen des naturforschenden Vereines in Brünn* 4, 3-44.
- Meyerowitz, E. M., 1995: Die Genetik der Blütenentwicklung. *Spektrum der Wissenschaft*, 42-49.
- Michelmore, R., I. Paran and R. V. Kesseli, 1991: Identification of marker linked to disease resistance gene by bulk segregant analysis: a rapid method to detect markers in specific genomic regions using segregating populations. *Proc. Natl. Acad. Sci. USA* 88, 9828-9832.
- Mohan Ram, H. Y. and V. S. Jaiswal, 1972: Induction of Male Flowers on Female Plants of *Cannabis sativa* by Gibberellins and its Inhibition by Abscisic Acid. *Planta* 105, 263-266.
- Morgan, T. H., C. B. Bridges and A. H. Sturtevant, 1925: Genetics of Drosophila. *Bibliogr. Genetica* 2, 1-262.
- Mueller, U. G., L. La. R. Wolfenbarger, 1999: AFLP genotyping and fingerprinting. *Tree* 14, 389-394.
- Mullis, K. and F. Faloona, F., 1987: Specific synthesis of DNA *in vitro* via a polymerase-catalyzed chain reaction. *Methods Enzymol.* 155, 335-350.

- Mullis, K. B., F. A. Faloona, S. J. Scharf, R. K. Saiki, G. T. Horn and H. A. Ehrlich, 1986: Specific enzymatic amplification of DNA *in vitro*: the polymerase chain reaction. Cold Spring Harbor Symp. Quant. Biol. 51: 263-273.
- Nakao, S., S. Matsunaga, A. Sakai, T. Kuroiwa and S. Kawano, 2002: RAPD isolation of a Y chromosome specific ORF in a dioecious plant, *Silene latifolia*. Genome 45, 413-20.
- Negi, M. S., M. Devic, M. Delseny and M. Lakshmikumaran, 2000: Identification of AFLP fragments linked to seed coat colour in *Brassica juncea* and conversion to a SCAR marker for rapid selection. Theor. Appl. Genet. 101, 146-152.
- Neuer, H. und R. von Sengbusch, 1943: Die Geschlechtsvererbung bei Hanf und die Züchtung eines monözischen Hanfes. Der Züchter 15, 49-62.
- Neuer, H., E. Prieger und R. von Sengbuch, 1946: Hanfzüchtung. I. Die Steigerung des Faserertrages von Hanf. Der Züchter 17/18, 33-39.
- Neuhaus-Url, G. and G. Neuhaus, 1993: The use of the nonradioactive digoxigenin chemiluminescent technology for plant genomic Southern blot hybridization: a comparison with radioactivity. Transgenic Research 2, 115-120.
- Nishina, Y., Y. Sinoto and D. Sato, 1940: Effects of fast neutrons upon plants. 3. cytological observations on the abnormal forms of *Fagopyrum* and *Cannabis*. Cytology (Tokyo) 10, 458-466.
- Nishiyama, I., 1940: Studien über künstlich erzeugte polyploide Pflanzen. III. Meiosis in tetraploidem Hanf. (japanisch), Bot. and Zool. 8, 47-52.
- Nishiyama, I., 1941: Studien über künstlich erzeugte polyploide Pflanzen. V. Die Züchtung von langfasrigen Sorten durch Verdopplung der Chromosomenzahl bei Hanf. (japanisch), Bot. and Zool. 9, 173-179.
- Nishiyama, I., I. Yamada and M. Mezaki, 1947: Studien über künstlich erzeugte polyploide Pflanzen. XI. Wechsel im Geschlechtsverhältnis in einer Nachkommenschaft von tetraploiden Hanf. (japanisch), Seihen Ziho 3, 144-151.
- Nova, 2002: Nachwachsende Rohstoffe, Statistiken und Grafiken. Das Internet-Dienstleistungszentrum für die Naturfaserwirtschaft und nachwachsende Rohstoffe. www.nova-institut.de
- Olson, M., L. Hood, C. Cantor and D. Botstein, 1989: A common language for physical mapping of the human genome. Science 245, 1434-1435.
- Orita, M., H. Iwahana, H. Kanazawa, K. Hayashi and T. Sekiya, 1989: Detection of polymorphisms of human DNA by gel electrophoresis as single-strand conformation polymorphisms. Proc. Natl. Acad. Sci. USA 86, 2766-2770.

- Ovesná, J., K. Poláková and L. Leišová, 2002: DNA Analysis and their Application in Plant Breeding. Czech. J. Genet. Plant. Breed. 38, 29-40.
- Paran, I. and R. W. Michelmore, 1993: Development of reliable PCR-based markers linked to downy mildew resistance genes in lettuce. Theor. Appl. Genet. 85, 985-993.
- Patzak, J., P. Vejl, S. Skupinová and V. Nesvadba, 2002: Identifikation of sex in F₁ progenies of Hop (*Humulus lupulus* L.) by molecular marker. Rostilinná Výroba 48, 318-321.
- Peil, A., V. Korzun, V. Schubert, E. Schumann, W. E. Weber und M. Röder, 1998: The application of wheat microsatellites to identify disomic *Triticum aestivum*- *Aegilops markgrafii* addition lines. Theor. Appl. Genet. 96, 138-146.
- Peil, A., H. Flachowsky, E. Schumann and W. E. Weber, 2003: Sex-linked AFLP markers indicate a pseudo autosomal region in hemp (*Cannabis sativa* L.). Theor. Appl. Genet. in press.
- Petri, G., P. Veszki, P. Oroszlán and G. Turiák, 1996: Cannabinoid content of foreign and domestic fibre hemp varieties cultivated in hungary. Acta Agronomica Hungarica 42, 133-140.
- Polley, A., E. Seigner and M. W. Ganal, 1997: Identification of sex in hop (*Humulus lupulus*) using molecular markers. Genome 40, 357-361.
- Porceddu, A., E. Albertini, G. Barcaccia, E. Falistocco and M. Falcinelli, 2002: Linkage mapping in apomictic and sexual Kentucky bluegrass (*Poa pratensis* L.) genotypes using a two way pseudo-testcross strategy based on AFLP and SAMPL markers. Theor. Appl. Genet. 104, 273-280.
- Postma, W. P., 1946: Mitosis, meiosis en allopolloidie bij *Cannabis sativa* en *Spinacia oleracea*. H. D. Tjenk, Willink & Co., Amsterdam.
- Price, A. H., K. A. Steele, B. J. Moore, P. B. Barraclough and L. J. Clark, 2000: A combined RFLP and AFLP linkage map of upland rice (*Oryza sativa* L.) used to identify QTLs for root-penetration ability. Theor. Appl. Genet. 100, 49-56.
- Qu, L.-J., T. N. Foote, M. A. Roberts, T. A. Money, L. Aragón-Alcaide, J. W. Snape and G. Moore, 1998: A simple PCR-based method for scoring the ph1b deletion in wheat. Theor. Appl. Genet. 96, 371-375.
- Rae, S. J., C. Aldam, I. Dominguez, M. Hoebrechts, S. R. Barnes and K. J. Edwards, 2000: Development and incorporation of microsatellite markers into the linkage map of sugar beet (*Beta vulgaris* spp.). Theor. Appl. Genet. 100, 1240-1248.
- Rath, L., 1967: Heterosisversuche mit Hanf. Techn. Textilien 10, 247-260.
- Reamon-Büttner, S. M. and C. Jung, 2000: AFLP-derived STS markers for the identification of sex in *Asparagus officinalis* L.. Theor. Appl. Genet. 100, 432-438.

- Reamon-Büttner, S. M., T. Schmidt and C. Jung, 1999: AFLP's represent highly repetitive sequences in *Asparagus officinalis* L.. Chromosome Research 7, 297-304.
- Reamon-Büttner, S. M., J. Schondelmaier and C. Jung, 1998: AFLP markers tightly linked to the sex locus in *Asparagus officinalis* L.. Mol. Breed. 4, 91-98.
- Reddy, K. D., E. G. Abraham and J. Nagaraju, 1999: Microsatellites of the silkworm, *Bombyx mori*: abundance, polymorphism and strain characterization. Genome 42, 1057-1065.
- Reddy, O. U. K., E. A. Siddiq, N. P. Sarma, J. Ali, A. J. Hussain, P. Nimmakayala, P. Ramasamy, S. Pammi and A. S. Reddy, 2000: Genetic analysis of temperature-sensitive male sterility in rice. Theor. Appl. Genet. 100 (5), 794-801.
- Réjon, C. R., M. Jamilena, M. G. Ramos, J. S. Parker und M. R. Réjon, 1994: Cytogenetic and molecular analysis of the multiple sex-chromosome system of *Rumex acetosa*. Heredity 72, 209-215.
- Ren, N. and M. P. Timko, 2001: AFLP analysis of genetic polymorphism and evolutionary relationships among cultivated and wild *Nicotiana* species. Genome 44, 559-571.
- Reuter, K., 1987: Anbau und Verwendung von Hanf (*Cannabis sativa*) als Lieferant von Fasern, Samen und Rauschdrogen. Diplomarbeit, Rheinische-Friedrich-Wilhelm-Universität, Bonn.
- Riedel, M., 2000: Zytologische und molekularbiologische Untersuchungen beim Hanf. Diplomarbeit, Martin-Luther-Universität Halle-Wittenberg.
- Rivera, R., K. J. Edwards, J. H. A. Barker, G. M. Arnold, G. Ayad, T. Hodgkin and A. Karp, 1999: Isolation and characterization of polymorphic microsatellites in *Cocos nucifera* L.. Genome 42, 668-675.
- Rizet, G., 1946: Sexualité et polyploidie chez *Cannabis sativa*. C. r. Soc. Biol. Paris 140, 284-285.
- Ross, R., X. L. Ross, B. Rueger, T. Laengin and A. B. Reske-Kunz, 1999: Nonradioactive Detection of Differentially Expressed Genes Using Complex RNA or DNA Hybridization Probes. BioTechniques 26, 150-155.
- Ruiz Rejón, C., M. Jamilena, M. Garrido Ramos, J. Parker and M. Ruiz Rejón, 1994: A cytogenetic and molecular analysis of the multiple sex chromosome system of *Rumex acetosa*. Heredity. 79, 209-215.
- Saal, B. and G. Wricke, 1999: Development of simple sequece repeat markers in rye (*Secale cereale* L.). Genome 42, 964-972.

- Saghai Maroof, M. A., K. M. Soliman, R. A. Jorgensen and R. W. Allard, 1984: Ribosomal DNA spacer-length polymorphisms in barley: Mendelian inheritance, chromosomal location, and population dynamics. *Proc. Natl. Acad. Sci. USA* 81, 8014-8018.
- Sakamoto, K., Y. Akiyama, K. Fukui, H. Kamada and A. S. Satoh, 1998: Characterization, Genom Size and Morphologie of Sex Chromosomes in Hemp (*Cannabis sativa* L.). *Cytologia* 63, 459-464.
- Sakamoto, K., N. Ohmido, K. Fukui, H. Kamada and S. Satoh, 2000: Site-specific accumulation of a LINE-like retrotransposon in a sex chromosom of the dioecious plant *Cannabis sativa*. *Plant Mol. Biol.* 44, 723-732.
- Sakamoto, K., K. Shimomura, Y. Komeda, H. Kamada and A. S. Satoh, 1995: A male-associated DNA sequence in a dioecious plant, *Cannabis sativa* L.. *Plant Cell Physiol.* 36, 1549-1554.
- Saliba-Colombani, V., M. Causse, L. Gervais and J. Philouze, 2000: Efficiency of RFLP, RAPD, and AFLP markers for the construction of an intraspecific map of the tomato genome. *Genome* 43, 29-40.
- Savelkoul, P. M., H. M. Aarts, J. de Haas, L. Dijkshoorn, B. Duim, M. Otsen, J. W. Rademaker, L. Schouls and J. A. Lenstra, 1999: Amplified-fragment length polymorphism analysis: the state of an art. *J. Clinical Microbiol.* 37, 3083-3091.
- Schneider, K., 1942: Die Anatomie als Ursache des ungleichmäßigen chemischen Aufschlusses bei Hanf, Beeinflussung durch Ernährung, zugleich ein Beitrag zur Frage des Gummihanfes. *Faserforschung* 4, 141-165.
- Schneider, K., D. Borchardt, R. Schäfer-Pregl, N. Nagl, C. Glass, A. Jeppson, C. Gebhardt and F. Salamini, 1999: PCR-based cloning and segregation analysis of functional gene homologues in *Beta vulgaris*. *Mol. Gen. Genet.* 262, 515-524.
- Schumacher, K., J. Schondelmaier, E. Barzen, G. Steinrücken, D. Borchardt, W. E. Weber, C. Jung and F. Salamini, 1997: Combining different linkage maps in sugar beet (*Beta vulgaris* L.) to make one map. *Plant Breeding* 116, 23-38.
- Schumann, E., A. Peil and W. E. Weber, 1999: Preliminary results of a German field trial with different hemp (*Cannabis sativa* L.) accessions. *Genet. Res. Crop Evol.* 46: 399-407.
- Schuster, D. M., M. Darfler, J. E. Lee and A. Rashtchian, 1998: Improved Sensitivity and Specificity of RT-PCR. *Focus* 20, 34-35.
- Schuster, W., 1986: Ölpflanzen in Europa. DLG-Verlag, Frankfurt am Main, 108-111.

- Schwarz, G., W. Michalek, V. Mohler, G. Wenzel and A. Jahoor, 1999: Chromosome landing at the *Mla* locus in barley (*Hordeum vulgare* L.) by means of high-resolution mapping with AFLP markers. *Theor. Appl. Genet.* 98, 521-530.
- Scott, L. J., M. Cross, M. Shepherd, T. Maguire and R. J. Henry, 1999: Increasing the Efficiency of Microsatellite Discovery from Poorly Enriched Libraries in Coniferous Forest Species. *Plant Mol. Biol. Rep.* 17, 351-354.
- Sebastian, R. L., E. C. Howell, G. J. King, D. F. Marshall and M. J. Kearsey, 2000: An integrated AFLP and RFLP *Brassica oleracea* linkage map from two morphologically distinct doubled-haploid mapping populations. *Theor. Appl. Genet.* 100, 75-81.
- Sefc, K. M., F. Regner, E. Turetschek, J. Glossl and H. Steinkellner, 1999: Identification of microsatellite sequences in *Vitis riparia* and their applicability for genotyping of different *Vitis* species. *Genome* 42, 1-7.
- Shan, X., T. K. Blake and L. E. Talbert, 1999: Conversion of AFLP to sequence-specific PCR markers. *Theor. Appl. Genet.* 98, 1072-1078.
- Sharked, H., K. Kashkush, H. Ozkan, M. Feldman and A. A. Levy, 2001: Sequence Elimination and Cytosine Methylation Are Rapid and Reproducible Responses of the Genome to Wide Hybridization and Allopolyploidy in Wheat. *Plant Cell* 13, 1749-1759.
- Sharma, A., R. Sharma and H. Machii, 2000: Assessment of genetic diversity in a *Morus* germplasm collection using fluorescence-based AFLP markers. *Theor. Appl. Genet.* 101, 1049-1055.
- Sharma, P. C., B. Hüttel, P. Winter, G. Kahl, R. C. Gardner and K. Weising, 1995a: The potential of microsatellites for hybridization- and PCR-based fingerprinting of chickpea (*Cicer arietinum* L.) and related species. *Electrophoresis* 16, 1755-1761.
- Sharma, P.C., P. Winter, T. Bünger, B. Hüttel, F. Weigand, K. Weising and G. Kahl, 1995b: Abundance and polymorphism of di-, tri- and tetranucleotide tandem repeats in chickpea (*Cicer arietinum* L.). *Theor. Appl. Genet.* 90, 90-96.
- Shirkot P., D. R. Sharma and T. Mohapatra, 2002: Molecular identification of sex in *Actinidia deliciosa* by RAPD markers. *Scientia Horticulturae* 94, 33-99.
- Shure, M., S. Wessler and N. V. Fedoroff, 1983: Molecular identification and isolation of the waxy locus in maize. *Cell* 35, 225-233.
- Sinoto, Y., 1929: Chromosome studies in some dioecious plants, with special reference to the allosomes. *Cytologia* 1, 109-191.
- Sloop, K. W. and S. J. Rhodes, 2000: Combined Hot start and Touchdown PCR Protocols Optimize Amplification of Difficult Targets. *Biochemica* 1, 35-36.

- Small, E., 1979: The species problem in *Cannabis* (science and semantics). Cropus-Verlag, Toronto.
- Southern, E. M., 1975: Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J. Mol. Biol.* 98, 503-517.
- Stam, P., 1993: Construction of integrated genetic linkage maps by means of a new computer package JOINMAP. *Plant J.* 3, 739-744.
- Staub, J. E., F. C. Serquen and M. Gupta, 1996: Genetic markers, map construction, and their application in plant breeding. *Hortscience* 31, 729-741.
- Stavolone, L., D. Alioto, A. Ragozzino and J.-F. Laliberté, 1998: Variability Among Turnip Mosaic Potyvirus Isolates. *Phytopathology* 88, 1200-1204.
- Stirling, B., G. Newcombe, J. Vrebalov, I. Bosdet and H. D. Bradshaw Jr., 2001: Suppressed recombination around the *MXC3* locus, a major gene for resistance to poplar leaf rust. *Theor. Appl. Genet.* 103, 1129-1137.
- Stitou, S., R. Jiménez, D. de la Guardia and M. Burgos, 2001: Silent ribosomal cistrons are located at the pairing segment of the postreductional sex chromosomes of *Apodemus sylvaticus* (Rodentia, Muridae). *Heredity* 86, 128-133.
- Sung, S.-K., G.-H. Yu, J. Nam, D.-H. Jeong and G. An, 1999: Developmentally regulated expression of two MADS-box genes, *MdMADS3* and *MdMADS4*, in the morphogenesis of flower buds and fruits in apple. *Planta* 210, 519-528.
- Suzuki, Y., T. Sekiya and K. Hayashi, 1991: Allele-specific PCR: A method for amplification and sequence determination of a single component among a mixture of sequence variants. *Anal. Biochem.* 192, 82-84.
- Takenaka, Y., 1951: Studies on sex-differentiation in higher plants. *Jahresbericht Nat. Inst. der Genetik (Japan 1949-1950)* 53, 32-33.
- Taramino, G. and S. V. Tingey, 1996: Simple sequence repeats for germplasm analysis and mapping in maize. *Genome* 39, 277-287.
- Taura, F., S. Morimoto and Y. Shoyama, 1996: Purification and Characterisation of Cannabidiolic-acid Synthase from *Cannabis sativa* L.. *J. Biol. Chemistry.* 271, 17411-17416.
- Taura, F., H. Matsushita, S. Morimoto and Y. Masayama, 2000: Tetrahydrocannabinolic acid synthase gene. Patent: JP 2000078979-A 13, Taishopharmaceut co Ltd.
- Tautz, D., 1989: Hypervariability of simple sequences as a general source for polymorphic DNA. *Nuc. Ac. Res.* 17, 6463
- Taylor, B. and A. Powell, 1985: Isolation of plant DNA and RNA. *Focus* 4, 4-6.

- Terauchi, R. and G. Kahl, 1999: Mapping of the *Dioscorea tokoro* genome: AFLP markers linked to sex. *Genome* 42, 752-762.
- Tischler, G., 1953: Allgemeine Pflanzenkaryologie. In: Linsbauer, K.: Handbuch der Pflanzenanatomie. Band. 2, Gebr. Bornträger, Berlin.
- Tobler, F., 1940: Beitrag zur Kenntnis der Ligninbildung im Holzkörper von *Cannabis sativa*. Vortragsband - Botanisches Institut der Technischen Hochschule Dresden, 143-148.
- Tozaki, T., S. Inoue, M. Mashima, M. Ohta, N. Miura and M. Tomita, 2000: Sequence analysis of trinucleotide repeat microsatellites from an enrichment library of the equine genome. *Genome* 43, 354-365.
- Urasaki, N., M. Tokumoto, K. Tarora, Y. Ban, T. Kayano, H. Tanaka, H. Oku, I. Chinen and R. Terauchi, 2002: A male and hermaphrodite specific RAPD marker for papaya (*Carica papaya* L.). *Theor. Appl. Genet.* 104, 281-285.
- Uzunova, M. I. and W. Ecke, 1999: Abundance, polymorphism and genetic mapping of microsatellites in oilseed rape (*Brassica napus* L.). *Plant Breeding* 118, 323-326.
- van de Wiel, C., P. Arens and B. Vosman, 1998: Microsatellite fingerprinting in lettuce (*Lactuca sativa* L.) and wild relatives. *Plant Cell Reports* 17, 837-842.
- van Heusden, A. W., J. W. van Ooijen, R. V. Ginkel, W. H. J. Verbeek, W. A. Wietsma and C. Kik, 2000: A genetic map of an interspecific cross in *Allium* based on amplified fragment length polymorphism (AFLPTM) markers. *Theor. Appl. Genet.* 100, 118-126.
- Vinutzer, B. A., A. Patocchi, L. Gianfranceschi, S. Tartarini, H. B. Zhang, C. Gessler and S. Sansavini, 2001: Apple (*Malus* sp.) contains receptor-like genes homologous to the *Cf* resistance gene family of tomato with a cluster of such genes co-segregating with *Vf* apple scab resistance. unpublished, <http://www.ncbi.nlm.nih.gov/>
- von Sengbusch, R., 1943: Die Geschlechtsvererbung bei Hanf und die Züchtung eines monözischen Hanfes. *Der Züchter* 15, 49-62.
- von Sengbusch, R., 1952: Ein weiterer Beitrag zur Vererbung des Geschlechts bei Hanf als Grundlage für die Züchtung eines monözischen Hanfes. *Z. für Pflanzenzüchtg* 31, 319-338.
- Vos, P., R. Hogers, M. Bleeker, M. Reijans, T. van der Lee, M. Hornes, A. Frijters, J. Pot, J. Peleman, M. Kuiper and M. Zabeau, 1995: AFLP: a new technique for DNA fingerprinting. *Nuc. Ac. Res.* 23, 4407-4414.
- Wagner, D. B., G. R. Furnier, M. A. Saghai-Marooof, S. M. Williams, B. P. Dancik and R. W. Allard, 1987: Chloroplast DNA polymorphisms in lodgepole and jack pines and their hybrids. *Proc. Natl. Acad. Sci. USA* 84, 2097-2100.

- Warmke, H. E. and A. F. Blakslee, 1939: Sex mechanism in polyploids of *Melandrium*. Science 89, 391-392.
- Weber, J. L. and P. E. May, 1989: Abundant class of human DNA polymorphism which can be typed using the polymerase chain reaction. Am. J. Hum. Genet. 44, 388-396.
- Weber, W. E. and G. Wricke, 1994: Genetic Markers in Plant Breeding. Adv. Plant Breeding 18, Paul Parey, Berlin, 1-105.
- Wei, F., K. Gobelman-Werner, S. M. Morroll, J. Kurth, L. Mao, R. Wing, D. Leister, P. Schulze-Lefert and R. P. Wise, 1999: The *Mla* (Powdery Mildew) Resistance Cluster Is Associated With Three NBS-LRR Gene Families and Suppressed Recombination Within a 240-kb DNA Interval on Chromosome 5S (1HS) of Barley. Genetics 153, 1929-1948.
- Weigel, D. and E. M. Meyerowitz, 1994: The ABCs of floral homeotic genes. Cell 78, 203-209.
- Weising, K., R. G. Atkinson and R. C. Gardener, 1995: Genomic Fingerprinting by Microsatellite-primed PCR: A Critical Evaluation. Research 4, 249-255.
- Weising, K., R. W. M. Fung, D. J. Keeling, R. G. Atkinson and R. C. Gardner, 1996: Characterization of microsatellites from *Actinidia chinensis*. Mol. Breeding 2, 117-131.
- Weising, K., D. Kaemmer, F. Weigand, J. T. Epplen and G. Kahl, 1992: Oligonucleotide fingerprinting reveals probe-dependent levels of informativeness in chickpea (*Cicer arietinum* L.). Genome 35, 436-442.
- Weising, K., D. Ramser, D. Kaemmer, G. Kahl and J. T. Epplen, 1991: Oligonucleotide fingerprinting in plants and fungi. In: T. Burke, G. Dolf, A. J. Jeffreys, and R. Wolff (Eds.), DNA Fingerprinting Approaches and Applications, Birkhauser Verlag, Basel, Switzerland, 112-123.
- Welsh, J., K. Chada, S. S. Dalal, D. Ralph, R. Cheng and M. McClelland, 1992: Arbitrarily primed PCR fingerprinting of RNA. Nuc. Ac. Res. 20, 4965-4970.
- Westergard, M., 1953: Über den Mechanismus der Geschlechtsbestimmung bei *Melandrium album*. Naturwissenschaften 9, 253-260.
- Williams, J. G. K., A. R. Kubelik, K. J. Livak and J. A. Rafalski, 1990: DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. Nuc. Ac. Res. 18, 6531-6535.
- Wong, A., M. R. Forbes and M. L. Smith, 2001: Characterization of AFLP markers in damselflies: prevalence of codominant markers and implications for population genetic applications. Genome 44, 677-684.

- Wu, R. L., Y. F. Han, J. J. Hu, L. Li, M. L. Li and Z.-B. Zeng, 2000: An integrated genetic map of *Populus* based on amplified fragment length polymorphisms. *Theor. Appl. Genet.* 100, 1249-1256.
- Wyman, A. R. and R. White, 1980: A highly polymorphic locus in human DNA. *Proc. Nat. Acad. Sci. USA* 77, 6754-6758.
- Xu, M., A. E. Melchinger, X. C. Xia and T. Luebberstedt, 1999: High-resolution mapping of loci conferring resistance to sugarcane mosaic virus in maize using RFLP, SSR and AFLP markers. *Mol. Gen. Genet.* 261, 574-581.
- Xu, M., X. Li and S. S. Korban, 2000: AFLP-Based Detection of DNA Methylation. *Plant Mol. Biol. Rep.* 18, 361-368.
- Xu, M., E. Huaracha and S. S. Korban, 2001: Development of sequence-characterized amplified regions (SCARs) from amplified fragment length polymorphism (AFLP) markers tightly linked to the *Vf* gene in apple. *Genome* 44, 63-70.
- Yamada, I., 1943: Die Geschlechtschromosomen des Hanfes. *Seihen Ziho* 2, 64-68.
- Yao, L.-J., Y. H. Dong, A. Kvarnheden and B. Morris, 1999: Seven MADS-box Genes in Apple are Expressed in Different Parts of the Fruit. *J. Amer. Soc. Hort. Sci.* 124, 8-13.
- Yap, E. P. and JO'D. McGee, 1992: Non-isotopic SSCP detection in PCR products by ethidium bromide staining. *Trends Genet.* 8, 49.
- Yu, G.-X. and R. P. Wise, 2000: An anchored AFLP-and retrotransposon-based map of diploid *Avena*. *Genome* 43, 736-749.
- Zhang H., V. S. Stilio, F. Rehman, A. Avery, D. Mulcahy and R. Kesseli, 1998: Y chromosome specific markers and the evolution of dioecy in the genus *Silene* Y. *Genome* 41, 141-147.