

6 Literaturverzeichnis

- Adam, K.P., Thiel, R., Zapp, J. (1999). Incorporation of 1-[1-¹³C]deoxy-D-xylulose in chamomile sesquiterpenes. *Arch. Biochem. Biophys.* **369**, 127-132.
- Adam, P., Eisenreich, W., Fellermeier, M., Hecht, S., Rohdich, F., Schuhr, C.A., Wungsintaweekul, J., Zenk, M.H., Bacher, A. (2000). International Patent Application PCT/EP01/06255.
- Agranoff, B.W., Eggerer, H., Henning, U., Lynen, F. (1960). Biosynthesis of terpenes. VII. Isopentenyl pyrophosphate isomerase. *J. Biol. Chem.* **235**, 326-332.
- Ahlemeyer, B., Krieglstein, J. (2003). Pharmacological studies supporting the therapeutic use of *Ginkgo biloba* extract for Alzheimer's disease. *Pharmacopsychiatry* **36**, S8-S14.
- Altincicek, B., Kollas, A.K., Sanderbrand, S., Wiesner, J., Hintz, M., Beck, E., Jomaa, H. (2001a). GcpE is involved in the 2-C-methyl-D-erythritol 4-phosphate pathway of isoprenoid biosynthesis in *Escherichia coli*. *J. Bacteriol.* **183**, 2411-2416.
- Altincicek, B., Kollas, A., Eberl, M., Wiesner, J., Sanderbrand, S., Hintz, M., Beck, E., Jomaa, H. (2001b). *LytB*, a novel gene of the 2-C-methyl-D-erythritol 4-phosphate pathway of isoprenoid biosynthesis in *Escherichia coli*. *FEBS Lett.* **499**, 37-40.
- Altincicek, B., Duin, E.C., Reichenberg, A., Hedderich, R., Kollas, A.K., Hintz, M., Wagner, S., Wiesner, J., Beck, E., Jomaa, H. (2002). *LytB* protein catalyzes the terminal step of the 2-C-methyl-D-erythritol-4-phosphate pathway of isoprenoid biosynthesis. *FEBS Lett.* **532**, 437-440.
- Anderson, M.S., Muehlbacher, M., Street, I.P., Proffitt, J., Poulter, C.D. (1989). Isopentenyl diphosphate:dimethylallyl diphosphate isomerase. An improved purification of the enzyme and isolation of the gene from *Saccharomyces cerevisiae*. *J. Biol. Chem.* **264**, 19169-19175.
- Araki, N., Kusumi, K., Masamoto, K., Niwa, Y., Iba, K. (2000). Temperature-sensitive *Arabidopsis* mutant defective in 1-deoxy-D-xylulose 5-phosphate synthase within the plastid non-mevalonate pathway for isoprenoid biosynthesis. *Physiol. Plant.* **108**, 19-24.
- Araujo, A.P.U., Oliva, G., Henrique-Silva, F., Garratt, R.C., Cáceres, O., Beltramini, L.M. (2000). Influence of the histidine tail on the structure and activity of recombinant chlorocatechol 1,2-dioxygenase. *Biochem. Biophys. Research Comm.* **272**, 480-484.
- Arfin, S.M., Umbarger, H.E. (1969). Purification and properties of the acetohydroxy acid isomeroreductase of *Salmonella typhimurium*. *J. Biochem.* **244**, 1118-1127.
- Arigoni, D., Sagner, S., Latzel, C., Eisenreich, W., Bacher, A., Zenk, M.H. (1997). Terpenoid biosynthesis from 1-deoxy-D-xylulose in higher plants by intramolecular skeletal rearrangement. *Proc. Natl. Acad. Sci. USA* **94**, 10600-10605.

- Arigoni, D., Eisenreich, W., Latzel, C., Sagner, S., Radykewicz, T., Zenk, M.H., Bacher, A. (1999). Dimethylallyl pyrophosphate is not the committed precursor of isopentenyl pyrophosphate during terpenoid biosynthesis from 1-deoxyxylulose in higher plants. *Proc. Natl. Acad. Sci. USA* **96**, 1309-1314.
- Ausubel, F.M., Brent, R., Kingston, R.E., Moore, D.D., Seidman, J.G., Smith, J.A., Struhl, K. (1999). *Short Protocols in Molecular Biology*, 4th edition. A compendium of methods from current protocols in molecular biology. Wiley New York.
- Banthorpe, D.V., Charlwood, B.V., Francis, M.J.O. (1972). The biosynthesis of monoterpenes. *Chem. Rev.* **72**, 115-155.
- Banthorpe, D.V., Doonan, S., Gutowski, J.A. (1977). Isopentenyl pyrophosphate isomerase from pig liver. *Arch. Biochem. Biophys.* **184**, 381-390.
- Baulcombe, D.C. (1999). Fast forward genetics based on virus-induced gene silencing. *Curr. Opin. Plant Biol.* **2**, 109-113.
- Baulcombe, D. (2004). RNA silencing in plants. *Nature* **431**, 356-363.
- Bick, J.A., Lange, B.M. (2003). Metabolic cross talk between cytosolic and plastidial pathways of isoprenoid biosynthesis: unidirectional transport of intermediates across the chloroplast envelope membrane. *Arch. Biochem. Biophys.* **415**, 146-154.
- Blagg, B.S., Poulter, C.D. (1999). Synthesis of 1-deoxy-D-xylulose and 1-deoxy-D-xylulose-5-phosphate. *J. Org. Chem.* **64**, 1508-1511.
- Bochar, D.A., Friesen, J.A., Stauffacher, C.V., Rodwell, V.W. (1999). Biosynthesis of mevalonic acid from acetyl-CoA. *Comprehensive Natural Products Chemistry. Isoprenoids Including Carotenoids and Steroids*. D.E. Cane (Ed.) **2**, 15-44, Pergamon, Oxford, UK.
- Bouvier, F., d'Harlingue, A., Suire, C., Backhaus, R.A., Camara, B. (1998). Dedicated roles of plastid transketolases during the early onset of isoprenoid biogenesis in pepper fruits. *Plant Physiol.* **117**, 1423-1431.
- Bradford, M.M. (1976). A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**, 248-254.
- Broers, S.T.J. (1994). Über die frühen Stufen der Biosynthese von Isoprenoiden in *Escherichia coli*. Dissertation Nr. 10978, ETH Zürich, Schweiz.
- Burch-Smith, T.M., Anderson, J.C., Martin, G.B., Dinesh-Kumar, S.P. (2004). Applications and advantages of virus-induced gene silencing for gene function studies in plants. *The Plant Journal.* **39**, 734-746.
- Campbell, M., Hahn, F.M., Poulter, C.D., Leustek, T. (1997). Analysis of the isopentenyl diphosphate isomerase gene family from *Arabidopsis thaliana*. *Plant Mol. Biol.* **36**, 323-328.

Campos, N., Rodriguez-Concepcion, M., Seemann, M., Rohmer, M., Boronat, A. (2001). Identification of *gcpE* as a novel gene of the 2-C-methyl-D-erythritol 4-phosphate pathway for isoprenoid biosynthesis in *Escherichia coli*. *FEBS Lett.* **488**, 170-173.

Carrigan, C.N., Poulter, C.D. (2003). Zinc is an essential cofactor for type I isopentenyl diphosphate:dimethylallyl diphosphate isomerase. *J. Am. Chem. Soc.* **125**, 9008-9009.

Cassera, M.B., Gozzo, F.B., D'Alexandri, F.L., Merino, E.F., del Portillo, H.A., Peres, V.J., Almeida, I.C., Eberlin, M.N., Wunderlich, G., Wiesner, J., Jomaa, H., Kimura, E.A., Katzin, A.M. (2004). The methylerythritol phosphate pathway is functionally active in all intraerythrocytic stages of *Plasmodium falciparum*. *J. Biol. Chem.* **279**, 51749-51750.

Charon, L., Pale-Grosdemange, C., Rohmer, M. (1999). On the reduction steps in the mevalonate independent 2C-methyl-D-erythritol 4-phosphate (MEP) pathway for isoprenoid biosynthesis in the bacterium *Zymomonas mobilis*. *Tetrahedron Lett.* **40**, 7231-7234.

Cunningham, F.X., Jr., Gantt, E. (2000). Identification of multi-gene families encoding isopentenyl diphosphate isomerase in plants by heterologous complementation in *Escherichia coli*. *Plant Cell Physiol.* **41**, 119-123.

Cunningham, F.X., Jr., Lafond, T.P., Gantt, E. (2000). Evidence of a role for LytB in the nonmevalonate pathway of isoprenoid biosynthesis. *J. Bacteriol.* **182**, 5841-5848.

Davisson, V.J., Woodside, A.B., Poulter C.D. (1985). Synthesis of allylic and homoallylic isoprenoid pyrophosphates. *Methods Enzymol.* **110**, 130-144.

Davisson, V.J., Woodside, A.B., Neal, T.R., Stremler, K.E., Muehlbacher, M., Poulter C.D. (1986). Phosphorylation of isoprenoid alcohols. *J. Org. Chem.* **51**, 4768-4779.

Disch, A., Rohmer, M. (1998). On the absence of the glyceraldehyd 3-phosphate/pyruvate pathway for isoprenoid biosynthesis in fungi and yeasts. *FEMS Microbiol. Lett.* **168**, 201-208.

Disch, A., Hemmerlin, A., Bach, T.J., Rohmer, M. (1998). Mevalonate-derived isopentenyl diphosphate is the biosynthetic precursor of ubiquinone prenyl side chain in tobacco BY-2 cells. *Biochem. J.* **331**, 615-621.

Dogbo, O., Camara, B. (1987). Purification of isopentenyl pyrophosphate isomerase and geranylgeranyl pyrophosphate synthase from *Capsicum* chromoplasts by affinity chromatography. *Biochim. Biophys. Acta* **920**, 140-148.

Durbecq, V., Sainz, G., Oudjama, Y., Clantin, B., Bompard-Gilles, C., Tricot, C., Caillet, J., Stalon, V., Droogmans, L., Villeret, V. (2001). Crystal structure of isopentenyl diphosphate:dimethylallyl diphosphate isomerase. *EMBO J.* **20**, 1530-1537.

Duvold, T., Bravo, J.-M., Pale-Grosdemange, C., Rohmer, M. (1997a). Biosynthesis of 2C-methyl-D-erythritol, a putative C₅ intermediate in the mevalonate independent pathway for isoprenoid biosynthesis. *Tetrahedron Lett.* **38**, 4769-4772.

Duvold, T., Cali, P., Bravo, J.-M., Rohmer, M. (1997b). Incorporation of 2C-methyl-D-erythritol, a putative isoprenoid precursor in the mevalonate-independent pathway, into ubiquinone and menaquinone of *Escherichia coli*. *Tetrahedron Lett.* **38**, 6181-6184.

- Eberl, M., Hintz, M., Reichenberg, A., Kollas, A.K., Wiesner, J., Jomaa, H. (2003). Microbial isoprenoid biosynthesis and human $\gamma\delta$ T cell activation. *FEBS Lett.* **544**, 4-10.
- Eisenreich, W., Schwarz, M., Cartayrade, A., Arigoni, D., Zenk, M.H., Bacher, A. (1998). The deoxyxylulose phosphate pathway of terpenoid biosynthesis in plants and microorganisms. *Chemistry & Biology* **5**, R221-R233.
- Eisenreich, W., Rohdich, F., Bacher, A. (2001). Deoxyxylulose phosphate pathway to terpenoids. *Trends Plant Sci.* **6**, 78-84.
- Eisenreich, W., Bacher, A., Arigoni, D., Rohdich, F. (2004). Biosynthesis of isoprenoids via the non-mevalonate pathway. *Cell Mol. Life Sci.* **61**, 1401-1426.
- Enfissi, E.M.A., Fraser, P.D., Lois, L.M., Boronat, A., Schuch, W., Bramley, P.M. (2005). Metabolic engineering of the mevalonate and nonmevalonate isopentenyl diphosphate-forming pathways for the production of health-promoting isoprenoids in tomato. *Plant Biotechnol. J.* **3**, 17-27.
- Englard, S., Seifter, S. (1990). Precipitation techniques. *Methods Enzymol.* **182**, 285-300.
- Estevez, J.M., Cantero, A., Reindl, A., Reichler, S., Leon, P. (2001). 1-Deoxy-D-xylulose-5-phosphate synthase, a limiting enzyme for plastidic isoprenoid biosynthesis in plants. *J. Biol. Chem.* **276**, 22901-22909.
- Fellermeier, M. (2000). Untersuchungen zum Alternativen Terpenbiosyntheseweg in Pflanzen. Dissertation, Ludwig-Maximilians-Universität München.
- Fellermeier, M., Raschke, M., Sagner, S., Wungsintaweekul, J., Schuhr, C.A., Hecht, S., Kis, K., Radykewicz, T., Adam, P., Rohdich, F., Eisenreich, W., Bacher, A., Arigoni, D., Zenk, M.H. (2001). Studies on the nonmevalonate pathway of terpene biosynthesis. The role of 2C-methyl-D-erythritol 2,4-cyclodiphosphate in plants. *Eur. J. Biochem.* **268**, 6302-6310.
- Flesch, G., Rohmer, M. (1988). Prokaryotic hopanoids: the biosynthesis of the bacteriohopane skeleton. *Eur. J. Biochem.* **175**, 405-411.
- Flügge, U.I., Gao, W. (2005). Transport of isoprenoid intermediates across chloroplast envelope membranes. *Plant Biol.* **7**, 91-97.
- Fraser, P.D., Pinto, M.E., Holloway, D.E., Bramley, P.M. (2000). Technical advance: application of high-performance liquid chromatography with photodiode array detection to the metabolic profiling of plant isoprenoids. *Plant J.* **24**, 551-558.
- Gao, W., Loeser, R., Raschke, M., Dessoy, M.A., Fulhorst, M., Alpermann, H., Wessjohann, L.A., Zenk, M.H. (2002). (E)-4-Hydroxy-3-methylbut-2-enyl diphosphate: an intermediate in the formation of terpenoids in plant chromoplasts. *Angew. Chem. Int. Ed.* **41**, 2604-2607.
- Gao, W., Raschke, M., Alpermann, H., Zenk, M.H. (2003). A facile enzymatic synthesis of isotopically labeled 2-methyl-D-erythritol 2,4-cyclodiphosphate by spinach chloroplast stroma. *Helv. Chem. Acta* **86**, 3568-3577.

- Gerharz, T., Zelic, B., Takors, R., Bott, M. (2001). Method for the fermentative production of pyruvate. German patent application DE10129711.4.
- Gerharz, T. (2003). Pyruvate production based on acetate auxotrophic *Escherichia coli* strains. *Berichte des Forschungszentrums Jülich* i-xi, 1-133.
- Giner, J.L., Ferris, W.V., Jr. (2002). Synthesis of 2-C-methyl-D-erythritol 2,4-cyclopyrophosphate. *Org. Lett.* **4**, 1225-1226.
- Guevara-Garcia, A., San Roman C., Arroyo, A., Cortes, M.E., Gutierrez-Nava, M. de la Luz, Leon, P. (2005). Characterization of the Arabidopsis *clb6* mutant illustrates the importance of posttranscriptional regulation of the methyl-D-erythritol 4-phosphate pathway. *Plant Cell.* **17**, 628-643.
- Gutierrez-Nava, M. de la Luz, Gillmor, C.S., Jimenez, L.F., Guevara-Garcia, A. Leon, P. (2004). Chloroplast biogenesis genes act cell and noncell autonomously in early chloroplast development. *Plant Physiology* **135**, 471-482.
- Guzman, M. (2003). Cannabinoids: potential anticancer agents. *Nat. Rev. Cancer.* **3**, 745-755.
- Hahn, F.M., Baker, J.A., Poulter, C.D. (1996a). Open reading frame 176 in the photosynthesis gene cluster of *Rhodobacter capsulatus* encodes *idi*, a gene for isopentenyl diphosphate isomerase. *J. Bacteriol.* **178**, 619-624.
- Hahn, F.M., Xuan, J.W., Chambers, A.F., Poulter, C.D. (1996b). Human isopentenyl diphosphate:dimethylallyl diphosphate isomerase: overproduction, purification, and characterization. *Arch. Biochem. Biophys.* **332**, 30-34.
- Hahn, F.M., Hurlburt, A.P., Poulter, C.D. (1999). *Escherichia coli* open reading frame 696 is *idi*, a nonessential gene encoding isopentenyl diphosphate isomerase. *J. Bacteriol.* **181**, 4499-4504.
- Hanahan, D. (1983). Studies on the transformation of *Escherichia coli* with plasmids. *J. Mol. Biol.* **166**, 557-580.
- Hanahan, D., Jessee, J., Bloom, F.R. (1991). Plasmid transformation of *Escherichia coli* and other bacteria. *Methods Enzymol.* **204**, 63-113.
- Hecht, S., Kis, K., Eisenreich, W., Amslinger, S., Wungsintaweekul, J., Herz, S., Rohdich, F., Bacher, A. (2001a). Enzyme-assisted preparation of isotope-labeled 1-deoxy-d-xylulose 5-phosphate. *J. Org. Chem.* **66**, 3948-3952.
- Hecht, S., Wungsintaweekul, J., Rohdich, F., Kis, K., Radykewicz, T., Schuhr, C.A., Eisenreich, W., Richter, G., Bacher, A. (2001b). Biosynthesis of terpenoids: efficient multistep biotransformation procedures affording isotope-labeled 2C-methyl-D-erythritol 4-phosphate using recombinant 2C-methyl-D-erythritol 4-phosphate synthase. *J. Org. Chem.* **66**, 7770-7775.
- Hecht, S., Eisenreich, W., Adam, P., Amslinger, S., Kis, K., Bacher, A., Arigoni, D., Rohdich, F. (2001c). Studies on the nonmevalonate pathway to terpenes: the role of the GcpE (IspG) protein. *Proc. Natl. Acad. Sci. USA* **98**, 14837-14842.

- Hellens, R.P., Edwards, E.A., Leyland, N.R., Bean, S., Mullineaux, P.M. (2000). pGreen: a versatile and flexible binary Ti vector for *Agrobacterium*-mediated plant transformation. *Plant Mol. Biol.* **42**, 819-832.
- Hemmerlin, A., Hoeffler, J.F., Meyer, O., Tritsch, D., Kagan, I.A., Grosdemange-Billiard, C., Rohmer, M., Bach, T.J. (2003). Cross-talk between the cytosolic mevalonate and the plastidial methylerythritol phosphate pathways in tobacco bright yellow-2 cells. *J. Biol. Chem.* **278**, 26666-26676.
- Herz, S., Wungsintaweekul, J., Schuhr, C.A., Hecht, S., Lüttgen, H., Sagner, S., Fellermeier, M., Eisenreich, W., Zenk, M.H., Bacher, A., Rohdich, F. (2000). Biosynthesis of terpenoids: YgbB protein converts 4-diphosphocytidyl-2C-methyl-D-erythritol 2-phosphate to 2C-methyl-D-erythritol 2,4-cyclodiphosphate. *Proc. Natl. Acad. Sci. USA* **97**, 2486-2490.
- Hill, R.E., Himmeldirk, K., Kennedy, I.A., Panolski, R.M., Sayer, B.G., Wolf, E., Spenser, I.D. (1996). The biogenetic anatomy of vitamin B₆. A ¹³C NMR investigation of the biosynthesis of pyridoxol in *Escherichia coli*. *J. Biol. Chem.* **271**, 30426-30435.
- Hintz, M., Reichenberg, A., Altincicek, B., Bahr, U., Gschwind, R.M., Kollas, A.K., Beck, E., Wiesner, J., Eberl, M., Jomaa, H. (2001). Identification of (E)-4-hydroxy-3-methyl-but-2-enyl pyrophosphate as a major activator for human $\gamma\delta$ T cells in *Escherichia coli*. *FEBS Lett.* **509**, 317-322.
- Hohmann, C. (2004). Antibiotika-Resistenzen. Die Situation ist nicht dramatisch. *Pharm. Ztg.* **149**, 732-733.
- Jez, J.M., Ferrer, J.L., Bowman, M.E., Dixon, R.A., Noel, J.P. (2000). Dissection of malonyl-coenzyme A decarboxylation from polyketide formation in the reaction mechanism of a plant polyketide synthase. *Biochemistry* **39**, 890-902.
- Jomaa, H., Wiesner, J., Sanderbrand, S., Altincicek, B., Weidemeyer, C., Hintz, M., Türbachova, I., Eberl, M., Zeidler, J., Lichtenthaler, H.K., Soldati, D., Beck, E. (1999). Inhibitors of the nonmevalonate pathway of isoprenoid biosynthesis as antimalarial drugs. *Science* **285**, 1573-1576.
- Jones, B.L., Porter, J.W. (1985). Enzymatic synthesis of phytoene. *Methods Enzymol.* **110**, 209-220.
- Joyard, J., Dorne, A.J., Douce, R. (1987). Use of thermolysin to probe the cytosolic surface of the outer envelope membranes from plastids. *Methods Enzymol.* **148**, 195-206.
- Kaneda, K., Kuzuyama, T., Takagi, M., Hayakawa, Y., Seto, H. (2001). An unusual isopentenyl diphosphate isomerase found in the mevalonate pathway gene cluster from *Streptomyces* sp. strain CL190. *Proc. Natl. Acad. Sci. USA* **98**, 932-937.
- Kennedy, I.A., Hemscheidt, T., Britten, J., Spenser, I.D. (1995). 1-Deoxy-D-xylulose. *Can. J. Chem.* **73**, 1329-1337.
- Kis, K., Wungsintaweekul, J., Eisenreich, W., Zenk, M.H., Bacher, A. (2000). An efficient preparation of 2C-methyl-D-erythritol 4-phosphoric acid and its derivatives. *J. Org. Chem.* **65**, 587-592.

Köhler, S., Delwiche, C.F., Denny, P.W., Tilney, L.G., Webster, P., Wilson, R.J., Palmer, J.D., Roos, D.S. (1997). A plastid of probable green algal origin in apicomplexan parasites. *Science* **275**, 1485-1489.

Kollas, A.K., Duin, E.C., Eberl, M., Altincicek, B., Hintz, M., Reichenberg, A., Henschker, D., Henne, A., Steinbrecher, I., Ostrovsky, D.N., Hedderich, R., Beck, E., Jomaa, H., Wiesner, J. (2002). Functional characterization of GcpE, an essential enzyme of the non-mevalonate pathway of isoprenoid biosynthesis. *FEBS Lett.* **532**, 432-436.

Koppisch, A.T., Blagg, B.S., Poulter, C.D. (1999). Synthesis of 2-C-methyl-D-erythritol 4-phosphate: the first pathway-specific intermediate in the methylerythritol phosphate route to isoprenoids. *Org. Lett.* **2**, 215-217.

Kuzuyama, T., Takahashi, S., Watanabe, H., Seto, H. (1998a). Direct formation of 2C-methyl-D-erythritol 4-phosphate from 1-deoxy-D-xylulose 5-phosphate by 1-deoxy-D-xylulose 5-phosphate reductoisomerase, a new enzyme in the non-mevalonate pathway to isopentenyl diphosphate. *Tetrahedron Lett.* **39**, 4509-4512.

Kuzuyama, T., Shimizu, T., Takahashi, S., Seto, H. (1998b). Fosmidomycin, a specific inhibitor of 1-deoxy-D-xylulose 5-phosphate reductoisomerase in the nonmevalonate pathway for terpenoid biosynthesis. *Tetrahedron Lett.* **39**, 7913-7916.

Kuzuyama, T., Takagi, M., Takahashi, S., Seto, H. (2000a). Cloning and characterization of 1-deoxy-D-xylulose 5-phosphate synthase from *Streptomyces* sp. Strain CL190, which uses both the mevalonate and nonmevalonate pathways for isopentenyl diphosphate biosynthesis. *J. Bacteriol.* **182**, 891-897.

Kuzuyama, T., Takagi, M., Kaneda, K., Dairi, T., Seto, H. (2000b). Formation of 4-(cytidine 5'-diphospho)-2C-methyl-D-erythritol from 2C-methyl-D-erythritol 4-phosphate by 2C-methyl-D-erythritol 4-phosphate cytidyltransferase, a new enzyme in the nonmevalonate pathway. *Tetrahedron Lett.* **41**, 703-706.

Kuzuyama, T., Takagi, M., Kaneda, K., Watanabe, H., Dairi, T., Seto, H. (2000c). Studies on the nonmevalonate pathway: conversion of 4-(cytidine 5'-diphospho)-2C-methyl-D-erythritol to its 2-phospho derivative by 4-(cytidine 5'-diphospho)-2C-methyl-D-erythritol kinase. *Tetrahedron Lett.* **41**, 2925-2928.

Kuzuyama, T. (2003). Mevalonate and nonmevalonate pathways for the biosynthesis of isoprene units. *Biosci. Biotechnol. Biochem.* **66**, 1619-1627.

Laemmli, U.K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680-685.

Lange, B.M., Wildung, M.R., McCaskill, D., Croteau, R. (1998). A family of transketolases that directs isoprenoid biosynthesis via a mevalonate-independent pathway. *Proc. Natl. Acad. Sci. USA* **95**, 2100-2104.

Lange, B.M., Croteau, R. (1999a). Isoprenoid biosynthesis via a mevalonate-independent pathway in plants: cloning and heterologous expression of 1-deoxy-D-xylulose-5-phosphate reductoisomerase from peppermint. *Arch. Biochem. Biophys.* **365**, 170-174.

- Lange, B.M., Croteau, R. (1999b). Isopentenyl diphosphate biosynthesis via a mevalonate-independent pathway: isopentenyl monophosphate kinase catalyzes the terminal enzymatic step. *Proc. Natl. Acad. Sci. USA* **96**, 13714-13719.
- Laule, O., Fürholz, A., Chang, H.S., Zhu, T., Wang, X., Heifetz, P.B., Gruißsem, W., Lange, M. (2003). Crosstalk between cytosolic and plastidial pathways of isoprenoid biosynthesis in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **100**, 6866-6871.
- Laupitz, R., Hecht, S., Amslinger, S., Zepeck, F., Kaiser, J., Richter, G., Schramek, N., Steinbacher, S., Huber, R., Arigoni, D., Bacher, A., Eisenreich, W., Rohdich, F. (2004). Biochemical characterization of *Bacillus subtilis* type II isopentenyl diphosphate isomerase, and phylogenetic distribution of isoprenoid biosynthesis pathways. *Eur. J. Biochem.* **271**, 2658-2669.
- Lawrence, D.L., Cline, K., Moore, G.A. (1997). Chromoplast development in ripening tomato fruit: identification of cDNAs for chromoplast-targeted proteins and characterization of a cDNA encoding a plastid-localized low-molecular-weight heat shock protein. *Plant Mol. Biol.* **33**, 483-492.
- Lell, B., Ruangweerayut, R., Wiesner, J., Missinou, M.A., Schindler, A., Baranek, T., Hintz, M., Hutchinson, D., Jomaa, H., Kremsner, P.G. (2003). Fosmidomycin, a novel chemotherapeutic agent for malaria. *Antimicrob. Agents Chemother.* **47**, 735-738.
- Lichtenthaler, H.K. (1987). Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. *Methods Enzymol.* **148**, 350-382.
- Lichtenthaler, H.K. (1999). The 1-deoxy-D-xylulose-5-phosphate pathway of isoprenoid biosynthesis in plants. *Annu. Rev. Plant Physiol. Mol. Biol.* **50**, 47-65.
- Lois, L.M., Campos, N., Putra, S.R., Danielsen, K., Rohmer, M., Boronat, A. (1998). Cloning and characterization of a gene from *Escherichia coli* encoding a transketolase-like enzyme that catalyzes the synthesis of D-1-deoxyxylulose 5-phosphate, a common precursor for isoprenoid, thiamin, and pyridoxol biosynthesis. *Proc. Natl. Acad. Sci. USA* **95**, 2105-2110.
- Lüttgen, H., Rohdich, F., Herz, S., Wungsintaweekul, J., Hecht, S., Schuhr, C.A., Fellermeier, M., Sagner, S., Zenk, M.H., Bacher, A., Eisenreich, W. (2000). Biosynthesis of terpenoids: YchB protein of *Escherichia coli* phosphorylates the 2-hydroxy group of 4-diphosphocytidyl-2C-methyl-D-erythritol. *Proc. Natl. Acad. Sci. USA* **97**, 1062-1067.
- Mandel, M., Higa, A. (1970). Calcium-dependent bacteriophage DNA infection. *J. Mol. Biol.* **53**, 159-162.
- Martin, V.J., Pitera, D.J., Withers, S.T., Newman, J.D., Keasling, J.D. (2003). Engineering a mevalonate pathway in *Escherichia coli* for production of terpenoids. *Nat. Biotechnol.* **21**, 796-802.
- McAteer, S., Coulson, A., McLennan, N., Masters, M. (2001). The *lytB* gene of *Escherichia coli* is essential and specifies a product needed for isoprenoid biosynthesis. *J. Bacteriol.* **183**, 7403-7407.

McCaskill, D., Croteau, R. (1993). Procedure for the isolation and quantification of the intermediates of the mevalonic acid pathway. *Anal. Biochem.* **215**, 142-149.

McCaskill, D., Croteau, R. (1999). Isopentenyl diphosphate is the terminal product of the deoxyxylulose-5-phosphate pathway for terpenoid biosynthesis in plants. *Tetrahedron Lett.* **40**, 653-656.

Mullis, K.B., Fallona, F.A. (1987). Specific synthesis of DNA *in vitro* via a polymerase-catalyzed reaction. *Methods Enzymol.* **155**, 335-350.

Nakamura, A., Shimada, H., Masuda, T., Ohta, H., Takamiya, K. (2001). Two distinct isopentenyl diphosphate isomerases in cytosol and plastid are differentially induced by environmental stresses in tobacco. *FEBS Lett.* **506**, 61-64.

Nicholas, K.B., Nicholas, H.B. Jr. (1997). GeneDoc: a tool for editing and annotating multiple sequence alignments. Distributed by the author.

Okuhara, M., Kuroda, Y., Goto, T., Okamoto, M., Terano, H., Kohsaka, M., Aoki, H., Imanaka, H. (1980). Studies on new phosphonic acid antibiotics. III. Isolation and characterization of FR-31564, FR-32863 and FR-33289. *J. Antibiotics* **33**, 24-28.

Page, J.E., Hause, G., Raschke, M., Gao, W., Schmidt, J., Zenk, M.H., Kutchan, T.M. (2004). Functional analysis of the final steps of the 1-deoxy-D-xylulose 5-phosphate (DXP) pathway to isoprenoids in plants using virus-induced gene silencing. *Plant Physiol.* **134**, 1401-1413.

Patterson, D.R. (1987). Rohm and Haas Co., US Patent 4693742.

Porath, J., Carlsson, J., Olsson, I., Belfrage, G. (1975). Metal chelate affinity chromatography, a new approach to protein fractionation. *Nature* **258**, 598-599.

Putra, S.R., Lois, L.M., Campos, N., Boronat, A., Rohmer, M. (1998). Incorporation of [2,3-¹³C₂]- and [2,4-¹³C₂]-D-1-deoxyxylulose into ubiquinone of *Escherichia coli* via the mevalonate-independent pathway for isoprenoid biosynthesis. *Tetrahedron Lett.* **39**, 23-26.

Qureshi, N., Porter, J.W. (1981). Conversion of acetyl-coenzyme A to isopentenyl pyrophosphate. *Biosynthesis of Isoprenoid Compounds* **1**, 47-94, Porter, J.W. and Spurgeon, S.L. (eds), Wiley New York.

Ralph, S.A., Van Dooren, G.G., Waller, R.F., Crawford, M.J., Fraunholz, M.J., Foth, B.J., Tonkin, C.J., Roos, D.S., McFadden, G.I. (2004). Tropical infectious diseases: metabolic maps and functions of the *Plasmodium falciparum* apicoplast. *Nat. Rev. Microbiol.* **2**, 203-216.

Ramos-Valdivia, A.C., van der Heijden, R., Verpoorte, R. (1997a). Isopentenyl diphosphate isomerase: a core enzyme in isoprenoid biosynthesis. A review of its biochemistry and function. *Nat. Prod. Rep.* **14**, 591-603.

Ramos-Valdivia, A.C., van der Heijden, R., Verpoorte, R., Camara, B. (1997b). Purification and characterization of two isoforms of isopentenyl-diphosphate isomerase from elicitor-treated *Cinchona robusta* cells. *Eur. J. Biochem.* **249**, 161-170.

- Raschke, M., Fellermeier, M., Zenk, M.H. (2004). A high-performance liquid chromatography method for the analysis of intermediates of the deoxyxylulose phosphate pathway. *Anal. Biochem.* **335**, 235-243.
- Ratcliff, F., Martin-Hernandez, A.M., Baulcombe, D.C. (2001). Tobacco rattle virus as a vector for analysis of gene function by silencing. *Plant J.* **25**, 237-245.
- Robertson, D. (2004). VIGS vectors for gene silencing: many targets, many tools. *Annu. Rev. Plant Biol.* **55**, 495-519.
- Rodriguez-Concepcion, M., Campos, N., Lois, L.M., Maldonado, C., Hoeffler, J.F., Grosdemange-Billiard, C., Rohmer, M., Boronat, A. (2000). Genetic evidence of branching in the isoprenoid pathway for the production of isopentenyl diphosphate and dimethylallyl diphosphate in *Escherichia coli*. *FEBS Lett.* **473**, 328-332.
- Rodriguez-Concepcion, M., Boronat, A. (2002). Elucidation of the methylerythritol phosphate pathway for isoprenoid biosynthesis in bacteria and plastids. A metabolic milestone achieved through genomics. *Plant Physiol.* **130**, 1079-1089.
- Rodriguez-Concepcion, M., Fores, O., Martinez-Garcia, J.F., Gonzalez, V., Phillips, M.A., Ferrer, A., Boronat, A. (2004). Distinct light-mediated pathways regulate the biosynthesis and exchange of isoprenoid precursors during Arabidopsis seedling development. *Plant Cell* **16**, 144-156.
- Rohdich, F., Wungsintaweekul, J., Fellermeier, M., Sagner, S., Herz, S., Kis, K., Eisenreich, W., Bacher, A., Zenk, M.H. (1999). Cytidine 5'-triphosphate-dependent biosynthesis of isoprenoids: YgbP protein of *Escherichia coli* catalyzes the formation of 4-diphosphocytidyl-2C-methylerythritol. *Proc. Natl. Acad. Sci. USA* **96**, 11758-11763.
- Rohdich, F., Wungsintaweekul, J., Eisenreich, W., Richter, G., Schuhr, C.A., Hecht, S., Zenk, M.H., Bacher, A. (2000a). Biosynthesis of terpenoids: 4-diphosphocytidyl-2C-methyl-D-erythritol synthase of *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **97**, 6451-6456.
- Rohdich, F., Wungsintaweekul, J., Lüttgen, H., Fischer, M., Eisenreich, W., Schuhr, C.A., Fellermeier, M., Schramek, N., Zenk, M.H., Bacher, A. (2000b). Biosynthesis of terpenoids: 4-diphosphocytidyl-2C-methyl-D-erythritol kinase from tomato. *Proc. Natl. Acad. Sci. USA* **97**, 8251-8256.
- Rohdich, F., Eisenreich, W., Wungsintaweekul, J., Hecht, S., Schuhr, C.A., Bacher, A. (2001a). Biosynthesis of terpenoids. 2C-Methyl-D-erythritol 2,4-cyclodiphosphate synthase (IspF) from *Plasmodium falciparum*. *Eur. J. Biochem.* **268**, 3190-3197.
- Rohdich, F., Kis, K., Bacher, A., Eisenreich, W. (2001b). The non-mevalonate pathway of isoprenoids: genes, enzymes and intermediates. *Curr. Opin. Chem. Biol.* **5**, 535-540.
- Rohdich, F., Hecht, S., Gärtner, K., Adam, P., Krieger, C., Amslinger, S., Arigoni, D., Bacher, A., Eisenreich, W. (2002). Studies on the nonmevalonate terpene biosynthetic pathway: metabolic role of IspH (LytB) protein. *Proc. Natl. Acad. Sci. USA* **99**, 1158-1163.

Rohdich, F., Zepeck, F., Adam, P., Hecht, S., Kaiser, J., Laupitz, R., Gräwert, T., Amslinger, S., Eisenreich, W., Bacher, A., Arigoni, D. (2003). The deoxyxylulose phosphate pathway of isoprenoid biosynthesis: studies on the mechanisms of the reactions catalyzed by IspG and IspH protein. *Proc. Natl. Acad. Sci. USA* **100**, 1586-1591.

Rohdich, F., Bacher, A., Eisenreich, W. (2004). Perspectives in anti-infective drug design. The late steps in the biosynthesis of the universal terpenoid precursors, isopentenyl diphosphate and dimethylallyl diphosphate. *Bioorg. Chem.* **32**, 292-308.

Rohmer, M., Knani, M., Simonin, P., Sutter, B., Sahm, H. (1993). Isoprenoid biosynthesis in bacteria: a novel pathway for the early steps leading to isopentenyl diphosphate. *Biochem. J.* **295**, 517-524.

Rohmer, M. (1999a). The discovery of a mevalonate-independent pathway for isoprenoid biosynthesis in bacteria, algae and higher plants. *Nat. Prod. Rep.* **16**, 565-574.

Rohmer, M. (1999b). A mevalonate-independent route to isopentenyl diphosphate. *Comprehensive Natural Products Chemistry. Isoprenoids Including Carotenoids and Steroids*. D.E. Cane (Ed.) **2**, 45-67, Pergamon, Oxford, UK.

Roos, D.S., Crawford, M.J., Donald, R.G., Fraunholz, M., Harb, O.S., He, C.Y., Kissinger, J.C., Shaw, M.K., Striepen, B. (2002). Mining the Plasmodium genome database to define organellar function: what does the apicoplast do? *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **357**, 35-46.

Sacchettini, J.C., Poulter, C.D. (1997). Creating isoprenoid diversity. *Science* **277**, 1788-1789.

Sagner, S., Latzel, C., Eisenreich, W., Bacher, A., Zenk, M.H. (1998). Differential incorporation of 1-deoxy-D-xylulose into monoterpenes and carotenoids in higher plants. *J. Chem. Soc. Chem. Commun.*, 221-222.

Saiki, R.K., Scharf, S., Faloona, F., Mullis, K.B., Horn, G.T., Erlich, H.A., Arnheim, N. (1985). Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science* **230**, 1350-1354.

Sambrook, J., Russell, D.W. (2001). *Molecular cloning: A laboratory manual*, 3rd edition. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.

Sanger, F., Nicklen, S., Coulson, A.R. (1977). DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**, 5463-5467.

Satterwhite, D.M. (1985). Isopentenyl diphosphate isomerase. *Methods Enzymol.* **110**, 92-99.

Schreyer, R., Böck, A. (1973). Phenotypic suppression of a fructose-1,6-diphosphate aldolase mutation in *Escherichia coli*. *J. Bacteriol.* **115**, 268-276.

Schuhr, C.A., Hecht, S., Kis, K., Eisenreich, W., Wungsintaweeikul, J., Bacher, A., Rohdich, F. (2001). Studies on the non-mevalonate pathway - preparation and properties of isotope-labeled 2C-methyl-D-erythritol 2,4-cyclodiphosphate. *Eur. J. Org. Chem.*, 3221-3226.

- Schuhr, C.A., Radykewicz, T., Sagner, S., Latzel, C., Zenk, M.H., Arigoni, D., Bacher, A., Rohdich, F., Eisenreich, W. (2003). Quantitative assessment of crosstalk between the two isoprenoid biosynthesis pathways in plants by NMR spectroscopy. *Phytochem. Reviews*. **2**, 3-16.
- Schwarz, M.K. (1994). Terpen-Biosynthese in *Ginkgo biloba*: Eine überraschende Geschichte. Dissertation Nr. 10951, ETH Zürich, Schweiz.
- Schwender, J., Müller, C., Zeidler, J., Lichtenthaler, H.K. (1999). Cloning and heterologous expression of a cDNA encoding 1-deoxy-D-xylulose-5-phosphate reductoisomerase of *Arabidopsis thaliana*. *FEBS Lett.* **455**, 140-144.
- Schwenkreis, P., Tegenthoff, M. (2003). Therapeutic use of cannabinoids in neurology. *Schmerz* **17**, 367-373.
- Seemann, M., Bui, B.T., Wolff, M., Tritsch, D., Campos, N., Boronat, A., Marquet, A., Rohmer, M. (2002). Isoprenoid biosynthesis through the methylerythritol phosphate pathway: the (E)-4-hydroxy-3-methylbut-2-enyl diphosphate synthase (GcpE) is a [4Fe-4S] protein. *Angew. Chem. Int. Ed.* **41**, 4337-4339.
- Seemann, M., Wegner, P., Schünemann, V., Bui, B.T., Wolff, M., Marquet, A., Trautwein, A.X., Rohmer, M. (2005). Isoprenoid biosynthesis in chloroplasts via the methylerythritol phosphate pathway: the (E)-4-hydroxy-3-methylbut-2-enyl diphosphate synthase (GcpE) from *Arabidopsis thaliana* is a [4Fe-4S] protein. *J. Biol. Inorg. Chem.* **10**, 131-137.
- Seto, H., Watanabe, H., Furihata, K. (1996). Simultaneous operation of the mevalonate and non-mevalonate pathway in the biosynthesis of isopentenyl diphosphate in *Streptomyces aeriovifer*. *Tetrahedron Lett.* **39**, 7979-7982.
- Shigi, Y. (1989). Inhibition of bacterial isoprenoid biosynthesis by fosmidomycin, a phosphonic acid-containing antibiotic. *J. Antimicrob. Chemoth.* **24**, 131-145.
- Sprenger, G.A., Schörken, U., Wiegert, T., Grolle, S., de Graaf, A.A., Taylor, S.V., Begley, T.P., Bringer-Meyer, S., Sahn, H. (1997). Identification of a thiamin-dependent synthase in *Escherichia coli* required for the formation of the 1-deoxy-D-xylulose 5-phosphate precursor to isoprenoids, thiamin, and pyridoxol. *Proc. Natl. Acad. Sci. USA* **94**, 12857-12862.
- Stahl, E., Kaltenbach, U. (1961). Thin-layer chromatography. VI. Trace analysis of sugar mixtures on layers of Kieselgur G. *J. Chromatog.* **5**, 351-355.
- Stock, I. (2004). Therapie der Malaria. *Med. Monatsschr. Pharm.* **27**, 260-272.
- Street, I.P., Poulter, C.D. (1990). Isopentenyl diphosphate:dimethylallyl diphosphate isomerase: construction of a high-level heterologous expression system for the gene from *Saccharomyces cerevisiae* and identification of an active-site nucleophile. *Biochemistry* **29**, 7531-7538.
- Street, I.P., Coffman, H.R., Baker, J.A., Poulter, C.D. (1994). Identification of Cys139 and Glu207 as catalytically important groups in the active site of isopentenyl diphosphate:dimethylallyl diphosphate isomerase. *Biochemistry* **33**, 4212-4217.

Takagi, M., Kuzuyama, T., Kaneda, K., Watanabe, H., Dairi, T., Seto, H. (2000). Studies on the nonmevalonate pathway: formation of 2C-methyl-D-erythritol 2,4-cyclodiphosphate from 2-phospho-4-(cytidine 5'-diphospho)-2C-methyl-D-erythritol. *Tetrahedron Lett.* **41**, 3395-3398.

Takahashi, S., Kuzuyama, T., Watanabe, H., Seto, H. (1998). A 1-deoxy-D-xylulose 5-phosphate reductoisomerase catalyzing the formation of 2C-methyl-D-erythritol 4-phosphate in an alternative nonmevalonate pathway for terpenoid biosynthesis. *Proc. Natl. Acad. Sci. USA* **95**, 9879-9884.

Taylor, S.V., Vu, L.D., Begley, T.P. (1998). Chemical and enzymatic synthesis of 1-deoxy-D-xylulose-5-phosphate. *J. Org. Chem.* **63**, 2375-2377.

Thiel, R., Adam, K.P. (1999). Synthesis of [5,5-²H₂]-1-deoxy-D-xylulose 5-phosphate. *Tetrahedron Lett.* **43**, 5307-5308.

Tholl, D., Croteau, R., Gershenzon, J. (2001). Partial purification and characterization of the short-chain prenyltransferases, geranyl diphosphate synthase and farnesyl diphosphate synthase, from *Abies grandis* (grand fir). *Arch. Biochem. Biophys.* **386**, 233-242.

Thompson, J.D., Higgins, D.G., Gibson, T.J. (1994). CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**, 4673-4680.

Van de Peer, Y., De Wachter, R. (1994). TREECON for Windows: a software package for the construction and drawing of evolutionary trees for the Microsoft Windows environment. *Comput. Applic. Biosci.* **10**, 569-570.

White, R.H. (1978). Stable isotope studies on the biosynthesis of the thiazole moiety of thiamin in *Escherichia coli*. *Biochemistry* **17**, 3833-3840.

Wiesner, J., Ortmann, R., Jomaa, H., Schlitzer, M. (2003). New antimalarial drugs. *Angew. Chem. Int. Ed.* **42**, 5274-5293.

Wolff, M., Seemann, M., Bui, B.T., Frapart, Y., Tritsch, D., Estrabot, A.G., Rodriguez-Concepcion, M., Boronat, A., Marquet, A., Rohmer, M. (2003). Isoprenoid biosynthesis via the methylerythritol phosphate pathway: the (E)-4-hydroxy-3-methylbut-2-enyl diphosphate reductase (LytB/IspH) from *Escherichia coli* is a [4Fe-4S] protein. *FEBS Lett.* **541**, 115-120.

Wong, S.C.K.; Hall, D.C., Josse, J. (1970). Constitutive inorganic pyrophosphatase of *Escherichia coli*. 3. Molecular weight and physical properties of the enzyme and its subunits. *J. Biol. Chem.* **245**, 4335-4345.

Wouters, J., Oudjama, Y., Barkley, S.J., Tricot, C., Stalon, V., Droogmans, L., Poulter, C.D. (2003). Catalytic mechanism of *Escherichia coli* isopentenyl diphosphate isomerase involves Cys-67, Glu-116, and Tyr-104 as suggested by crystal structures of complexes with transition state analogues and irreversible inhibitors. *J. Biol. Chem.* **278**, 11903-11908.

Wungsintaweekul, J. (2001a). Enzymes of the alternative terpenoid pathway in *Escherichia coli*. Dissertation, Ludwig-Maximilians-Universität München.

Wungsintaweekul, J., Herz, S., Hecht, S., Eisenreich, W., Feicht, R., Rohdich, F., Bacher, A., Zenk, M.H. (2001b). Phosphorylation of 1-deoxy-D-xylulose by D-xylulokinase of *Escherichia coli*. *Eur. J. Biochem.* **268**, 310-316.

Zhang, F.L., Casey, P.J. (1996). Protein prenylation: molecular mechanisms and functional consequences. *Annu. Rev. Biochem.* **65**, 241-269.

Zeidler, J., Schwender, J., Müller, C., Wiesner, J., Weidemeyer, C., Beck, E., Jomaa, H., Lichtenthaler, H.K. (1998). Inhibition of the non-mevalonate 1-deoxy-D-xylulose-5-phosphate pathway of plant isoprenoid biosynthesis by fosmidomycin. *Z. Naturforsch.* **53c**, 980-986.