

4. Literatur

- Aarts, N., Metz, M., Holub, E., Staskawicz, B.J., Daniels, M.J., and Parker, J.E.** (1998). Different requirements for *EDS1* and *NDR1* by disease resistance genes define at least two *R* gene-mediated signaling pathways in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **95**, 10306-10311.
- Abramovitch, R.B., Janjusevic, R., Stebbins, C.E., and Martin, G.B.** (2006). Type III effector AvrPtoB requires intrinsic E3 ubiquitin ligase activity to suppress plant cell death and immunity. *Proc. Natl. Acad. Sci. USA* **103**, 2851-2856.
- Albrecht, M., and Takken, F.L.** (2006). Update on the domain architectures of NLRs and R proteins. *Biochemical and Biophys. Res. Commun.* **339**, 459-462.
- Austin, M.J., Muskett, P., Kahn, K., Feys, B.J., Jones, J.D.G., and Parker, J.E.** (2002). Regulatory Role of *SGT1* in early *R* gene-mediated plant defences. *Science* **295**, 2077-2080.
- Axtell, M.J., and Staskawicz, B.J.** (2003). Initiation of *RPS2*-specified disease resistance in *Arabidopsis* is coupled to the AvrRpt2-directed elimination of RIN4. *Cell* **112**, 369-377.
- Axtell, M.J., Chisholm, S.T., Dahlbeck, D., and Staskawicz, B.J.** (2003). Genetic and molecular evidence that the *Pseudomonas syringae* type III effector protein AvrRpt2 is a cysteine protease. *Mol. Microbiol.* **49**, 1537-1546.
- Azevedo, C., Santos-Rosa, M.J., and Shirasu, K.** (2001). The U-box protein family in plants. *Trends Plant Sci.* **6**, 354-358.
- Azevedo, C., Sadanandom, A., Kitagawa, K., Freialdenhoven, A., Shirasu, K., and Schulze-Lefert, P.** (2002). The RAR1 interactor SGT1, an essential component of R gene-triggered disease resistance. *Science* **295**, 2073-2076.
- Bai, J., Choi, S.-H., Ponciano, G., Leung, H., and Leach, J.E.** (2000). *Xanthomonas oryzae* pv. *oryzae* avirulence genes contribute differently and specifically to pathogen aggressiveness. *Mol. Plant-Microbe Interact.* **13**, 1322-1329.
- Ballvora, A., Pierre, M., Van den Ackerveken, G., Schornack, S., Rossier, O., Ganai, M., Lahaye, T., and Bonas, U.** (2001). Genetic mapping and functional analysis of the tomato *Bs4* locus, governing recognition of the *Xanthomonas campestris* pv. *vesicatoria* AvrBs4 protein. *Mol. Plant-Microbe Interact.* **14**, 629-638.
- Bendahmane, A., Kanyuka, K., and Baulcombe, D.C.** (1999). The *Rx* gene from potato controls separate virus resistance and cell death responses. *Plant Cell* **11**, 781-791.
- Bendahmane, A., Farnham, G., Moffett, P., and Baulcombe, D.C.** (2002). Constitutive gain-of-function mutants in a nucleotide binding site-leucine rich repeat protein encoded at the *Rx* locus of potato. *Plant J.* **32**, 195-204.
- Bent, A.F., Kunkel, B.N., Dahlbeck, D., Brown, K.L., Schmidt, R., Giraudat, L.J., and Staskawicz, B.J.** (1994). *RPS2* of *Arabidopsis thaliana* : a leucine-rich repeat class of disease resistance genes. *Science* **265**, 1856-1860.
- Bieri, S., Mauch, S., Shen, Q.-H., Peart, J., Devoto, A., Casais, C., Ceron, F., Schulze, S., Steinbiß, H.-H., Shirasu, K., and Schulze-Lefert, P.** (2004). RAR1 positively controls steady state levels of barley MLA resistance proteins and enables sufficient MLA6 accumulation for effective resistance. *Plant Cell* **16**, 3480-3495.
- Biffen, R.H.** (1905). Mendel's laws of inheritance and wheat breeding. *J Agr. Sci.* **1**, 1-48.
- Blair, M.W., Garris, A.J., Iyer, A.S., Chapman, B., Kresovich, S., and McCouch, S.R.** (2003). High resolution genetic mapping and candidate gene identification at the

- xa5* locus for bacterial blight resistance in rice (*Oryza sativa* L.). *Theor. Appl. Genet.* **107**, 62-73.
- Blatch, G.L., and Lässle, M.** (1999). The tetratricopeptide repeat: a structural motif mediating protein-protein interactions. *Bioessays* **21**, 932-939.
- Bohs, L., and Olmstead, R.** (1997). Phylogenetic relationships in *Solanum* (Solanaceae) based on *ndhF* sequences. *Syst. Botany* **22**, 5-17.
- Bonas, U., and Lahaye, T.** (2002). Plant disease resistance triggered by pathogen-derived molecules: refined models of specific recognition. *Curr. Opin. Microbiol.* **5**, 44-50.
- Bonas, U., Stall, R.E., and Staskawicz, B.** (1989). Genetic and structural characterization of the avirulence gene *avrBs3* from *Xanthomonas campestris* pv. *vesicatoria*. *Mol. Gen. Genet.* **218**, 127-136.
- Bonas, U., Conrads-Strauch, J., and Balbo, I.** (1993). Resistance in tomato to *Xanthomonas campestris* pv. *vesicatoria* is determined by alleles of the pepper-specific avirulence gene *avrBs3*. *Mol. Gen. Genet.* **238**, 261-269.
- Boyce, D.C., Nam, J., and Dangl, J.L.** (1998). The *Arabidopsis thaliana* *RPM1* disease resistance gene product is a peripheral plasma membrane protein that is degraded coincident with the hypersensitive response. *Proc. Natl. Acad. Sci. USA* **95**, 15849-15854.
- Bracha-Drori, K., Shichrur, K., Katz, A., Oliva, M., Angelovici, R., Yalovsky, S., and Ohad, N.** (2004). Detection of protein-protein interactions in plants using bimolecular fluorescence complementation. *Plant J.* **40**, 419-427.
- Bustin, S.A.** (2000). Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays. *J. Mol. Endocrin.* **25**, 169-193.
- Büttner, D., and Bonas, U.** (2006). Who comes first? How plant pathogenic bacteria orchestrate type III secretion. *Curr. Opin. Microbiol.* **9**, 193-200.
- Canteros, B., Minsavage, G., Bonas, U., Pring, D., and Stall, R.** (1991). A gene from *Xanthomonas campestris* pv. *vesicatoria* that determines avirulence in tomato is related to *avrBs3*. *Mol. Plant-Microbe Interact.* **4**, 628-632.
- Century, K.S., Shapiro, A.D., Repetti, P.P., Dahlbeck, D., Holub, E., and Staskawicz, B.J.** (1997). *NDR1*, a pathogen-induced component required for *Arabidopsis* disease resistance. *Science* **278**, 1963-1965.
- Chu, Z., Yuan, M., Yao, J., Ge, X., Yuan, B., Xu, C., Li, X., Fu, B., Li, Z., Bennetzen, J.L., Zhang, Q., and Wang, S.** (2006). Promoter mutations of an essential gene for pollen development result in disease resistance in rice. *Genes Dev.* **20**, 1250-1255.
- Coppinger, P., Repetti, P.P., Day, B., Dahlbeck, D., Mehlert, A., and Staskawicz, B.J.** (2004). Overexpression of the plasma membrane-localized NDR1 protein results in enhanced bacterial disease resistance in *Arabidopsis thaliana*. *Plant J.* **40**, 225-237.
- Cunnac, S., Occhialini, A., Barberis, P., Boucher, C., and Genin, S.** (2004). Inventory and functional analysis of the large Hrp regulon in *Ralstonia solanacearum*: identification of novel effector proteins translocated to plant host cells through the type III secretion system. *Mol. Microbiol.* **53**, 115-128.
- Dangl, J.L., and Jones, J.D.G.** (2001). Plant pathogens and integrated defence responses to infection. *Nature* **411**, 826-833.
- Dangl, J.L., and McDowell, J.M.** (2006). Two modes of pathogen recognition by plants. *Proc. Natl. Acad. Sci. USA* **103**, 8575-8576.
- De Feyter, R., and Gabriel, D.W.** (1991). At least six avirulence genes are clustered on a 90-kilobase plasmid in *Xanthomonas campestris* pv. *malvacearum*. *Mol. Plant-Microbe Interact.* **4**, 423-432.

- De Feyter, R., Yang, Y.O., and Gabriel, D.W.** (1993). Gene-for-genes interactions between cotton *R*-genes and *Xanthomonas campestris* pv. *malvacearum* avr genes. *Mol. Plant-Microbe Interact.* **6**, 225-237.
- De Feyter, R., McFadden, H., and Dennis, L.** (1998). Five avirulence genes from *Xanthomonas campestris* pv. *malvacearum* cause genotype-specific cell death when expressed transiently in cotton. *Mol. Plant-Microbe Interact.* **11**, 698-701.
- De Ilarduya, O.M., and Kaloshian, I.** (2001). *Mi-1.2* transcripts accumulate ubiquitously in resistant *Lycopersicon esculentum*. *J. Nematol.* **33**, 116-120.
- de la Fuente van Bentem, S., Vossen, J.H., de Vries, K.J., van Wees, S., Tameling, W.I., Dekker, H.L., de Koster, C.G., Haring, M.A., Takken, F.L., and Cornelissen, B.J.** (2005). Heat shock protein 90 and its co-chaperone protein phosphatase 5 interact with distinct regions of the tomato I-2 disease resistance protein. *Plant J.* **43**, 284-298.
- de la Fuente van Bentem, S., Vossen, J.H., Haring, M.A., and Cornelissen, B.J.C.** (2003). Tomato protein phosphatase 5 interacts with the LRR domain of multiple plant resistance gene products in the yeast two-hybrid system. meeting report
- Deslandes, L., Olivier, J., Theulieres, F., Hirsch, J., Feng, D.X., Bittner-Eddy, P., Beynon, J., and Marco, Y.** (2002). Resistance to *Ralstonia solanacearum* in *Arabidopsis thaliana* is conferred by the recessive *RRS1-R* gene, a member of a novel family of resistance genes. *Proc. Natl. Acad.Sci. USA* **99**, 2404-2409.
- Deslandes, L., Olivier, J., Peeters, N., Feng, D.X., Khounlotham, M., Boucher, C., Somssich, I., Genin, S., and Marco, Y.** (2003). Physical interaction between *RRS1-R*, a protein conferring resistance to bacterial wilt, and PopP2, a type III effector targeted to the plant nucleus. *Proc. Natl. Acad.Sci. USA* **100**, 8024-8029.
- Dinesh-Kumar, S.P., Tham, W.H., and Baker, B.J.** (2000). Structure-function analysis of the tobacco mosaic virus resistance gene *N*. *Proc. Natl. Acad.Sci. USA* **97**, 14789-14794.
- Dixon, R.A., and Harrison, M.J.** (1990). Activation, structure, and organization of genes involved in microbial defense in plants. *Adv. Genet.* **28**, 165-234.
- Dodds, P.N., Lawrence, G.J., and Ellis, J.G.** (2001). Six amino acid changes confined to the leucine-rich repeat β -strand/ β -turn motif determine the difference between the P and P2 rust resistance specificities in flax. *Plant Cell* **13**, 163-178.
- Dodds, P.N., Lawrence, G.J., Catanzariti, A.M., Teh, T., Wang, C.I., Ayliffe, M.A., Kobe, B., and Ellis, J.G.** (2006). From the Cover: Direct protein interaction underlies gene-for-gene specificity and coevolution of the flax resistance genes and flax rust avirulence genes. *Proc. Natl. Acad.Sci. USA* **103**, 8888-8893.
- Doidge, E.** (1921). A tomato canker. *Ann. Appl. Biol.* **7**, 407-430.
- Durrant, W.E., and Dong, X.** (2004). Systemic acquired resistance. *Ann. Rev. Phytopathol.* **42**, 185-209.
- Dye, D.** (1978). Genus IX. *Xanthomonas*. Dowson 1939. In A proposed nomenclature and classification for plant pathogenic bacteria, M. Young, D. Dye, F. Bradbury, C. Panagopoulos, und C. Robbs, eds, pp. 153-177.
- Ellis, J., Catanzariti, A.M., and Dodds, P.** (2006). The problem of how fungal and oomycete avirulence proteins enter plant cells. *Trends Plant Sci.* **11**, 61-63.
- Erickson, F.L., Holzberg, S., Calderon-Urrea, A., Handley, V., Axtell, M., Corr, C., and Baker, B.** (1999). The helicase domain of the TMV replicase proteins induces the *N*-mediated defence response in tobacco. *Plant J.* **18**, 67-75.

- Ezuka, A., Horino, O., Toriyama, K., Shinoda, H., and Morinaka, T.** (1975). Inheritance of resistance of rice variety Wase Aikoku 3 to *Xanthomonas oryzae*. Bull. Tokai-kinki Nat. Agric. Exp. Stn. **28**, 124-130.
- Falk, A., Feys, B.J., Frost, L.N., Jones, J.D.G., Daniels, M.J., and Parker, J.E.** (1999). *EDS1*, an essential component of *R* gene-mediated disease resistance in *Arabidopsis* has homology to eukaryotic lipases. Proc. Natl. Acad. Sci. USA **96**, 3292-3297.
- Flor, H.H.** (1942). Inheritance of pathogenicity in *Melampsora lini*. Phytopathology **32**, 653-669.
- Flor, H.H.** (1971). Current status of the gene-for-gene concept. Ann. Rev. Phytopathol. **9**, 275-296.
- Gabriel, D.W., and Rolfe, B.G.** (1990). Working models of specific recognition in plant-microbe interactions. Ann. Rev. Phytopathol. **28**, 365-391.
- Gabriel, D.W., Burges, A., and Lazo, G.R.** (1986). Gene-for-gene interactions of five cloned avirulence genes from *Xanthomonas campestris* pv *malvacearum* with specific resistance genes in cotton. Proc. Natl. Acad. Sci. USA **83**, 6415-6419.
- Gardner, M., and Kendrick, J.** (1921). Bacterial spot of tomato. J. Agr. Res. **21**, 123-156.
- Ginalski, K., Elofsson, A., Fischer, D., and Rychlewski, L.** (2003). 3D-Jury: a simple approach to improve protein structure predictions. Bioinformatics **19**, 1015-1018.
- Goggin, F.L., Shah, G., Williamson, V.M., and Ullman, D.E.** (2004). Developmental regulation of *Mi*-mediated aphid resistance is independent of *Mi-1.2* transcript levels. Mol. Plant-Microbe Interact. **17**, 532-536.
- Gómez-Gómez, L., and Boller, T.** (2000). FLS2: An LRR receptor-like kinase involved in the perception of the bacterial elicitor flagellin in *Arabidopsis*. Mol. Cell **5**, 1003-1011.
- Gómez-Gómez, L., Felix, G., and Boller, T.** (1999). A single locus determines sensitivity to bacterial flagellin in *Arabidopsis thaliana*. Plant J. **18**, 277-284.
- Grant, J.J., Chini, A., Basu, D., and Loake, G.J.** (2003). Targeted activation tagging of the *Arabidopsis* NBS-LRR gene, *ADR1*, conveys resistance to virulent pathogens. Mol. Plant-Microbe Interact. **16**, 669-680.
- Grant, M.R., Godiard, L., Straube, E., Ashfield, T., Lewald, J., Sattler, A., Innes, R.W., and Dangl, J.L.** (1995). Structure of the *Arabidopsis RPM1* gene enabling dual specificity disease resistance. Science **269**, 843-846.
- Greenberg, J.T., and Yao, N.** (2004). The role and regulation of programmed cell death in plant-pathogen interactions. Cell. Microbiol. **6**, 201-211.
- Gu, K., Yang, B., Tian, D., Wu, L., Wang, D., Sreekala, C., Yang, F., Chu, Z., Wang, G.L., White, F.F., and Yin, Z.** (2005). *R* gene expression induced by a type-III effector triggers disease resistance in rice. Nature **435**, 1122-1125.
- Gürlebeck, D., Szurek, B., and Bonas, U.** (2005). Dimerization of the bacterial effector protein AvrBs3 in the plant cell cytoplasm prior to nuclear import. Plant J. **42**, 175-187.
- Gürlebeck, D., Thieme, F., and Bonas, U.** (2006). Type III effector proteins from the plant pathogen *Xanthomonas* and their role in the interaction with the host plant. J. Plant Physiol. **163**, 233-255.
- Hammond-Kosack, K.E., and Jones, J.D.G.** (1996). Resistance gene-dependent plant defense responses. Plant Cell **8**, 1773-1791.
- Heath, M.C.** (2000). Nonhost resistance and nonspecific plant defenses. Curr. Opin. Plant Biol. **3**, 315-319.

- Herbers, K., Conrads-Strauch, J., and Bonas, U.** (1992). Race-specificity of plant resistance to bacterial spot disease determined by repetitive motifs in a bacterial avirulence protein. *Nature* **356**, 172-174.
- Hibberd, A.M., Bassett, M.J., and Stall, R.E.** (1987). Allelism tests of three dominant genes for hypersensitive resistance to bacterial spot of pepper. *Phytopathology* **77**, 1304-1307.
- Holt, B.F., 3rd, Belkhadir, Y., and Dangl, J.L.** (2005). Antagonistic control of disease resistance protein stability in the plant immune system. *Science* **309**, 929-932.
- Hopkins, C.M., White, F.F., Choi, S.H., Guo, A., and Leach, J.E.** (1992). Identification of a family of avirulence genes from *Xanthomonas oryzae* pv. *oryzae*. *Mol. Plant-Microbe Interact.* **5**, 451-459.
- Howles, P., Lawrence, G., Finnegan, J., McFadden, H., Ayliffe, M., Dodds, P., and Ellis, J.** (2005). Autoactive alleles of the flax L6 rust resistance gene induce non-race-specific rust resistance associated with the hypersensitive response. *Mol. Plant-Microbe Interact.* **18**, 570-582.
- Hueck, C.J.** (1998). Type III protein secretion systems in bacterial pathogens of animals and plants. *Microbiol. Mol. Biol. Rev.* **62**, 379-433.
- Hwang, C.-F., and Williamson, V.M.** (2003). Leucine-rich repeat-mediated intramolecular interactions in nematode recognition and cell death signaling by the tomato resistance protein Mi. *Plant J.* **34**, 585-593.
- Iyer, A.S., and McCouch, S.R.** (2004). The rice bacterial blight resistance gene *xa5* encodes a novel form of disease resistance. *Mol. Plant-Microbe Interact.* **17**, 1348-1354.
- Jamir, Y., Guo, M., Oh, H.-S., Petnicki-Ocwieja, T., Chen, S., Tang, X., Dickman, M.B., Collmer, A., and R. Alfano, J.** (2004). Identification of *Pseudomonas syringae* type III effectors that can suppress programmed cell death in plants and yeast. *Plant J.* **37**, 554-565.
- Janjusevic, R., Abramovitch, R.B., Martin, G.B., and Stebbins, C.E.** (2006). A bacterial inhibitor of host programmed cell death defenses is an E3 ubiquitin ligase. *Science* **311**, 222-226.
- Jia, Y., McAdams, S.A., Bryan, G.T., Hershey, H.P., and Valent, B.** (2000). Direct interaction of resistance gene and avirulence gene products confers rice blast resistance. *EMBO J.* **19**, 4004-4014.
- Jinek, M., Rehwinkel, J., Lazarus, B.D., Izaurralde, E., Hanover, J.A., and Conti, E.** (2004). The superhelical TPR-repeat domain of O-linked GlcNAc transferase exhibits structural similarities to importin alpha. *Nat. Struct. Mol. Biol.* **11**, 1001-1007.
- Johnsson, N., and Varshavsky, A.** (1994). Split ubiquitin as a sensor of protein interactions *in vivo*. *Proc. Natl. AcadSci. USA* **91**, 10340-10344.
- Jones, D.A., Thomas, C.M., Hammond-Kosack, K.E., Balint-Kurti, P.J., and Jones, J.D.G.** (1994). Isolation of the tomato *Cf-9* gene for resistance to *Cladosporium fulvum* by transposon tagging. *Science* **266**, 789-793.
- Kawasaki, T., Nam, J., Boyes, D.C., Holt, B.F., Hubert, D.A., Wiig, A., and Dangl, J.L.** (2005). A duplicated pair of Arabidopsis RING-finger E3 ligases contribute to the RPM1- and RPS2-mediated hypersensitive response. *The Plant J.* **44**, 258-270.
- Kay, S., Boch, J., and Bonas, U.** (2005). Characterization of AvrBs3-like effectors from a Brassicaceae pathogen reveals virulence and avirulence activities and a protein with a novel repeat architecture. *Mol. Plant-Microbe Interact.* **18**, 838-848.

- Keen, N.T.** (1990). Gene-for-gene complementarity in plant-pathogen interactions. *Ann. Rev. Genet.* **24**, 447-463.
- Kerppola, T.K.** (2006). Visualization of molecular interactions by fluorescence complementation. *Nat. Rev. Mol. Cell Biol.* **7**, 449-456.
- Kim, M.G., da Cunha, L., McFall, A.J., Belkhadir, Y., DeRoy, S., Dangl, J.L., and Mackey, D.** (2005). Two *Pseudomonas syringae* type III effectors inhibit RIN4-regulated basal defense in *Arabidopsis*. *Cell* **121**, 749-759.
- Klement, Z.** (1963). Rapid detection of the pathogenicity of phytopathogenic pseudomonads. *Nature* **199**, 299-300.
- Knapp, S.** (2002). Tobacco to tomatoes: a phylogenetic perspective on fruit diversity in the Solanaceae. *J. Exp. Bot.* **53**, 2001-2022.
- Kobe, B., and Deisenhofer, J.** (1995). A structural basis of the interactions between leucine rich repeats and protein ligands. *Nature* **374**, 183-186.
- Krüger, J., Thomas, C.M., Golstein, C., Dixon, M.S., Smoker, M., Tang, S., Mulder, L., and Jones, J.D.** (2002). A tomato cysteine protease required for *Cf-2*-dependent disease resistance and suppression of autonecrosis. *Science* **296**, 744-747.
- Lahaye, T., and Bonas, U.** (2001). Molecular secrets of bacterial type III effector proteins. *Trends Plant Sci.* **6**, 479-485.
- Lam, B., and Blumwald, E.** (2002). Domains as functional building blocks of plant proteins. *Trends Plant Sci.* **7**, 544-549.
- Lam, E., Kato, N., and Lawton, M.** (2001). Programmed cell death, mitochondria and the plant hypersensitive response. *Nature* **411**, 848-853.
- Lee, B.M., Park, Y.J., Park, D.S., Kang, H.W., Kim, J.G., Song, E.S., Park, I.C., Yoon, U.H., Hahn, J.H., Koo, B.S., Lee, G.B., Kim, H., Park, H.S., Yoon, K.O., Kim, J.H., Jung, C.H., Koh, N.H., Seo, J.S., and Go, S.J.** (2005). The genome sequence of *Xanthomonas oryzae* pathovar *oryzae* KACC10331, the bacterial blight pathogen of rice. *Nucleic Acids Res* **33**, 577-586.
- Leipe, D.D., Koonin, E.V., and Aravind, L.** (2004). STAND, a class of P-loop NTPases including animal and plant regulators of programmed cell death: multiple, complex domain architectures, unusual phyletic patterns, and evolution by horizontal gene transfer. *J. Mol. Biol.* **343**, 1-28.
- Leister, R.T., Dahlbeck, D., Day, B., Li, Y., Chesnokova, O., and Staskawicz, B.J.** (2005). Molecular genetic evidence for the role of SGT1 in the intramolecular complementation of Bs2 protein activity in *Nicotiana benthamiana*. *Plant Cell* **17**, 1268-1278.
- Levy, M., Edelbaum, O., and Sela, I.** (2004). Tobacco mosaic virus regulates the expression of its own resistance gene *N*. *Plant Physiol.* **135**, 2392-2397.
- Li, P., Long, J.Y., Huang, Y.C., Zhang, Y., and Wang, J.S.** (2004). *AvrXa3*: A novel member of *avrBs3* gene family from *Xanthomonas oryzae* pv. *oryzae* has a dual function. *Progr. Nat. Sci.* **14**, 774-780.
- Liang, B., Yu, T.G., Guo, B., Yang, C., Dai, L.Y., and Shen, D.L.** (2004). Cloning and characterization of a novel avirulence gene (*arp3*) from *Xanthomonas oryzae* pv. *oryzae*. *DNA Seq.* **15**, 110-117.
- Liu, Y., Schiff, M., Serino, G., Deng, X.-W., and Dinesh-Kumar, S.P.** (2002). Role of SCF ubiquitin-ligase and the COP9 signalosome in the *N* gene-mediated resistance response to tobacco mosaic virus. *Plant Cell* **14**, 1483-1496.
- Liu, Y., Burch-Smith, T., Schiff, M., Feng, S., and Dinesh-Kumar, S.P.** (2004). Molecular chaperone Hsp90 associates with resistance protein *N* and its signaling proteins SGT1 and Rar1 to modulate an innate immune response in plants. *J. Biochem.* **279**, 2101-2108.

- Lu, R., Malcuit, I., Moffett, P., Ruiz, M.T., Peart, J., Wu, A.-J., Rathjen, J.P., Bendahmane, A., Day, L., and Baulcombe, D.C.** (2003). High throughput virus-induced gene silencing implicates heat shock protein 90 in plant disease resistance. *EMBO J.* **22**, 5690-5699.
- Luck, J.E., Lawrence, G.J., Dodds, P.N., Shepherd, K.W., and Ellis, J.G.** (2000). Regions outside of the leucine-rich repeats of flax rust resistance proteins play a role in specificity determination. *Plant Cell* **12**, 12367-11377.
- Mackey, D., Holt, B.F., Wiig, A., and Dangl, J.L.** (2002). RIN4 interacts with *Pseudomonas syringae* type III effector molecules and is required for RPM1-mediated resistance in *Arabidopsis*. *Cell* **108**, 743-754.
- Mackey, D., Belkhadir, Y., Alonso, J.M., Ecker, J.R., and Dangl, J.L.** (2003). *Arabidopsis* RIN4 is a target of the type III virulence effector AvrRpt2 and modulates RPS2-mediated resistance. *Cell* **112**, 379-489.
- Mackey, D.M.** (2003). Manipulation of Plant Signaling by Bacterial Effector Proteins.
- Marois, E., Van den Ackerveken, G., and Bonas, U.** (2002). The *Xanthomonas* type III effector protein AvrBs3 modulates plant gene expression and induces cell hypertrophy in the susceptible host. *Mol. Plant-Microbe Interact.* **15**, 637-646.
- Martin, G.B., Brommonschenkel, S.H., Chunwongse, J., Frary, A., Ganai, M.W., Spivey, R., Wu, T.Y., Earle, E.D., and Tanksley, S.D.** (1993). Map-based cloning of a protein-kinase gene conferring disease resistance in tomato. *Science* **262**, 1432-1436.
- McHale, L., Tan, X., Koehl, P., and Michelmore, R.W.** (2006). Plant NBS-LRR proteins: adaptable guards. *Genome Biol.* **7**, 212.
- Mestre, P., and Baulcombe, D.C.** (2006). Elicitor-mediated oligomerization of the tobacco N disease resistance protein. *Plant Cell* **18**, 491-501.
- Meyers, B.C., Shen, K.A., Rohani, P., Gaut, B.S., and Michelmore, R.W.** (1998). Receptor-like genes in the major resistance locus of lettuce are subject to divergent selection. *Plant Cell* **10**, 1833-1846.
- Meyers, B.C., Kozik, A., Griego, A., Kuang, H., and Michelmore, R.W.** (2003). Genome-Wide Analysis of NBS-LRR-Encoding Genes in *Arabidopsis*. *Plant Cell* **15**, 809-834.
- Meyers, B.C., Dickerman, A.W., Michelmore, R.W., Sivaramakrishnan, S., Sobral, B.W., and Young, N.D.** (1999). Plant disease resistance genes encode members of an ancient and diverse protein family within the nucleotide-binding superfamily. *Plant J.* **20**, 317-332.
- Milligan, S.B., Bodeau, J., Yaghoobi, J., Kaloshian, I., Zabel, P., and Williamson, V.M.** (1998). The root knot nematode resistance gene *Mi* from tomato is a member of the leucine zipper, nucleotide binding, leucine-rich repeat family of plant genes. *Plant Cell* **10**, 1307-1319.
- Mindrinos, M., Katagiri, F., Yu, G.L., and Ausubel, F.M.** (1994). The *A. thaliana* disease resistance gene *RPS2* encodes a protein containing a nucleotide-binding site and leucine-rich repeats. *Cell* **78**, 1089-1099.
- Minsavage, G., Jones, J., R., S., Miller, S., and Ritchie, D.** (1999). Hypersensitive resistance in *Capsicum pubescens* PI 235047 to *Xanthomonas campestris* pv. *vesicatoria* (*Xcv*) is elicited by *avrBs3-2*. *Phytopathology* **89**, S53.
- Moffett, P., Farnham, G., Peart, J., and Baulcombe, D.C.** (2002). Interaction between domains of a plant NBS-LRR protein in disease resistance-related cell death. *EMBO J.* **21**, 4511-4519.
- Muskett, P., and Parker, J.** (2003). Role of SGT1 in the regulation of plant R gene signalling. *Microbes Infect.* **5**, 969-976.

- Neuwald, A.F., Aravind, L., Spouge, J.L., and Koonin, E.V. (1999). AAA+: A class of chaperone-like ATPases associated with the assembly, operation, and disassembly of protein complexes. *Genome Res.* **9**, 27-43.
- Nimchuk, Z., Marois, E., Kjemtrup, S., Leister, R.T., Katagiri, F., and Dangl, J.L. (2000). Eukaryotic fatty acylation drives plasma membrane targeting and enhances function of several type III effector proteins from *Pseudomonas syringae*. *Cell* **101**, 353-363.
- Nürnberg, T., Brunner, F., Kemmerling, B., and Piater, L. (2004). Innate immunity in plants and animals: striking similarities and obvious differences. *Immunol. Rev.* **198**, 249-266.
- Oldroyd, G.E.D., and Staskawicz, B.J. (1998). Genetically engineered broad-spectrum disease resistance in tomato. *Proc. Natl. Acad. Sci. USA* **95**, 10300-10305.
- Paal, J., Henselewski, H., Muth, J., Meksem, K., Menendez, C.M., Salamini, F., Ballvora, A., and Gebhardt, C. (2004). Molecular cloning of the potato *Gro1-4* gene conferring resistance to pathotype Ro1 of the root cyst nematode *Globodera rostochiensis*. *Plant J.* **38**, 285-297.
- Pan, Q.L., Wendel, J., and Fluhr, R. (2000). Divergent evolution of plant NBS-LRR resistance gene homologues in dicot and cereal genomes. *J. Mol. Evol.* **50**, 203-213.
- Parker, J.E., Coleman, M.J., Szabo, V., Frost, L.N., Schmidt, R., van der Biezen, E.A., Moores, T., Dean, C., Daniels, M.J., and Jones, J.D.G. (1997). The *Arabidopsis* downy mildew resistance gene *RPP5* shares similarity to the Toll and Interleukin-1 receptors with *N* and *L6*. *Plant Cell* **9**, 879-894.
- Parniske, M., Hammond-Kosack, K.E., Golstein, C., Thomas, C.M., Jones, D.A., Harrison, K., Wulff, B.B.H., and Jones, J.D.G. (1997). Novel disease resistance specificities result from sequence exchange between tandemly repeated genes at the *Cf-4/9* locus of tomato. *Cell* **91**, 821-832.
- Peart, J.R., Lu, R., Sadanandom, A., Malcuit, I., Moffett, P., Brice, D.C., Schauser, L., Jaggard, D.A.W., Xiao, S., Coleman, M.J., Dow, M., Jones, J.D.G., Shirasu, K., and Baulcombe, D.C. (2002). Ubiquitin ligase-associated protein SGT1 is required for host and nonhost disease resistance in plants. *PNAS* **99**, 10865-10869.
- Pierre, M., Noël, L., Lahaye, T., Ballvora, A., Veuskens, J., Galal, M., and Bonas, U. (2000). High-resolution genetic mapping of the pepper resistance locus *Bs3* governing recognition of the *Xanthomonas campestris* pv. *vesicatora* AvrBs3 protein. *Theor. Appl. Genet.* **101**, 255-263.
- Pink, D.A.C. (2002). Strategies using genes for non-durable disease resistance. *Euphytica* **124**, 227-236.
- Porter, B.W., Chittoor, J.M., Yano, M., Sasaki, T., and White, F.F. (2003). Development and mapping of markers linked to the rice bacterial blight Resistance gene *Xa7*. *Crop Sci.* **43**, 1484-1492.
- Puri, N., Jenner, C., Bennett, M., Stewart, R., Mansfield, J., Lyons, N., and Taylor, J. (1997). Expression of *avrPphB*, an avirulence gene from *Pseudomonas syringae* pv. *phaseolicola*, and the delivery of signals causing the hypersensitive reaction in bean. *Mol. Plant-Microbe Interact.* **10**, 247-256.
- Rathjen, J., and Moffett, P. (2003). Early signal transduction events in specific plant disease resistance. *Curr. Opin. Plant Biol.* **6**, 300-306.
- Riedl, S.J., Li, W., Chao, Y., Schwarzenbacher, R., and Shi, Y. (2005). Structure of the apoptotic protease-activating factor 1 bound to ADP. *Nature* **434**, 926-933.
- Ronald, P.C., Salmeron, J.M., Carland, F.M., and Staskawicz, B.J. (1992). The cloned avirulence gene *avrPto* induces disease resistance in tomato cultivars containing the *Pto* resistance gene. *J. Bacteriol.* **174**, 1604-1611.

- Rooney, H.C., Van't Klooster, J.W., van der Hoorn, R.A., Joosten, M.H., Jones, J.D., and de Wit, P.J. (2005). Cladosporium Avr2 inhibits tomato Rcr3 protease required for Cf-2-dependent disease resistance. *Science* **308**, 1783-1786.
- Salanoubat, M., Genin, S., Artiguenave, F., Gouzy, J., Mangenot, S., Arlat, M., Billault, A., Brottier, P., Camus, J.C., Cattolico, L., Chandler, M., Choisine, N., Claudel-Renard, C., Cunnac, S., Demange, N., Gaspin, C., Lavie, M., Moisan, A., Robert, C., Saurin, W., Schiex, T., Siguier, P., Thebault, P., Whalen, M., Wincker, P., Levy, M., Weissenbach, J., and Boucher, C.A. (2002). Genome sequence of the plant pathogen *Ralstonia solanacearum*. *Nature* **415**, 497-502.
- Salmeron, J.M., Oldroyd, G.E.D., Rommens, C.M.T., Scofield, S.R., Kim, H.S., Lavelle, D.T., Dahlbeck, D., and Staskawicz, B.J. (1996). Tomato *Prf* is a member of the leucine-rich repeat class of plant disease resistance genes and lies embedded within the *Pto* kinase gene-cluster. *Cell* **86**, 123-133.
- Schlögelhofer, P., Garzon, M., Kerzendorfer, C., Nizhynska, V., and Bachmair, A. (2006). Expression of the ubiquitin variant ubR48 decreases proteolytic activity in Arabidopsis and induces cell death. *Planta* **223**, 684-697.
- Schornack, S., Peter, K., Bonas, U., and Lahaye, T. (2005). Expression levels of *avrBs3*-like genes affect recognition specificity in tomato *Bs4*- but not in pepper *Bs3*-mediated perception. *Mol. Plant-Microbe Interact.* **18**, 1215-1225.
- Schornack, S., Meyer, A., Römer, P., Jordan, T., and Lahaye, T. (2006). Gene-for-gene-mediated recognition of nuclear-targeted AvrBs3-like bacterial effector proteins. *J. Plant Physiol.* **163**, 256-272.
- Schornack, S., Ballvora, A., Gürlebeck, D., Peart, J., Baulcombe, D., Baker, B., Ganai, M., Bonas, U., and Lahaye, T. (2004). The tomato resistance protein Bs4 is a predicted non-nuclear TIR-NB-LRR protein that mediates defense responses to severely truncated derivatives of AvrBs4 and overexpressed AvrBs3. *Plant J.* **37**, 46-60.
- Schulze-Lefert, P. (2004). Plant immunity: the origami of receptor activation. *Curr. Biol.* **14**, R22-24.
- Shao, F., Golstein, C., Ade, J., Stoutemyer, M., Dixon, J.E., and Innes, R.W. (2003). Cleavage of *Arabidopsis* PBS1 by a bacterial type III effector. *Science* **301**, 1230-1233.
- Shen, K.A., Chin, D.B., Arroyo-Garcia, R., Ochoa, O.E., Lavelle, D.O., Wroblewski, T., Meyers, B.C., and Micheltore, R.W. (2002). *Dm3* is one member of a large constitutively expressed family of nucleotide binding site-leucine-rich repeat encoding genes. *Mol. Plant-Microbe Interact.* **15**, 251-261.
- Shirasu, K., and Schulze-Lefert, P. (2003). Complex formation, promiscuity and multi-functionality: protein interactions in disease-resistance pathways. *Trends Plant Sci.* **8**, 252-258.
- Sidhu, G.S., Khush, G.S., and Mew, T.W. (1978). Genetic-analysis of bacterial-blight resistance in 74 cultivars of rice, *Oryza-sativa*-L. *Theor. Appl. Genet.* **53**, 105-111.
- Sobczak, M., Avrova, A., Jupowicz, J., Phillips, M.S., Ernst, K., and Kumar, A. (2005). Characterization of susceptibility and resistance responses to potato cyst nematode (*Globodera* spp.) infection of tomato lines in the absence and presence of the broad-spectrum nematode resistance *Hero* gene. *Mol. Plant-Microbe Interact.* **18**, 158-168.
- Staskawicz, B.J., Dahlbeck, D., and Keen, N.T. (1984). Cloned avirulence gene of *Pseudomonas syringae* pv. *glycinea* determines race specific incompatibility on *Glycine max.* (L.) Merr. *Proc. Natl. Acad. Sci. USA* **81**, 6024-6028.

- Stokes, T.L., and Richards, E.J.** (2002). Induced instability of two *Arabidopsis* constitutive pathogen-response alleles. *Proc. Natl. Acad. Sci. USA* **99**, 7792-7796.
- Stokes, T.L., Kunkel, B.N., and Richards, E.J.** (2002). Epigenetic variation in *Arabidopsis* disease resistance. *Genes Dev.* **16**, 171-182.
- Swiderski, M.R., and Innes, R.W.** (2001). The *Arabidopsis* *PBS1* resistance gene encodes a member of a novel protein kinase subfamily. *Plant J.* **26**, 101-112.
- Szurek, B., Marois, E., Bonas, U., and Van den Ackerveken, G.** (2001). Eukaryotic features of the *Xanthomonas* type III effector AvrBs3: protein domains involved in transcriptional activation and the interaction with nuclear import receptors from pepper. *Plant J.* **26**, 523-534.
- Szurek, B., Rossier, O., Hause, G., and Bonas, U.** (2002). Type III-dependent translocation of the *Xanthomonas* AvrBs3 protein into the plant cell. *Mol. Microbiol.* **46**, 13-23.
- Taddei, A., Hediger, F., Neumann, F.R., and Gasser, S.M.** (2004). The function of nuclear architecture: a genetic approach. *Annu. Rev. Genet.* **38**, 305-345.
- Takken, F.L., Albrecht, M., and Tameling, W.I.** (2006). Resistance proteins: molecular switches of plant defence. *Curr. Opin. Plant Biol.*
- Tameling, W.I.L., Elzinga, S.D.J., Darmin, P.S., Vossen, J.H., Takken, F.L.W., Haring, M.A., and Cornelissen, B.J.C.** (2002). The tomato *R* gene products I-2 and Mi-1 are functional ATP binding proteins with ATPase activity. *Plant Cell* **14**, 2929-2939.
- Tao, Y., Yuan, F., Leister, R.T., Ausubel, F.M., and Katagiri, F.** (2000). Mutational analysis of the *Arabidopsis* nucleotide binding site-leucine-rich repeat resistance gene *RPS2*. *Plant Cell* **12**, 2541-2554.
- Thompson, G.A., Jr., and Okuyama, H.** (2000). Lipid-linked proteins of plants. *Prog. Lipid Res.* **39**, 19-39.
- Thrower, J.S., Hoffman, L., Rechsteiner, M., and Pickart, C.M.** (2000). Recognition of the polyubiquitin proteolytic signal. *EMBO J.* **19**, 94-102.
- Tian, D., Traw, M.B., Chen, J.Q., Kreitman, M., and Bergelson, J.** (2003). Fitness costs of R-gene-mediated resistance in *Arabidopsis thaliana*. *Nature* **423**, 74-77.
- Tör, M., Gordon, P., Cuzick, A., Eulgem, T., Sinapidou, E., Mert-Turk, F., Can, C., Dangl, J.L., and Holub, E.B.** (2002). *Arabidopsis* SGT1b is required for defense signaling conferred by several downy mildew resistance genes. *Plant Cell* **14**, 993-1003.
- Van den Ackerveken, G., Marois, E., and Bonas, U.** (1996). Recognition of the bacterial avirulence protein AvrBs3 occurs inside the host plant cell. *Cell* **87**, 1307-1316.
- Van der Biezen, E.A., and Jones, J.D.G.** (1998a). The NB-ARC domain: a novel signalling motif shared by plant resistance gene products and regulators of cell death in animals. *Curr. Biol.* **8**, R226-R227.
- Van der Biezen, E.A., and Jones, J.D.G.** (1998b). Plant disease-resistance proteins and the gene-for-gene concept. *Trends Biochem. Sci.* **23**, 454-456.
- Van der Hoorn, R.A.L., Laurent, F., Roth, R., and De Wit, P.J.G.M.** (2000). Agroinfiltration is a versatile tool that facilitates comparative analyses of *Avr9/Cf-9*-induced and *Avr4/Cf-4*-induced necrosis. *Mol. Plant-Microbe Interact.* **13**, 439-446.
- Van der Vossen, E.A.G., Rouppe van der Voort, J.N.A.M., Kanyuka, K., Bendahmane, A., Sandbrink, H., Baulcombe, D.C., Bakker, J., Stiekema, W.J., and Klein-Lankhorst, R.M.** (2000). Homologues of a single resistance-gene cluster in potato confer resistance to distinct pathogens: a virus and a nematode. *Plant J.* **23**, 567-576.

- van Doorn, W.G., and Woltering, E.J.** (2005). Many ways to exit? Cell death categories in plants. *Trends Plant Sci.* **10**, 117-122.
- Vera Cruz, C.M., Bai, J., Ona, I., Leung, H., Nelson, R.J., Mew, T.W., and Leach, J.E.** (2000). Predicting durability of a disease resistance gene based on an assessment of the fitness loss and epidemiological consequences of avirulence gene mutation. *Proc. Natl. Acad. Sci. USA* **97**, 13500-13505.
- Vidal, S., Cabrera, H., Andersson, R.A., Fredriksson, A., and Valkonen, J.P.T.** (2002). Potato gene *Y-1* is an *N* gene homolog that confers cell death upon infection with *potato virus Y*. *Mol. Plant-Microbe Interact.* **15**, 717-727.
- Vos, P., Simons, G., Jesse, T., Wijbrandi, J., Heinen, L., Hogers, R., Frijters, A., Groenendijk, J., Diergaarde, P., Reijans, M., Fierens-Onstenk, J., de Both, M., Peleman, J., Liharska, T., Hontelez, J., and Zabeau, M.** (1998). The tomato *Mi-1* gene confers resistance to both root-knot nematode and potato aphids. *Nature Biotechnol.* **16**, 1365-1369.
- Warren, R.F., Henk, A., Mowery, P., Holub, E., and Innes, R.W.** (1998). A mutation within the leucine-rich repeat domain of the Arabidopsis disease resistance gene *RPS5* partially suppresses multiple bacterial and downy mildew resistance genes. *Plant Cell* **10**, 1439-1452.
- Wendehenne, D., Durner, J., and Klessig, D.** (2004). Nitric oxide: a new player in plant signalling and defence responses. *Curr. Opin. Plant Biol.* **7**, 449-455
- White, F.F., Yang, B., and Johnson, L.B.** (2000). Prospects for understanding avirulence gene function. *Curr. Opin. Plant Biol.* **3**, 291-298.
- Whitham, S., McCormick, S., and Baker, B.** (1996). The *N* gene of tobacco confers resistance to tobacco mosaic virus in transgenic tomato. *Proc. Natl. Acad. Sci. USA* **93**, 8776-8781.
- Whitham, S., Dinesh-Kumar, S.P., Choi, D., Hehl, R., Corr, C., and Baker, B.** (1994). The product of the tobacco mosaic virus resistance gene *N*: similarity to Toll and the Interleukin-1 receptor. *Cell* **78**, 1101-1115.
- Wichmann, G., and Bergelson, J.** (2004). Effector genes of *Xanthomonas axonopodis* pv. *vesicatoria* promote transmission and enhance other fitness traits in the field. *Genetics* **166**, 693-706.
- Xinghua, L., Duanpin, Z., Yuefeng, X., Qifa, Z., and Heping, G.** (1996). Mapping a new gene for resistance to bacterial blight using RFLP markers. *Internatl. Rice Res. Notes* **21**, 30.
- Xu, Y., Tao, X., Shen, B., Horng, T., Medzhitov, R., Manley, J.L., and Tong, L.** (2000). Structural basis for signal transduction by the Toll/interleukin-1 receptor domains. *Nature* **408**, 111-115.
- Yang, B., and White, F.F.** (2004). Diverse members of the AvrBs3/PthA family of type III effectors are major virulence determinants in bacterial blight disease of rice. *Mol. Plant-Microbe Interact.* **17**, 1192-1200.
- Yang, B., Sugio, A., and White, F.F.** (2005). Avoidance of host recognition by alterations in the repetitive and C-terminal regions of AvrXa7, a type III effector of *Xanthomonas oryzae* pv. *oryzae*. *Mol. Plant-Microbe Interact.* **18**, 142-149.
- Yang, B., Sugio, A., and White, F.F.** (2006). *Os8N3* is a host disease-susceptibility gene for bacterial blight of rice. *Proc. Natl. Acad. Sci. USA* **103**, 10503-10508.
- Yang, B., Zhu, W., Johnson, L.B., and White, F.F.** (2000). The virulence factor AvrXa7 of *Xanthomonas oryzae* pv. *oryzae* is a type III secretion pathway-dependent nuclear-localized double-stranded DNA-binding protein. *Proc. Natl. Acad. Sci. USA* **97**, 9807-9812.

- Yang, Y.N., and Gabriel, D.W.** (1995). *Xanthomonas* avirulence/pathogenicity gene family encodes functional plant nuclear targeting signals. *Mol. Plant-Microbe Interact.* **8**, 627-631.
- Yang, Y.N., De Feyter, R., and Gabriel, D.W.** (1994). Host-specific symptoms and increased release of *Xanthomonas citri* and *X. campestris* pv. *malvacearum* from leaves are determined by the 102-bp tandem repeats of *pthA* and *avrb6*, respectively. *Mol. Plant-Microbe Interact.* **7**, 345-355.
- Yoshimura, A., Mew, T.W., Khush, G.S., and Omura, T.** (1983). Inheritance of resistance to bacterial blight in rice cultivar Cas209. *Phytopathology* **73**, 1409-1412.
- Yoshimura, S., Yoshimura, A., Iwata, N., McCouch, S.R., Abenes, M.L., Baraoidan, M.R., Mew, T.W., and Nelson, R.J.** (1995). Tagging and combining bacterial-blight resistance genes in rice using RAPD and RFLP markers. *Mol. Breeding* **1**, 375-387.
- Yoshimura, S., Yamanouchi, U., Katayose, Y., Toki, S., Wang, Z.X., Kono, I., Kurata, N., Yano, M., Iwata, N., and Sasaki, T.** (1998). Expression of *Xa1*, a bacterial blight-resistance gene in rice, is induced by bacterial inoculation. *Proc. Natl. Acad. Sci. USA* **95**, 1663-1668.
- Yoshimura, S., A., Yoshimura, A., Saito, N., Kishimoto, M., Kawase, M., Yano, M., Nakagahra, M., Ogawa, T., and Iwata, N.** (1992). RFLP analysis of introgressed chromosomal segments in three near-isogenic lines of rice for bacterial blight resistance genes, *Xa-1*, *Xa-3*, *Xa-4*. *Jpn. J. Genet.* **67**, 29-37.
- Young, N.D.** (2000). The genetic architecture of resistance. *Curr. Opin. Plant Biol.* **3**, 285-290.
- Zhang, Y., Dorey, S., Swiderski, M., and Jones, J.D.G.** (2004). Expression of *RPS4* in tobacco induces an AvrRps4-independent HR that requires EDS1, SGT1 and HSP90. *Plant J.* **40**, 213-224.
- Zhong, Y.M., Jiang, G.H., Chen, X.W., Xia, Z.H., Li, X.B., Zhu, L.H., and Zhai, W.X.** (2003). Identification and gene prediction of a 24 kb region containing *xa5*, a recessive bacterial blight resistance gene in rice (*Oryza sativa* L.). *Chin. Sci. Bull.* **48**, 2725-2729.
- Zhu, W.G., Yang, B., Chittoor, J.M., Johnson, L.B., and White, F.F.** (1998). AvrXa10 contains an acidic transcriptional activation domain in the functionally conserved C terminus. *Mol. Plant-Microbe Interact.* **11**, 824-832.
- Zipfel, C., Robatzek, S., Navarro, L., Oakeley, E.J., Jones, J.D., Felix, G., and Boller, T.** (2004). Bacterial disease resistance in Arabidopsis through flagellin perception. *Nature* **428**, 764-767.