

9. Literaturverzeichnis

1. Ainamo J, Barmes D, Beagrie G, Cutress T, Martin J, Sardo-Infirri J: Development of the World Health Organization (WHO) community periodontal index of treatment needs (CPITN). *Int Dent J* 32 (1982) 281-291
2. American Academy of Pediatric Dentistry: Treatment of temporomandibular disorders in children: summary statements and recommendations. *J Am Dent Assoc* 120 (1990) 265, 267, 269
3. American Academy of Orofacial Pain: Orofacial Pain, Guidelines for Assessment, Diagnosis and Management. Quintessence, Chicago, 1996
4. Bader G, Lavigne G: Sleep bruxism; an overview of an oromandibular sleep movement disorder. Review article. *Sleep Med Rev* 4 (2000) 27-43
5. Bartlett DW, Coward PY, Nikkah C, Wilson RF: The prevalence of tooth wear in a cluster sample of adolescent schoolchildren and its relationship with potential explanatory factors. *Br Dent J* 184 (1998) 125-129
6. Bernal M, Tsamtsouris A: Signs and symptoms of temporomandibular joint dysfunction in 3 to 5 year old children. *J Pedod* 10 (1986) 127-140
7. Bragdon EE, Light KC, Costello NL, Sigurdsson A, Bunting S, Bhalang K, Maixner W: Group differences in pain modulation: pain-free women compared to pain-free men and to women with TMD. *Pain* 96 (2002) 227-237
8. Bush FM, Harkins SW, Harrington WG, Price DD: Analysis of gender effects on pain perception and symptom presentation in temporomandibular pain. *Pain* 53 (1993) 73-80
9. Carlson CR, Bertrand PM, Ehrlich AD, Maxwell AW, Burton RG: Physical self-regulation training for the management of temporomandibular disorders. *J Orofac Pain* 15 (2001) 47-55
10. Carlsson GE, Egermark I, Magnusson T: Predictors of bruxism, other oral parafunctions, and tooth wear over a 20-year follow-up period. *J Orofac Pain* 17 (2003) 50-57
11. Chuang SY: Incidence of temporomandibular disorders (TMDs) in senior dental students in Taiwan. *J Oral Rehabil* 29 (2002) 1206-1211
12. Cochran W: *Sampling Techniques*. Wiley, New York, 1977
13. Crombez G, Baeyens F, de Laat A, Goosens L: Bruxisme: een literatuuroverzicht. *Gedragstherapie* jrg. 24, nr. 3 (1991) 157-172

14. De Boever JA, Carlson GE, Klineberg IJ: Need for occlusal therapy and prosthodontic treatment in the management of temporomandibular disorders. Part II. Tooth loss and prosthodontic treatment. *J Oral Rehabil* 27 (2000) 647-659
15. De Boever JA, van den Berghe L: Longitudinal study of functional conditions in the masticatory system in Flemish children. *Community Dent Oral Epidemiol* 15 (1987) 100-103
16. De Vis H, De Boever JA, van Cauwenberghe P: Epidemiologic survey of functional conditions of the masticatory system in Belgian children aged 3-6 years. *Community Dent Oral Epidemiol* 12 (1984) 203-207
17. Drangsholt M, LeResche L: Temporomandibular disorder pain. IASP Press, Seattle, 1999
18. Droukas B, Lindee C, Carlsson GE: Relationship between occlusal factors and signs and symptoms of mandibular dysfunction. A clinical study of 48 dental students. *Acta Odontol Scand* 42 (1984) 277-283
19. Dworkin SF, LeResche L: Research diagnostic criteria for temporomandibular disorders: review, criteria, examinations and specifications, critique. *J Craniomandib Disord* 6 (1992) 301-355
20. Egermark I, Carlsson GE, Magnusson T: A 20-year longitudinal study of subjective symptoms of temporomandibular disorders from childhood to adulthood. *Acta Odontol Scand* 59 (2001) 40-48
21. Egermark-Eriksson I, Carlsson GE, Ingervall B: Prevalence of mandibular dysfunction and orofacial parafunction in 7-, 11- and 15-year-old Swedish children. *Eur J Orthod* 3 (1981) 163-172
22. Fillingim RB, Maixner W: Sex-related factors in temporomandibular disorders. IASP Press, Seattle, 2000
23. Fletcher RH, Fletcher SW, Wagner EH: *Klinische Epidemiologie*. Ullstein Medical Verlagsgesellschaft mbH & Co., Wiesbaden, 1999
24. Goho C, Jones HL: Association between primary dentition wear and clinical temporomandibular dysfunction signs. *Pediatr Dent* 13 (1991) 263-266
25. Goulet JP, Clark GT, Flack VF, Liu C: The reproducibility of muscle and joint tenderness detection methods and maximum mandibular movement measurement for the temporomandibular system. *J Orofac Pain* 12 (1998) 17-26

26. Goulet JP, Lavigne GJL und JP: Jaw pain prevalence among French-speaking Canadians in Quebec and related symptoms of temporomandibular disorders. *J Dent Res* 74 (1995) 1738-1744
27. Hanamura H, Houston F, Rylander H, Carlsson GE, Haraldson T, Nyman S: Periodontal status and bruxism. A comparative study of patients with periodontal disease and occlusal parafunctions. *J Periodontol* 58 (1987) 173-176
28. Helkimo M: Studies on function and dysfunction of the masticatory system. II. Index for anamnestic and clinical dysfunction and occlusal state. *Sven Tandlak Tidskr* 67 (1974) 101-121
29. Hirsch C: Kraniomandibuläre Dysfunktionen (CMD) bei Kindern und Jugendlichen - Prävalenz, Beeinträchtigungen und Einflüsse der physischen Entwicklung. Habilitationsschrift, MLU Halle-Wittenberg, Halle/Saale, 2003a
30. Hirsch C, John MT: Prävalenz kraniomandibulärer Dysfunktionen (CMD) bei Kindern und Jugendlichen. *DZZ* (58) 10 (2003b) 589-592
31. Hirsch C, John MT: Szenarien zu Krankheitskosten in der Diagnostik und Therapie kraniomandibulärer Dysfunktionen in Deutschland. *DZZ* (60) 2 (2005) 108-111
32. Huang GJ, LeResche L, Critchlow CW, Martin MD, Drangsholt MT: Risk factors for diagnostic subgroups of painful temporomandibular disorders (TMD). *J Dent Res* 81 (2002) 284-288
33. Hublin C, Kaprio J, Partinen M, Koskenvuo M: Sleep bruxism based on self-report in a nationwide twin cohort. *J Sleep Res* 7 (1998) 61-67
34. Johansson A, Haraldson T, Omar R, Kiliaridis S, Carlsson GE: A system for assessing the severity and progression of occlusal tooth wear. *J Oral Rehabil* 20 (1993) 125-131
35. John M, Hirsch C, Reiber T: Häufigkeit, Bedeutung und Behandlungsbedarf kraniomandibulärer Dysfunktionen. *Zeitschr f Gesundheitswissenschaften* 9 (2001) 136-155
36. John M, Wefers K-P: Orale Dysfunktionen bei den Erwachsenen. Deutscher Ärzteverlag, Köln, 1999
37. John MT, Frank H, Lobbezoo F, Drangsholt M, Dette KE. No association between incisal tooth wear and temporomandibular disorders. *J Prosthet Dent* 87 (2002a) 197-203

38. John MT, Hirsch C, Reiber T, Dworkin SF: Translating the Research Diagnostic Criteria for Temporomandibular Disorders into German: Evaluation of content and process. *J Orofac Pain* (im Druck) (2006)
39. John MT, Hirsch C, Drangsholt MT, Mancl LA, Setz JM: Overbite and overjet are not related to self-report of temporomandibular disorder symptoms. *J Dent Res* 81 (2002b) 164-169
40. Kieser JA, Groeneveld HT: Relationship between juvenile bruxing and craniomandibular dysfunction. *J Oral Rehabil* 25 (1998) 662-665
41. Kononen M, Waltimo A, Nystrom M: Does clicking in adolescence lead to painful temporomandibular joint locking? *Lancet* 347 (1996) 1080-1081
42. Korkhaus G: Prophylaxe und Frühbehandlung der Zahnstellungs- und Kieferanomalien. Hermann Meuser, Leipzig & Berlin, 1932
43. Koutantji M, Pearce SA, Oakley DA: The relationship between gender and family history of pain with current pain experience and awareness of pain in others. *Pain* 77 (1998) 25-31
44. Kupfer J, Felder H, Brähler E: Zur Genese geschlechtsspezifischer Somatisierung. Westdt. Verlag, Wiesbaden, 1999
45. Lavigne G, Manzini G: *Bruxism*. W.B. Saunders Co, Philadelphia, 2000
46. Lavigne G, Montplaisir JY: *Bruxism: epidemiology* In: J. R. Fricton and R. Dubner (Hrsg): *Orofacial Pain and Temporomandibular Disorders*. Raven Press, New York, 1995
47. Leroy R, Bogaerts K, Lesaffre E, Declerck D: The emergence of permanent teeth in Flemish children. *Community Dent Oral Epidemiol* 31 (2003) 30-39
48. List T, Wahlund K, Wenneberg B, Dworkin SF: TMD in children and adolescents: prevalence of pain, gender differences, and perceived treatment need. *J Orofac Pain* 13 (1999) 9-20
49. Lobbezoo F, Lavigne GJ: Do bruxism and temporomandibular disorders have a cause-and-effect relationship? *J Orofac Pain* 11 (1997) 15-23
50. Lobbezoo F, Naeije M: Bruxism is mainly regulated centrally, not peripherally. *J Oral Rehabil* 28 (2001) 1085-1091
51. Lobbezoo F, Rompre PH, Soucy JP, Iafrancesco C, Turkewicz J, Montplaisir JY, Lavigne, GJ: Lack of associations between occlusal and cephalometric measures, side

- imbalance in striatal D2 receptor binding, and sleep-related oromotor activities. *J Orofac Pain* 15 (2001) 64-71
52. Locker D, Slade G: Prevalence of symptoms associated with temporomandibular disorders in a Canadian population. *Community Dent Oral Epidemiol* 16 (1988) 310-313
53. Magnusson T, Carlsson GE, Egermark I: Changes in subjective symptoms of craniomandibular disorders in children and adolescents during a 10-year period. *J Orofac Pain* 7 (1993) 76-82
54. Magnusson T, Egermark I, Carlsson GE: A longitudinal epidemiologic study of signs and symptoms of temporomandibular disorders from 15 to 35 years of age. *J Orofac Pain* 14 (2000) 310-319
55. Marbach JJ, Lennon MC, Dohrenwend BP: Candidate risk factors for temporomandibular pain and dysfunction syndrome: psychosocial, health behavior, physical illness and injury. *Pain* 34 (1988) 139-151
56. Marbach JJ, Raphael KG, Dohrenwend BP, Lennon MC: The validity of tooth grinding measures: etiology of pain dysfunction syndrome revisited. *J Am Dent Assoc* 120 (1990) 327-333
57. Meltzer JW, Hochstim JR: Reliability and validity of survey data on physical health. *Public Health Rep* 85 (1970) 1075-1086
58. Meng HP, Dibbets JM, van der Weele LT, Boering G: Symptoms of temporomandibular joint dysfunction and predisposing factors. *J Prosthet Dent* 57 (1987) 215-222
59. Micheelis W, Reich E: Dritte Deutsche Mundgesundheitsstudie - DMS III. Deutscher Ärzte Verlag, Köln, 1999
60. Mohl ND, Dixon DC: Current status of diagnostic procedures for temporomandibular disorders. *J Am Dent Assoc* 125 (1994) 56-64
61. Molina OF, dos Santos Junior J, Nelson SJ, Nowlin T: Profile of TMD and Bruxer compared to TMD and nonbruxer patients regarding chief complaint, previous consultations, modes of therapy, and chronicity. *Cranio* 18 (2000) 205-219
62. Nilner M: Relationships between oral parafunctions and functional disturbances and diseases of the stomatognathic system among children aged 7-14 years. *Acta Odontol Scand* 41 (1983a) 167-172

63. Nilner M: Relationships between oral parafunctions and functional disturbances in the stomatognathic system among 15- to 18-year-olds. *Acta Odontol Scand* 41 (1983b) 197-201
64. Nydell A, Helkimo M, Koch G: Craniomandibular disorders in children--a critical review of the literature. *Swed Dent J* 18 (1994) 191-205
65. Okeson JP, O'Donnell JP: Standards for temporomandibular evaluation in the pediatric patient. *Pediatr Dent* 11 (1989) 329-330
66. Principato JJ, Barwell DR: Biofeedback training and relaxation exercises for treatment of temporomandibular joint dysfunction. *Otolaryngology* 86 (1978) 766-769
67. Pullinger AG, Seligman DA: The degree to which attrition characterizes differentiated patient groups of temporomandibular disorders. *J Orofac Pain* 7 (1993) 196-208
68. Round JM, Jones DA, Honour JW, Nevill AM: Hormonal factors in the development of differences in strength between boys and girls during adolescence: a longitudinal study. *Ann Hum Biol* 26 (1999) 49-62
69. Rugh JD, Solberg WK: Electromyographic studies of bruxist behavior before and during treatment. *J Calif Dent Assoc* 3 (1975) 56-59
70. Sari S, Sonmez H: Investigation of the relationship between oral parafunctions and temporomandibular joint dysfunction in Turkish children with mixed and permanent dentition. *J Oral Rehabil* 29 (2002) 108-112
71. Schmitter M, Ohlmann B, John MT, Hirsch C, Rammelsberg P: Research Diagnostic Criteria for Temporomandibular Disorders: A Calibration and Reliability Study. *Cranio* 23 (2005) 212-218
72. Seligman DA, Pullinger AG, Solberg WK: The prevalence of dental attrition and its association with factors of age, gender, occlusion, and TMJ symptomatology. *J Dent Res* 67 (1988) 1323-1333
73. Shi CS, Wang HY: Influence of an occlusal splint on integrated electromyography of the masseter muscles. *J Oral Rehabil* 18 (1991) 253-256
74. Skeppar J, Nilner M: Treatment of craniomandibular disorders in children and young adults. *J Orofac Pain* 7 (1993) 362-369
75. Thorpy MJ: Classification of sleep disorders. *J Clin Neurophysiol* 7 (1990) 67-81
76. Treacy K: Awareness/relaxation training and transcutaneous electrical neural stimulation in the treatment of bruxism. *J Oral Rehabil* 26 (1999) 280-287

77. Tsolka P, Walter JD, Wilson RF, Preiskel HW: Occlusal variables, bruxism and temporomandibular disorders: a clinical and kinesiographic assessment. *J Oral Rehabil* 22 (1995) 849-856
78. Türp JC, John M, Nilges P, Jürgens J: Recommendations for the standardized evaluation and classification of patients with painful temporomandibular disorders. *Schmerz* 14 (2000) 416-428
79. Van der Weele LT, Dibbets JM: Helkimo's index: a scale or just a set of symptoms? *J Oral Rehabil* 14 (1987) 229-237
80. Vanderas AP: Relationship between oral parafunctions and craniomandibular dysfunction in children and adolescents: a review. *ASDC J Dent Child* 61 (1994) 378-381
81. Vanderas AP: Relationship between craniomandibular dysfunction and oral parafunctions in Caucasian children with and without unpleasant life events. *J Oral Rehabil* 22 (1995) 289-294
82. Vanderas AP, Papagiannoulis L: Multifactorial analysis of the aetiology of craniomandibular dysfunction in children. *Int J Paediatr Dent* 12 (2002) 336-346
83. Wahlund K, List T, Dworkin SF: Temporomandibular disorders in children and adolescents: reliability of a questionnaire, clinical examination, and diagnosis. *J Orofac Pain* 12 (1998) 42-51
84. Wanman A, Agerberg G: Mandibular dysfunction in adolescents. II. Prevalence of signs. *Acta Odontol Scand* 44 (1986a) 55-62
85. Wanman A, Agerberg G: Two-year longitudinal study of symptoms of mandibular dysfunction in adolescents. *Acta Odontol Scand* 44 (1986b) 321-331
86. Warren MP, Fried JL: Temporomandibular disorders and hormones in women. *Cells Tissues Organs* 169 (2001) 187-192
87. Wedel A, Carlsson GE: Retrospective review of 350 patients referred to a TMJ clinic. *Community Dent Oral Epidemiol* 11 (1983) 69-73
88. Westling L, Mattiasson A: Background factors in craniomandibular disorders: reported symptoms in adolescents with special reference to joint hypermobility and oral parafunctions. *Scand J Dent Res* 99 (1991) 48-54
89. WHO (World Health Organization): Basic methods. WHO, Genf, 1997
90. Widmalm SE, Christiansen RL, Gunn SM: Oral parafunctions as temporomandibular disorder risk factors in children. *Cranio* 13 (1995a) 242-246

91. Widmalm SE, Gunn SM, Christiansen RL, Hawley LM: Association between CMD signs and symptoms, oral parafunctions, race and sex, in 4-6-year-old African-American and Caucasian children. *J Oral Rehabil* 22 (1995b) 95-100
92. Yap AU: Effects of stabilization appliances on nocturnal parafunctional activities in patients with and without signs of temporomandibular disorders. *J Oral Rehabil* 25 (1998) 64-68
93. Zwijnenburg A, John M, Reiber T: Schmerz als bestimmender Faktor für den subjektiven Behandlungsbedarf kranio-mandibulärer Dysfunktionen. Schlüterscher Verlag, Hannover, 2002