

References:

- Acton S., Rigotti A., Landschulz K.T., Xu S., Hobbs H.H., & Krieger M. (1996) Identification of scavenger receptor SR-BI as a high density lipoprotein receptor. *Science* 271: 518-520.
- Adams M.R, Golden D.L., Anthony M.S, Register T.C., Williams J.K (2002) The inhibitory effect of soy protein isolate on atherosclerosis in mice does not require the presence of LDL receptors or alteration of plasma lipoproteins. *J. Nutr.* 132: 43-49.
- Adlercreutz H. (1990). Western diets and western diseases: Some hormonal and biochemical mechanisms and associations. *Scand. J. Lab. Clin. Invest.* 50 (Suppl. 201): 3-23.
- Ait Yahita D., Madani S., Prost E., Bouchenak M. & Belleville J., (2004) Fish protein improves blood pressure but alters HDL2 and HDL 3 composition and tissue lipoprotein lipase activities in spontaneously hypertensive rats. *Eur. J. Nutr.* 44, 10-17.
- Ali A.A., Velasquez M.T., Hansen C.T., Mohamed A.I., Bhathena S.J., (2004) Effects of soybean isoflavones, probiotics, and their interactions on lipid metabolism and endocrine system in an animal model of obesity and diabetes. *J. Nutr. Biochem.* 15 583-590.
- Anderson J.W., Johnstone B. M., & Cook- Newwell M. E. (1995) Meta analysis of the effects of soy protein intake on serum lipids. *N. Engl. J. Med.* 333: 276-282.
- Ascencio C, Torres N, Isoard-Acosta F, Gómez-Pérez FJ, Hernández-Pando R, Tovar AR. (2004) Soy protein affects serum insulin and hepatic SREBP-1 mRNA and reduces fatty liver in rats. *J Nutr* 134: 522-529.
- Austin M.A. (1991) Plasma triglyceride and coronary heart disease. *Arterioscler. Thromb.* 11: 2-14.
- Austin M.A., Hokanson J.E., and Edwards K.L. (1998) Hypertriglyceridemia as a cardiovascular risk factor. *Am. J. Cardiol.* 81(4A): 7B-12B.
- Antony M.S., Clarkson T.B. & Williams J.K. (1998) Effects of soy isoflavones on atherosclerosis: potential mechanisms. *Am. J. Clin. Nutr.* 68: 1390S-1393S.

- Anthony M.S., Clarkson T.B., B.C., Wagner J.D. (1997) Soy protein versus soy phytoestrogens in the prevention of diet induced coronary artery atherosclerosis of male cynomolgous monkeys. *Arterioscler. Thromb. Vasc. Biol.* 17: 2524-2531.
- Bakhit R.M., Klein B.P., Essex-Sorlie D., Ham J.O., Erdman J.W., and Potter S.M. (1994) Intake of 25 g soybean protein reduces plasma cholesterol in men with elevated cholesterol concentrations. *J. Nutr.* 124: 213-222.
- Balmir F., stack R., Jeffrey E., Berber-Jimenez M.D., Wang L., and Potter S.M. (1996) An extract of soy flour influences serum cholesterol and thyroid hormones in rats and hamsters. *J. Nutr.* 126: 3046-3053.
- Barth C.A., Scholz-Ahrens K.E., de Vrese M., and Hotze A. (1990) Differences of plasma amino acids following casein or soy protein intake: significance for differences of serum lipid concentrations.
- Bassler and Buchholz H (1993) Methodenbuch Band III. Die chemische Untersuchung von Futtermitteln, *VDLUFA-Verlag, Darmstadt, Germany.*
- Bergeron N., Dishies Y., & Jacques H., (1992) Dietary fish protein modulates high density lipoprotein cholesterol and lipoprotein lipase activity in rabbits. *J. Nutr.* 122: 1731-1737.
- Bergeron N. and Jacques H. (1989) Influences of fish protein as compared to casein and soyprotein on serum lipids, and serum lipoprotein cholesterol levels in the rabbits. *Atherosclerosis* 78: 113-121.
- Bilia A.R., Salvini D., Mazzi G., Vincieri F.F, (2001) Characterization of Calendula Flower, Milk-Thistle Fruit, and Passion Flower Tinctures by HPLC-DAD and HPLC-MS. *Chromatographia* 53 3-4.
- Bradford M.M. (1976) A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72: 255-260.
- Brandsch C., Shukla A., Hirche F., Sangl G.I. & Eder K. (2006) Effects of proteins from beef, pork, and turkey meat on plasma and liver lipids of rats compared with casein and soy protein. *Nutr.* 22: 1162-1170.

- Butte W. (1983) Rapid method for the determination of fatty acid profiles from fats and oils using trimethylsulfonium hydroxide for transesterification. *J. Chromatogr.* 261: 142-145.
- Carroll K.K. (1991) Review of clinical studies on cholesterol-lowering response to soyprotein. *J. Am. Dietetic Assoc.* 91: 820-827.
- Carroll K.K., Kurowska E.M. (1995) Soy consumption and cholesterol reduction: review of animal and human studies. *J. Nutr.* 125: 594S-597S.
- Chirgwin J.M., Przybyla A.E., MacDonald R.J., Rutter W.J. (1979). Isolation of biologically active ribonucleic acid from sources enriched in ribonuclease. *Biochem.* 18: 5294-5299.
- Christiansen E.N., Lund J.S., Rorveit T., Rustan A.C. (1991) Effect of dietary n-3 and n-6 fatty acids on fatty acid desaturation in rat liver. *J. Biol. Chem.* 274: 471-477.
- Clarke R, Daly L, Robinson K, Naughten E, Cahalane S, Fowler B, Graham J. (1991) Hyperhomocysteinemia: an independent risk factor for vascular disease. *N Engl J Med* 324: 1149-55.
- Claudia T.P., Susan M. P., Julie C. H. and Niel F. S. (1998) Intake of Soy protein and Soy protein Extracts Influences Lipid Metabolism and Hepatic Gene Expression in Gerbils. *J. Nutr.* 128: 839-842.
- Damasceno N.R., Gidlund M.A., Goto H., Dias C.T., Okawabata F.S., Abdalla D.S. (2001) Casein and soy protein in experimental atherosclerosis: influence on hypercholesterolemia and lipoprotein oxidation. *Ann. Nutr. Metab* 45: 38-46.
- De Hoff J.L., Davidson J.H. & Kritchevsky D. (1978) An enzymatic assay for determining free and total cholesterol in tissues. *Clin. Chem.* 24: 433-435.
- Desvergne B, Wahli W. (1999) Peroxisome proliferator-activated receptors: nuclear control of metabolism. *Endocr. Rev.* 20: 649-688.
- Deutsch J. (1995). Glucose-6-phosphate dehydrogenase, In Bergmeyer H.U. (ed) Methods of enzymatic analysis 3: 190-197, *verlag chemie, VCH, Weinheim.*

- Dreyer C., Krey G., Mahfoudi A., Keler H., Givel F., Helftenbein G. and Wahli W. (1992) Control of the peroxisomal beta-oxidation pathway by a novel family of nuclear hormone receptors. *Cell*. 61: 879-887.
- Eaton S. (2002) Control of mitochondrial β -oxidation flux. *Prog. Lipid. Res.* 41: 197-239.
- Eder K. and Brandsch C. (2002) Effect of fatty acid composition of rapeseed oil on plasma lipids, fatty acid composition of tissues and susceptibility of low-density lipoprotein to lipid peroxidation in cholesterol-fed hamsters. *Eur. J. Lipid Sci. Technol.* 104: 3-13.
- Eder K. and Kirchgessner M. (1994) J. Trace Elem. Electrol. Health Dis. 8: 63.
- Ericsson C.G., Hamsten A., Nilsson J., Grip L., Svane B., and de Faire U. (1996) Angiographic assessment of effects of bezafibrate on progression of coronary artery disease in young male postinfarction patients. *Lancet* 347: 849-853.
- Field F.J., Albright E. & Mathur S.N. (1987) Regulation of cholesterol esterification by micellar cholesterol in CaCo-2 cells. *J. Lipid Res.* 28: 1057-1066.
- Forsythe W.A (1995) Soy protein, thyroid regulation and cholesterol metabolism. *J. Nutr.* 116: 1165-1171.
- Forsythe W.A., (1986) Comparison of dietary casein or soy protein effects on plasma lipids and hormone concentrations in the gerbil (*Meriones unguiculatus*). *J. Nutr.* 116 1165-1171.
- Foufelle F, Girard J, Ferre P. (1996) Regulation of lipogenic enzyme expression by glucose in liver and adipose tissue: a review of the potential cellular and molecular mechanisms. *Adv Enzyme Regul* 36: 199-226.
- Fukui K., Tachibana N., Fukuda Y., Takamatsu K., Sugano M., (2004) Ethanol washing does not attenuate the hypocholesterolemic potential of soy protein. *Nutrition* 29: 984-990.
- Gardner C.D., Newell K.A., Cherin R., Haskell W.L. (2001) The effects of soy protein with or without isoflavones relative to milk protein on plasma lipids in hypercholesterolemic postmenausal women. *Am. J. Clin. Nutr.* 73: 667-678

- Garg M.L., Thomson A.B., Clandinin M.T. (1998) Effect of dietary cholesterol and omega 3 fatty acids on lipid composition and delta 5-desaturase activity of liver microsomes. *J. Nutr.* 118: 661-668
- Gascon A., Jacques H., Moorjani S., Deshaies Y., Brun LD & Julien P. (1996) Plasma lipoprotein profile and lipolytic activities in response to the substitution of lean white fish for other animal protein sources in premenausal women. *Am. J. Nutr.* 63: 315-321.
- Genest J. Jr. Marcil M., Denis M., & Yu L. (1999) High density lipoprotein in health and disease. *J. Invest. Med.* 47: 31-42.
- Giroux I., Kurowska E.M., Freeman D.J., Carroll K.K., (1999) Addition of arginine but not glycine to lysine plus methionine-enriched diets modulates serum cholesterol and liver phospholipid in rabbits. *J. Nutr.* 129 1807-1813.
- Giudetti A.M., Beynen A.C. Lemmens A.G., Gnoni G.V. & Geelen M.J.H. (2003) Hepatic fatty acid metabolism in rats fed diets with different contents of C18:0, C18:1 cis and C18:1 trans isomers. *Br. J. Nutr.* 90: 887-893.
- Glass C. K. and Witztum J. L. (2001) "Atherosclerosis: the road ahead." *Cell* 104(4): 503-16.
- Grundy S. M. (1994) Second Report of the National Cholesterol Education Program (NCEP) expert panel on detection, evaluation and treatment of high blood cholesterol in adults (Adults treatment panel II). *Circulation* 89: 1329-1445.
- Grundy S.M. (1996) Lipids, nutrition, and coronary heart disease. In *Atherosclerosis and Coronary Artery Disease*. (Fuster, V., Ross, R., and Topol, E.J. eds) V.1: 45-48, Lippincott-Raven Publishers, Philadelphia.
- Grundy S.M., Cleeman J.I., Rifkind B.M., and Kuller L.H. (1999) Cholesterol lowering in the elderly population. *Arch. Intern. Med.* 159: 1670-1678.
- Gudbrandsen O.A., Wergedahl H., Liaset B., Espe M., Berge R.K. (2005) Dietary proteins with isoflavone content or low methionine-glycine and lysine-arginine ratios are hypocholesterolemic and lower the plasma homocysteine level in male Zucker fa/fa rats. *Br. J. Nutr.* 94: 321-326.

- Hagemeyer H., Scholz-Ahrens K.E., Schulte-Coerne H., and Barth C.A. (1990). Plasma amino acids and cholesterol following consumption of dietary casein or soy protein in minipigs. *J. Nutr.* 120: 1305-1311.
- Hara A. and Radin N.S. (1978) Lipid extraction of tissues with a low toxicity solvent. *Anal Biochem* 90: 420-426.
- Hojnacki J.L., Nicolosi R.J. & Hayes K.C. (1976) Densitometric quantification of neutral lipids on ammonium sulphate impregnated thin layer chromatograms. *J. Chromatogr.* 128: 133-139.
- Hokenson J.E. & Austin M.A. (1996) Plasma triglyceride level as a risk factor for cardiovascular disease independent of High-density lipoprotein cholesterol level: a meta-analysis of population-based prospective studies. *J. Cardiovasc. Risk* 3: 213-219
- Horigome T. and Cho Y.S. (1992). Dietary casein and soy bean protein affect the concentration of serum cholesterol, triglyceride and free amino acids in rats. *J. Nutr.* 122: 2273-2282.
- Hortan J.D., Bashmakov Y., Shimomura I. & Shimano H. (1998) Regulation of sterol regulatory element binding proteins in livers of fasted and refed mice. *Proc. Natl. Acad. Sci. USA* 95: 5987-5992.
- Horton J. D., Goldstein J. L. & Brown M. S. (2002) SREBPs: activators of the complete program of cholesterol and fatty acid synthesis in the liver. *J. Clin. Invest.* 109: 1125-1131.
- Hua X., Sakai J., Ho Y.K., Goldstein J.L., Brown M.S. (1995). *J. Bio. Chem.* 270: 29422-29427.
- Hua X., Yokoyama C., Wu J., Briggs M.R., Admon A., Goldstein J.L., Brown M.S., (1993) SREBP-2, a second basic-helix-loop-helix-leucine zipper protein that stimulates transcription by binding to a sterol regulatory element. *Proc. Natl. Acad. Sci U.S.A.* 90 11603-11607.
- Iritani N., Narita R., Fugita T. and Tanaka T. (1985) Effects of fish protein, soybean protein and casein on cholesterol turnover in rats. *J. Nutr. Sci. Vitaminol.* 31: 385-392.
- Iritani N, Nagashima K, Fukuda H, Katsurada A., and Tanaka T., (1986) Effects of Dietary Proteins on Lipogenic enzymes in Rat Liver. *J. Nutr.* 116: 190-197.

- Iritani N., Hosomi H., Fukuda H., and Tada K., (1996) Soybean Protein Suppresses Hepatic Lipogenic Enzyme Gene Expression in Wister Rats. *J. Nutr* 126: 380-388.
- Jacques H., Gascon Bergeron N., Lavigne C., Hurley C., Deshaies Y., Moorjani S. and Julien P. (1995) Role of dietary fish protein in the regulation of plasma lipids. *Can. J. Cardiol.* 11 (suppl G), 63G-71G.
- Jawie J, Nastalek P, Korbut R. (2004) Mouse models of experimental atherosclerosis. *J Physiol Pharmacol* 55:503-17.
- Jenkins D.J, Kendall C.W., Jackson C.J., Connelly P.W., Parker T., Faulkner D, Vidgen E., Cunnane S.C., Leiter L.A., Josse R.G., (2002) Effects of high- and low-isoflavones soyfoods on blood lipids, oxidized LDL, homocysteine, and blood pressure in hyperlipidemic men and women. *Am. J. Clin. Nutr.* 76: 365-372.
- Karpe F., Hamsten A., (1994) Determination of apolipoproteins B-48 and B-100 in triglyceride-rich lipoproteins by analytical SDS PAGE. *J. Lipid Res.* 35: 1311-1317.
- Kern M., Ellison D., Marroquin Y., Ambrose M. & Mosier (2002) Effects of soy protein supplemented with methionine on blood lipids and adiposity of rats *Nutr.* 18: 654-656.
- Kirk E.A., Sutherland P., Wang S.A., Chait A., and Le Boeuf R.C. (1998) Dietary isoflavones reduce plasma cholesterol and atherosclerosis in C 57BL/6 mice but not LDL receptor-deficient mice. *J. Nutr.* 128: 954-959
- Kjeldahl J., 1883, A new method for the determination of nitrogen in organic substances. *Ztschr. anal. Chem.*, 22, 366,
- Krauss R.M. (1998) Atherogenicity of triglyceride-rich lipoproteins. *Am. J. Cardiol.* 81(4A): 13B-17B.
- Kris-Etherton P.M., Etherton T.D., and Yu S. (1997) Efficacy of multiple dietary therapies in reducing cardiovascular disease risk factors. *Am. J. Clin. Nutr.* 65: 560-561.
- Kritchevsky D. and Czarnecki S.K. (2000) Influence of specific amino acids on experimental atherosclerosis in rabbits. *J. Nutr.* 130: (Suppl.): 673S-4S.
- Kritchevsky D., Tepper S.A., Czarnecki S.K., Klurfeld D.M., (1982) Atherogenicity of animal and vegetable protein. Influence of the lysine to arginine ratio. *Atherosclerosis* 41 429-431.

- Kurowska E.M., and Carroll K.K., (1991). Studies on the mechanism of induction of hypercholesterolemia in rabbits by high dietary levels of amino acids. *J. Nutr. Biochem.* 2: 656-662
- Lapre J.A., West C.E., Lovati M.R., Sirtori C.R., Beynen A.C. (1989) Dietary animal proteins and cholesterol metabolism in rats. *Int J Vitam Nutr Res.*59: 93-100.
- Lichtenstein A.H. (1998) Soy Protein, Isoflavones and Cardiovascular Disease Risk. *J. Nutr.* 128: 1589-1592
- Lichtenstein A.H., Jalbert S.M., Adlercreutz H., Goldin B.R., Rasmussen H., Schaefer E.J. & Ausman L.M. (2002) Lipoprotein response to diets high in soy or animal protein with and without isoflavones in moderately hypercholesterolemic subjects. *Arterioscler. Thromb. Vasc. Biol.* 22: 1852-1858.
- Li Z., Lamon-Fava S., Otvos J., Lichtenstein A. H., Velez-Carrasco W., McNamara J. R., Ordovas J.M., and Scaefar E.J. (2004) Fish consumption shifts lipoprotein subfractions to a less atherogenic Pattern in humans. *J.Nutr.* 134: 1724-1728.
- Lovati M.R., Manzoni C., Agostinelli P., Ciappellano S., Mannucci L., Sirtori C.R., (1991) Studies on the mechanism of the cholesterol lowering activity of soy proteins. *Nutr. Metab. Cardiovasc. Dis.* 1 18-24.
- Lovati M.R., Manzoni C., Canavesi A., Sirtori M., Vaccarino V., Marchi M., Gaddi G., Sirtori C.R., (1987) Soybean protein diet increases low density lipoprotein receptor activity in mononuclear cells from hypercholesterolemic patients. *J. Clin. Invest.* 80 1498-1502.
- Lovati M.R., Manzoni C., Corsini A., Granata A., Frattini R., Fumagalli R., Sirtori C.R., (1992) Low density lipoprotein receptor activity is modulated by soybean globulins in cell culture. *J. Nutr.* 122 1971-1978.
- Lovati M.R., Manzoni C., Corsini A., Granata A., Fumagalli R., Sirtori C.R., (1996) 7S globulin from soybean is metabolized in human cell cultures by a specific uptake and degradation system. *J. Nutr.* 126 2831-2842.
- Lovati M.R., Manzoni C., Gianazza E., Arnoldi A., Kuroswka E., Carroll K.K., Sirtori C.R. (2000) Soy protein peptides regulate cholesterol homeostasis in Hep G2 cells. *J. Nutr.* 130 2543-2549.
- Lovati M.R., Manzoni C., Gianazza E., Sirtori C.R., (1998) Soybean protein products as regulators of liver low-density lipoprotein receptors. I. Identification of active β -conglycinin subunits. *J. Agric. Food Chem.* 46 2474-2480.

- Lucas E.A., Khalil D.A., Daggy B.P., and Arjmandi B.H. (2001) Ethanol-extracted soy protein isolate does not modulate serum cholesterol in golden Syrian hamsters: a model of postmenopausal hypercholesterolemia. *J.Nutr.* 131 (2): 211-4
- Luskey K.L. (1988) Regulation of cholesterol synthesis: mechanism for control of HMG CoA reductase. *Recent Prog. Horm. Res.* 44: 35-51.
- Madani S., Lopez S., Blond J.P., Prost J., Belleville J. (1998) Highly purified soy protein is not hypocholesterolemic in rats but stimulates cholesterol synthesis and reduces polyunsaturated fatty acid biosynthesis. *J. Nutr.* 128: 1084-1091.
- Madani S., Prost J., Norce M., & Belleville J. (2003) VLDL metabolism in rats is affected by the concentration and source of dietary protein *J. Nutr.* 133: 4102-4106.
- Mandard S., Müller M., and Kersten S. (2004) Peroxisome proliferator-activated receptor α target genes. *CMLS, Cell. Mol. Life Sci.* 61: 393-416.
- Manzoni C., Lovati M.R., Gianazza E., Sirtori C.R., (1998) Soybean protein products as regulators of liver low-density lipoprotein receptors. II. α - α' Rich commercial soy concentrate and α' deficient mutant differently affect low-density lipoprotein receptor activation. *J. Agric. Food Chem.* 46 2481-2484.
- Marlett J.A. & Fisher M.H. (2002) A poorly fermented gel from psyllium seed husk increases excreta moisture and bile acid excretion in rats. *J. Nutr.* 132: 2638-2643.
- Mason T.M. (1998) The role of factors that regulate the synthesis and secretion of very low density lipoprotein by hepatocytes. *Crit. Rev. Clin. Lab. Sci.* 35: 461-487
- Meddings J., Spady D., Dietschy J., (1986) Kinetic constants for receptor-dependent and receptor-independent low density lipoprotein transport in the tissues of the rat and hamster. *J. Clin. Invest.* 77: 1474-1481.
- Morita T., Oh-hasi A., Takai K., Ikai M., Kasaoka S., and Kiriya S. (1997) Cholesterol-lowering effects of soybean, potato and rice protein depend on their low methionine contents in rats fed a cholesterol-free purified diet. *J.Nutr.* 127: 470-477.
- Mullen E., Brown R.M., Osborne T.F., Shay, (2004) Soy isoflavones affect sterol regulatory element binding proteins (SREBPs) and SREBP-regulated genes in HepG2 cells. *J. Nutr.* 134: 2942-2947.

- Murphy C, Parini P, Wang J, Björkhem I, Eggertsen G, Gäfvels M. (2005) Cholic acid as a key regulator of cholesterol synthesis, intestinal absorption and hepatic storage in mice. *Biochim Biophys Acta* 1735: 167-75.
- Nagata Y., Ishiwaki, N. & Sugano M. (1982) Studies on the mechanism of antihypercholesterolemic action of soy protein and soy protein type amino acid mixtures in relation to the casein counterparts in rats. *J. Nutr.* 112: 1614-1625.
- Nagata Y., Tanaka K., Sugano M. (1981) Serum and liver cholesterol levels of rats and mice fed soy-bean protein or casein. *J. Nutr. Sci. Vitaminol (Tokyo)* 27: 583-93.
- Nagaoka S., Miwa K., Eto M., Kuzuya Y., Hori G & Yamamoto K (1999) Soy protein peptic hydrolysate with bound phospholipids decreases micellar solubility and cholesterol absorption in rats and caco cells. *J. Nutr.* 112: 1614-1625.
- Nanami K., Oda H., & Yokogoshi H. (1996) Antihypercholesterolemic action of taurine on streptozotocin-diabetic rats or on rats fed a high cholesterol diet. *Adv. Exp. Med. Biol.* 403: 561-568.
- Naumann C. and Bassler R. (1976) Methodenbuch Band III. Die chemische Untersuchung von Futtermitteln, *VDLUFA-Verlag, Darmstadt, Germany, 1993.*
- Nehler M.R. Taylor L.M. & Porter J.M. (1997) Homocysteine as a risk factor for atherosclerosis: a review *Cardiovasc. Pathol.* 6: 1-9
- Nepokroef C.M., Lakshmanan M.R., Porter J.W. (1975). Fatty acid synthase from rat liver. *Methods Enzymol.* 35: 37-44.
- Ni W, Tsuda Z., Sakono M. and Imaizumi K (1998) Dietary soy protein isolate, compared with casein, reduces atherosclerotic lesion area in Apolipoprotein E- deficient mice. *J. Nutr.* 128: 1884-1889.
- Ni W., Yoshida S., Tsuda, Y., Nagao K., Sato M & Imaizumi K. (1999) Ethanol extracted soy protein isolate results in elevation of serum cholesterol in exogenously hypercholesterolemic rats. *Lipids* 34: 713-716.

- Park T. and Lee K. (1998) Dietary taurine supplementation reduces plasma and liver cholesterol and triglyceride levels in rats fed a high-cholesterol or a cholesterol-free diet. *Adv. Exp. Med. Biol.* 442: 319-25.
- Peluso M.R., Winters T.A., Shanahan M.F., Banz W.J. (2000) A cooperative interaction between soy protein and its isoflavone-enriched fraction lowers hepatic lipid in male obese Zucker rats and reduces blood platelet sensitivity in male Sprague-Dawley rats. *J Nutr* 130: 2333-42.
- Pfeuffer M. and Barth C.A. (1986) Modulation of very low density lipoprotein secretion by dietary protein is age dependent in rats. *Ann. Nutr. Metab.* 30: 281-288.
- Potter S. M. (1995) Overview of proposed mechanisms for the hypocholesterolemic effect of soy. *J. Nutr.* 125: 606S-611S
- Potter S.M. (1996) Soy protein and serum Lipids. *Curr Opin Lipidol.* 7: 260-264.
- Potter S.M., Baum J.A., Teng H., Stillman R.J., Shay N.F., and Erdman J.W. Jr. (1998) Soy protein and isoflavones: their effects on blood lipids and bone density in postmenopausal women. *Am. J. Clin. Nutr.* 68 (Suppl): 1375S-1379S.
- Rawson R.B. (2003) The SEBP pathway-Insights from insig and insects. *nature* 4: 631-640
- Reeves P.G, Nielsen F.H. and Fahey G.C. Jr. (1993) AIN-93 purified diets for laboratory rodents: Final report of the American Institute of Nutrition ad hoc Writing Committee on the reformulation of the AIN-76A rodent diet. *J Nutr* 123: 1939-1951.
- Refsum H, Ueland PM, Nygård O, Vollset SE. (1998) Homocysteine and cardiovascular disease. *Annu Rev Med* 49: 31-62.
- Ringseis R. and Kder K. (2005) Effects of dietary fat and oxidized cholesterol on gene expression in rat liver as assessed by cDNA array analysis. *Eur.J. Nutr.* 44: 231-241.
- Sanchez A., Hubbard R.W., Smit E., and Hilton G.F., (1988). Testing a mechanism of control in human cholesterol metabolism: Relation of arginine and glycine to insulin and glucagons. *Atherosclerosis* 71: 87-92.

- Sautier C., Dieng K., Flament C., Doucet C., Suquet J.P., Lemonnier D. (1983) Effects of whey protein, casein, soya-bean and sunflower proteins on the serum , tissue and faecal steroids in rats. *J. Nutr.* 49 (3): 313-319.
- Sautier C., Flament C., Doucet C., Suquet J. P., (1986) Effect of eight dietary proteins and their amino acid content on serum, hepatic and fecal steroids in rats. *Nutr. Rep Int*, 34: 1051-1061.
- Schuster R. (1988) Determination of amino acids in biological, pharmaceutical, plant and food samples by automated pre-column derivatization and high-performance liquid chromatography. *J. Chromatogr.* 431: 271-284.
- Schaefer E.J., Lichtenstein A.H., Lamon-Fava S., Contois J.H., Li Z., Rossmussen H., McNamara J.R. & Ordovas J.M. (1995) Effect of National Cholesterol Education Program Step 2 diets in normolipidemic and hyperlipidemic middle aged and elderly men and women. *Arterioscler. Throm. Vasc. Biol.*15: 1079-1085.
- Schaefer E.J., Lichtenstein A.H., Lamon-Fava S., Contois J.H., Li Z., Goldin B. R., Rossmussen H., McNamara J.R. & Ordovas J.M. (1996) Effect of National Cholesterol Education Program Step 2 diets relatively high or relatively low in fish-derived fatty acids on plasma lipoproteins in middle aged and elderly subjects. *Am. J. Clin. Nutr.* 63: 234-241.
- Semenkovich C.F. (1997) Regulation of fatty acid synthase (FAS). *Prog. Lipid Res.* 36: 43-53.
- Shelness G.S., Sellar J.A (2001) very low density lipoprotein assembly and secretion. *Curr. Opin. Lipidol.* 12: 151-157.
- Shimano H., Yahagik N., Amemiya-Kudo M., Hasty A.H., Osuga J., Tamura Y., Shionoiri F., Lizuka Y., Ohashi K., Harada K., Gotoda T., Ishibashi S. & Yamada N. (1999) Sterol regulatory element binding protein-1 as a key transcription factor for nutritional induction of lipogenic enzyme genes. *J. Biol. Chem.* 274: 35832-35839.
- Shimomura I., Shimano H., Horton J.D., Goldstein J. L. & Brown M.S. (1997). *J. Clin. Invest.* 99: 838-845.

- Shukla A., Bettzieche A., Hirche F., Brandsch C., Stangl G.I., Eder K. (2006 a) Dietary fish protein alters blood lipid concentrations and hepatic genes involved in cholesterol homeostasis in the rat model. *Br. J. Nutr.* 96: 674-682.
- Shukla A., Brandsch C., Bettzieche A., Hirche F., Stangl G.I., Eder K. (2006 b) Isoflavone-poor soy protein alters lipid metabolism of rats by SREBP-mediated down regulation of hepatic genes. *J. Nutr. Biochem.* (published online).
- Sirtori C.R., Galli G., Lovati M.R., Carrara P., Bosisio E., Kienle M.G., (1984) Effects of dietary proteins on the regulation of liver lipoprotein receptors in rats. *J. Nutr.* 114: 1493-1502.
- Sirtori, C.R., Lovati, M.R. (2001) Soy Proteins and Cardiovascular Disease. *Curr Atheroscler Rep.* 3: 47-53.
- Sirtori C.R., Lovati M.R., Castiglioni S., Duranti M., Magni C., Moranti S., D'Agostina A. & Arnoldi A. (2004) proteins of white lupin seed, a naturally isoflavone-poor legume, reduced cholesterolemia in rats and increase LDL receptor activity in HepG2 cells. *J. Nutr.* 134: 18-23.
- Sirtori C.R., Lovati M.R., Manzoni C., Gianazza E., Bondioli A., Staels B., Auwerx J. (1998) Reduction of serum cholesterol by soybean proteins: clinical experience and potential molecular mechanisms. *Nutr Metab Cardiovasc Dis* 8: 334-40.
- Sparks J. D., PhuIng T.L., Bolognino M., Cianci J., Khurana R., Peterson R.G., Soven M.P., Corsetti J.P. and Sparks C.E. (1998) Lipoproteins alterations in 10- and 20-weeks old Zucker diabetic fatty rats: hyperinsulinemic versus insulinopenic hyperglycemia. *Metabolism* 47: 1315-1324.
- Stamler J., Wentworth D., and Neaton J.D. (1986) Is relationship between serum cholesterol and risk of premature death from coronary heart disease continuous and graded? findings in 356,222 primary screenees of the Multiple Risk Factor Intervention Trial (MRFIT). *J. Am. Med. Assoc.* 256: 2823-2828.
- Sugano M., Tanaka K. and Idle T. (1982) Secretion of cholesterol, triglyceride and apolipoprotein A-I by isolated perfused liver from rats fed soybean protein and casein on their amino acid mixtures. *J. Nutr.* 112: 855-862.

- Sugiyama K, Kanamori H., Akachi T, and Yamakawa A., (1996). Amino acid composition of dietary proteins affects plasma cholesterol concentration through alteration of hepatic phospholipids metabolism in rats fed a cholesterol free diet. *J. Nutr. Biochem.* 7: 40-48
- Sugiyama K. and Muramatsu K. (1990) Significance of amino acid composition of dietary protein in regulation of plasma cholesterol. *J. Nutr. Sci. Vitaminol.* 36: 105S-110S.
- Sugiyama K., Ohkawa S. & Muramatsu K. (1986) Relationship between amino acid composition of diet and plasma cholesterol level in growing rats fed a high cholesterol diet. *J Nutr Sci Vitaminol* 32: 413-423.
- Sugiyama K., Yamakawa A.A., Kumazawa A., Saeki S. (1997) Methionine content of dietary proteins affects the molecular species composition of plasma phosphatidylcholine in rats fed a cholesterol-free diet. *J. Nutr.* 127: 600-607.
- Sülzle A., Hirsch F., and Eder K., (2004) Thermally oxidized fat upregulates the expression of target genes of PPAR α in rat liver. *J. Nutr.* 134: 1375-1383.
- Tachibana N., Matsumoto I., Fukui K., Arai S., Kato H., Abe K. & Takamatsu K. (2005) Intake of soy protein isolate alters hepatic gene expression in rats. *J. Agric. Food Chem.* 53: 4253-4257.
- Terpstra AHM, van Tintelen G., West C. (1982) The effect of semipurified diet containing different proportions of either casein or soybean protein on the concentration of cholesterol on whole serum, serum lipoproteins and liver in male and female rats. *Atherosclerosis* 42: 85-92.
- Terpstra AHM, van Tintelen G., West C. (1983) Serum lipid lipoprotein composition and liver cholesterol in genetically obese Zucker rat fed semipurified diets containing either casein or soy protein. *Ann. Nutr. Metab.* 27: 132-136.
- Toborek M., Kopiczna-Gzebieniak E., Drozd M. & Wiczorek M (1995) Increased lipid peroxidation as a mechanism of methionine- induced atherosclerosis in rabbits. *Atherosclerosis* 115: 217-224.

- Tovar A., Manzano N., Torres N. (2005) Metabolism of cholesterol and fatty acids in nephrotic syndrome and its regulation by sterol regulatory element binding protein (SREBP's). Effect of soy protein consumption. *Gac Med Mex* 141: 407-415.
- Tover A.R, Murguia F., Cruz C., et al. (2002) A Soy Protein Diet Alters Hepatic Lipid Metabolism, Gene Expression and Reduces Serum Lipids and renal Fibrogenic Cytokines in Rats with Chronic Nephrotic Syndrome. *J. Nutr.* 132: 2562-2569.
- Tsai Po-Jung & Huang Po-Chao (1999) Effects of isoflavones containing soyprotein isolate compared with fish protein on serum lipids and susceptibility of low density lipoprotein and liver lipids to in vitro oxidation in hamsters. *J. Nutr. Biochem.* 10: 631-637.
- Tso P., Drake D.S., Black D.D. & Sabesin S.M. (1984) Evidence for separate pathways of chylomicron and very low-density lipoprotein assembly and transport by rat small intestine. *Am. J. Physiol.* 247: G599-G610.
- Vallett S.M., Sanchez H.B., Rosenfeld J.M. & Osborne T.F. (1996) A direct role of sterol regulatory element binding protein in activation of 3-hydroxy-3-methylglutaryl coenzyme A reductase gene. *J Biol Chem* 271: 12247-12253.
- Van Raaij J.M.A., Katan M.B., West C.E., Hautvast J.G.A.J., and Hermus R.J.J. (1981) Effects of casein versus soy protein diets on serum cholesterol and lipoproteins in young healthy volunteers. *Am. J Clin Nutr.* 34: 1261-1271.
- Van Raaij J.M.A., Katan M.B., West C.E., Hautvast J.G.A.J. (1982) Influence of diets containing casein, soy protein isolate, and soy protein concentrate on serum cholesterol and lipoproteins in middle- aged volunteers. *Am. J. Clin. Nutr.* 35: 925-934.
- Vigna G.B., Pansini F., Bonaccorsi G., Albertazzi P., Donega P., Zanotti L., De Aloysio D., Mollica G., & Fellin R. (2000) Plasma lipoproteins in soy-treated postmenausal women: a double blind, placebo-controlled trial. *Nutr. Metab. Cardiovasc. Dis.* 10: 315-322.
- Vlahecevic Z.R., Pandak W.M. & Stravitz R.T. (1999) Regulation of bile acid biosynthesis. *Gastroenterol Clin North Am* 28: 1-25.

- Wang W.W., Smith E. O'Brian, Stuff J.E., Hachey D.L., Heird W.C. and Pownell H.J. (1998) Cholesterol lowering effect of soy protein in normocholesterolemic and hypercholesterolemic men. *Am. J. Clin. Nutr.* 68 (suppl), 1385S-1389S.
- Welch G.N. and Loscalzo J. (1998) Homocysteine and atherothrombosis. *N Engl J Med* 338: 1042-50.
- Wergedahl H., Liaset B., Gudbrandsen O.A., Lied E., Espe M., Muna Z., Mork S., & Berge R.K., (2004) Fish protein hydrolysate reduces plasma total cholesterol, increases the proportion of HDL cholesterol, and lowers acyl-CoA acyltransferase activity in liver of Zucker rats. *J.Nutr.* 134: 1320-1327.
- Woo CWH, Siow Y.L., Pierce G.N., Choy P.C., Minuk G.Y., Mymin D. & Karmin O. (2005) Hyperhomocysteinemia induces hepatic cholesterol biosynthesis and lipid accumulation via activation of transcription factors. *Am. J. Physiol. Endocrinol. Metab.* 288, E1002-E1010.
- Wright S.M., Salter A.M. (1998) Effect of soy protein on plasma cholesterol and bile acid excretion in hamsters. *Comp. Biochem. Physiol. B. Biochem. Mol. Biol* 119: 247-254.
- Yabe D., Brown M.S. & Goldstein J.L. (2002) Insig-2, a second endoplasmic reticulum protein that binds SCAP and blocks export of sterol regulatory element-binding proteins. *Proc. Natl. Acad. Sci. U.S.A.* 99: 12753-12758.
- Yan C.C., Bravo E., Cantafora A. (1993) Effect of taurine levels on liver lipid metabolism: an in vivo study in the rat. *Proc. Soc. Exp. Biol. Med.* 202: 88-96.
- Yang T., Espenshade P.J., Wright M.E., Yabe D., Gong Y., Aebersold R., Goldstein J.L. & Brown M.S. (2002) Crucial Step in Cholesterol Homeostasis: Sterols promote Binding of SCAP to INSIG-1, a Membrane Protein that Facilitates Retention of SREBPs in ER. *Cell* 110: 489-500.
- Zhang X. & Beynen A.C. (1993) Influence of dietary fish proteins on plasma and liver cholesterol concentrations in rats. *Br J Nutr* 69: 767-777.