

Anhang

Literaturverzeichnis

- Aloyz, R. S., Bamji, S. X., Pozniak, C. D., Toma, J. G., Atwal, J., Kaplan, D. R., and Miller, F. D. (1998). p53 is essential for developmental neuron death as regulated by the TrkA and p75 neurotrophin receptors. *J. Cell Biol.*, **143**:1691-1703.
- Anfinsen, C. B. (1973). Principles that govern the folding of protein chains. *Science*, **181**:223-230.
- Angeletti, R. H. and Bradshaw, R. A. (1971). Nerve growth factor from mouse submaxillary gland: amino acid sequence. *Proc. Natl. Acad. Sci. U. S. A.*, **68**:2417-2420.
- Apfel, S. C., Arezzo, J. C., Lipson, L., and Kessler, J. A. (1992). Nerve growth factor prevents experimental cisplatin neuropathy. *Ann. Neurol.*, **31**:76-80.
- Apfel, S. C. and Kessler, J. A. (1996). Neurotrophic factors in the treatment of peripheral neuropathy. *Ciba Found. Symp.*, **196**:98-108.
- Aurikko, J. P., Ruotolo, B. T., Grossmann, J. G., Moncrieffe, M. C., Stephens, E., Leppanen, V. M., Robinson, C. V., Saarma, M., Bradshaw, R. A., and Blundell, T. L. (2005). Characterization of symmetric complexes of nerve growth factor and the ectodomain of the pan-neurotrophin receptor, p75NTR. *J. Biol. Chem.*, **280**:33453-33460.
- Bai, Y., Milne, J. S., Mayne, L., and Englander, S. W. (1993). Primary structure effects on peptide group hydrogen exchange. *Proteins*, **17**:75-86.
- Baker, D., Sohl, J. L., and Agard, D. A. (1992). A protein-folding reaction under kinetic control. *Nature*, **356**:263-265.
- Bamji, S. X., Majdan, M., Pozniak, C. D., Belliveau, D. J., Aloyz, R., Kohn, J., Causing, C. G., and Miller, F. D. (1998). The p75 neurotrophin receptor mediates neuronal apoptosis and is essential for naturally occurring sympathetic neuron death. *J. Cell Biol.*, **140**:911-923.
- Barbacid, M. (1995). Neurotrophic factors and their receptors. *Curr. Opin. Cell Biol.*, **7**:148-155.
- Barde, Y. A. (1990). The nerve growth factor family. *Prog. Growth Factor Res.*, **2**:237-248.
- Bauskin, A. R., Zhang, H. P., Fairlie, W. D., He, X. Y., Russell, P. K., Moore, A. G., Brown, D. A., Stanley, K. K., and Breit, S. N. (2000). The propeptide of macrophage inhibitory cytokine (MIC-1), a TGF-beta superfamily member, acts as a quality control determinant for correctly folded MIC-1. *EMBO J.*, **19**:2212-2220.

- Bax, B., Blundell, T. L., Murray-Rust, J., and McDonald, N. Q. (1997). Structure of mouse 7S NGF: a complex of nerve growth factor with four binding proteins. *Structure*, **5**:1275-1285.
- Beattie, M. S., Harrington, A. W., Lee, R., Kim, J. Y., Boyce, S. L., Longo, F. M., Bresnahan, J. C., Hempstead, B. L., and Yoon, S. O. (2002). ProNGF induces p75-mediated death of oligodendrocytes following spinal cord injury. *Neuron*, **36**:375-386.
- Belliveau, D. J., Krivko, I., Kohn, J., Lachance, C., Pozniak, C., Rusakov, D., Kaplan, D., and Miller, F. D. (1997). NGF and neurotrophin-3 both activate TrkA on sympathetic neurons but differentially regulate survival and neuritogenesis. *J. Cell Biol.*, **136**:375-388.
- Berger, E. A. and Shooter, E. M. (1977). Evidence for pro-beta-nerve growth factor, a biosynthetic precursor to beta-nerve growth factor. *Proc. Natl. Acad. Sci. USA*, **74**:3647-3651.
- Bibel, M., Hoppe, E., and Barde, Y. A. (1999). Biochemical and functional interactions between the neurotrophin receptors trk and p75NTR. *EMBO J.*, **18**:616-622.
- Bohm, G., Muhr, R., and Jaenicke, R. (1992). Quantitative analysis of protein far UV circular dichroism spectra by neural networks. *Protein Eng*, **5**:191-195.
- Bothwell, M. A. and Shooter, E. M. (1977). Dissociation equilibrium constant of β nerve growth factor. *J. Biol. Chem.*, **252**:8532-8536.
- Böttinger, E. P., Factor, V. M., Tsang, M. L., Weatherbee, J. A., Kopp, J. B., Qian, S. W., Wakefield, L. M., Roberts, A. B., Thorgeirsson, S. S., and Sporn, M. B. (1996). The recombinant proregion of transforming growth factor beta1 (latency-associated peptide) inhibits active transforming growth factor beta1 in transgenic mice. *Proc. Natl. Acad. Sci. USA*, **93**:5877-5882.
- Bowie, J. U. and Sauer, R. T. (1989). Equilibrium dissociation and unfolding of the Arc repressor dimer. *Biochemistry*, **28**:7139-7143.
- Bradshaw, R. A., Blundell, T. L., Lapatto, R., McDonald, N. Q., and Murray-Rust, J. (1993). Nerve growth factor revisited. *Trends Biochem. Sci.*, **18**:48-52.
- Brann, A. B., Tcherpakov, M., Williams, I. M., Futerman, A. H., and Fainzilber, M. (2002). Nerve growth factor-induced p75-mediated death of cultured hippocampal neurons is age-dependent and transduced through ceramide generated by neutral sphingomyelinase. *J. Biol. Chem.*, **277**:9812-9818.
- Bresnahan, P. A., Leduc, R., Thomas, L., Thorner, J., Gibson, H. L., Brake, A. J., Barr, P. J., and Thomas, G. (1990). Human fur gene encodes a yeast KEX2-like endoprotease that cleaves pro-beta-NGF in vivo. *J. Cell Biol.*, **111**:2851-2859.
- Bryan, P. N. (2002). Prodomains and protein folding catalysis. *Chem. Rev.*, **102**:4805-4816.
- Buczek, O., Olivera, B. M., and Bulaj, G. (2004). Propeptide does not act as an intramolecular chaperone but facilitates protein disulfide isomerase-assisted folding of a conotoxin precursor. *Biochemistry*, **43**:1093-1101.

- Buevich, A. V., Shinde, U. P., Inouye, M., and Baum, J. (2001). Backbone dynamics of the natively unfolded pro-peptide of subtilisin by heteronuclear NMR relaxation studies. *J. Biomol. NMR*, **20**:233-249.
- Casaccia-Bonnet, P., Carter, B. D., Dobrowsky, R. T., and Chao, M. V. (1996). Death of oligodendrocytes mediated by the interaction of nerve growth factor with its receptor p75. *Nature*, **383**:716-719.
- Casademunt, E., Carter, B. D., Benzel, I., Frade, J. M., Dechant, G., and Barde, Y. A. (1999). The zinc finger protein NRIF interacts with the neurotrophin receptor p75(NTR) and participates in programmed cell death. *EMBO J.*, **18**:6050-6061.
- Chakrabarti, S., Sima, A. A., Lee, J., Brachet, P., and Dicou, E. (1990). Nerve growth factor (NGF), proNGF and NGF receptor-like immunoreactivity in BB rat retina. *Brain Res.*, **523**:11-15.
- Chao, M. V. (2003). Neurotrophins and their receptors: a convergence point for many signalling pathways. *Nat. Rev. Neurosci.*, **4**:299-309.
- Chao, M. V. and Hempstead, B. L. (1995). p75 and Trk: a two-receptor system. *Trends Neurosci.*, **18**:321-326.
- Chen, Y., Dicou, E., and Djakiew, D. (1997). Characterization of nerve growth factor precursor protein expression in rat round spermatids and the trophic effects of nerve growth factor in the maintenance of Sertoli cell viability. *Mol. Cell Endocrinol.*, **127**:129-136.
- Cohen, P., Sutter, A., Landreth, G., Zimmermann, A., and Shooter, E. M. (1980). Oxidation of tryptophan-21 alters the biological activity and receptor binding characteristics of mouse nerve growth factor. *J. Biol. Chem.*, **255**:2949-2954.
- Cohen, S. (1960). Purification of a nerve-growth promoting protein from the mouse salivary gland and its neuro-cytotoxic antiserum. *Proc. Natl. Acad. Sci. USA*, **46**:302-311.
- Creighton, T. E. (1991). Molecular chaperones. Unfolding protein folding. *Nature*, **352**:17-18.
- Daopin, S., Piez, K. A., Ogawa, Y., and Davies, D. R. (1992). Crystal structure of transforming growth factor-beta 2: an unusual fold for the superfamily. *Science*, **257**:369-373.
- Darling, T. L., Petrides, P. E., Beguin, P., Frey, P., Shooter, E. M., Selby, M., and Rutter, W. J. (1983). The biosynthesis and processing of proteins in the mouse 7S nerve growth factor complex. *Cold Spring Harb. Symp. Quant. Biol.*, **48 Pt 1**:427-434.
- Davey, F. and Davies, A. M. (1998). TrkB signalling inhibits p75-mediated apoptosis induced by nerve growth factor in embryonic proprioceptive neurons. *Curr. Biol.*, **8**:915-918.
- De Bernardes, Clark E., Schwarz, E., and Rudolph, R. (1999). Inhibition of aggregation side reactions during in vitro protein folding. *Methods Enzymol.*, **309**:217-236.
- De Young, L. R., Burton, L. E., Liu, J., Powell, M. F., Schmelzer, C. H., and Skelton, N. J. (1996). RhNGF slow unfolding is not due to proline isomerization: possibility of a cystine knot loop-threading mechanism. *Protein Sci.*, **5**:1554-1566.

- De Young, L. R., Schmelzer, C. H., and Burton, L. E. (1999). A common mechanism for recombinant human NGF, BDNF, NT-3, and murine NGF slow unfolding. *Protein Sci.*, **8**:2513-2518.
- Dechant, G. (2001). Molecular interactions between neurotrophin receptors. *Cell Tissue Res.*, **305**:229-238.
- Degnin, C., Jean, F., Thomas, G., and Christian, J. L. (2004). Cleavages within the prodomain direct intracellular trafficking and degradation of mature bone morphogenetic protein-4. *Mol. Biol. Cell*, **15**:5012-5020.
- Delsite, R. and Djakiew, D. (1999). Characterization of nerve growth factor precursor protein expression by human prostate stromal cells: a role in selective neurotrophin stimulation of prostate epithelial cell growth. *Prostate*, **41**:39-48.
- Dicou, E., Pflug, B., Magazin, M., Lehy, T., Djakiew, D., Ferrara, P., Nerriere, V., and Harvie, D. (1997). Two peptides derived from the nerve growth factor precursor are biologically active. *J Cell Biol*, **136**:389-398.
- Drinkwater, C. C., Suter, U., Angst, C., and Shooter, E. M. (1991). Mutation of tryptophan-21 in mouse nerve growth factor (NGF) affects binding to the fast NGF receptor but not induction of neurites on PC12 cells. *Proc. Biol. Sci.*, **246**:307-313.
- Eder, J., Rheinnecker, M., and Fersht, A. R. (1993). Folding of subtilisin BPN': role of the pro-sequence. *J. Mol. Biol.*, **233**:293-304.
- Edwards, R. H., Selby, M. J., Mobley, W. C., Weinrich, S. L., Hruby, D. E., and Rutter, W. J. (1988). Processing and secretion of nerve growth factor: expression in mammalian cells with a vaccinia virus vector. *Mol. Cell Biol.*, **8**:2456-2464.
- Edwards, R. H., Selby, M. J., and Rutter, W. J. (1986). Differential RNA splicing predicts two distinct nerve growth factor precursors. *Nature*, **319**:784-787.
- Eisenberg, D., Weiss, R. M., and Terwilliger, T. C. (1984). The hydrophobic moment detects periodicity in protein hydrophobicity. *Proc. Natl. Acad. Sci. USA*, **81**:140-144.
- Ernfors, P., Ibanez, C. F., Ebendal, T., Olson, L., and Persson, H. (1990). Molecular cloning and neurotrophic activities of a protein with structural similarities to nerve growth factor: developmental and topographical expression in the brain. *Proc. Natl. Acad. Sci. USA*, **87**:5454-5458.
- Esposito, D., Patel, P., Stephens, R. M., Perez, P., Chao, M. V., Kaplan, D. R., and Hempstead, B. L. (2001). The cytoplasmic and transmembrane domains of the p75 and Trk A receptors regulate high affinity binding to nerve growth factor. *J. Biol. Chem.*, **276**:32687-32695.
- Fahnestock, M., Michalski, B., Xu, B., and Coughlin, M. D. (2001). The precursor pro-nerve growth factor is the predominant form of nerve growth factor in brain and is increased in Alzheimer's disease. *Mol. Cell Neurosci.*, **18**:210-220.
- Fahnestock, M., Yu, G., and Coughlin, M. D. (2004a). ProNGF: a neurotrophic or an apoptotic molecule? *Prog. Brain Res.*, **146**:107-110.

- Fahnestock, M., Yu, G., Michalski, B., Mathew, S., Colquhoun, A., Ross, G. M., and Coughlin, M. D. (2004b). The nerve growth factor precursor proNGF exhibits neurotrophic activity but is less active than mature nerve growth factor. *J. Neurochem.*, **89**:581-592.
- Fairlie, W. D., Zhang, H. P., Wu, W. M., Pankhurst, S. L., Bauskin, A. R., Russell, P. K., Brown, P. K., and Breit, S. N. (2001). The propeptide of the transforming growth factor-beta superfamily member, macrophage inhibitory cytokine-1 (MIC-1), is a multifunctional domain that can facilitate protein folding and secretion. *J. Biol. Chem.*, **276**:16911-16918.
- Farhadi, H., Pareek, S., Day, R., Dong, W., Chretien, M., Bergeron, J. J., Seidah, N. G., and Murphy, R. A. (1997). Prohormone convertases in mouse submandibular gland: colocalization of furin and nerve growth factor. *J. Histochem. Cytochem.*, **45**:795-804.
- Fayard, B., Loeffler, S., Weis, J., Vogelín, E., and Kruttgen, A. (2005). The secreted brain-derived neurotrophic factor precursor pro-BDNF binds to TrkB and p75NTR but not to TrkA or TrkC. *J. Neurosci. Res.*, **80**:18-28.
- Feeney, B. and Clark, A. C. (2005). Reassembly of active caspase-3 is facilitated by the propeptide. *J. Biol. Chem.*, **280**:39772-39785.
- Fischer, G., Tradler, T., and Zarnt, T. (1998). The mode of action of peptidyl prolyl cis/trans isomerases in vivo: binding vs. catalysis. *FEBS Lett.*, **426**:17-20.
- Fox, T., de Miguel E., Mort, J. S., and Storer, A. C. (1992). Potent slow-binding inhibition of cathepsin B by its propeptide. *Biochemistry*, **31**:12571-12576.
- Frade, J. M., Rodriguez-Tebar, A., and Barde, Y. A. (1996). Induction of cell death by endogenous nerve growth factor through its p75 receptor. *Nature*, **383**:166-168.
- Freedman, R. B., Hirst, T. R., and Tuite, M. F. (1994). Protein disulphide isomerase: building bridges in protein folding. *Trends Biochem. Sci.*, **19**:331-336.
- Friedman, W. J. (2000). Neurotrophins induce death of hippocampal neurons via the p75 receptor. *J. Neurosci.*, **20**:6340-6346.
- Gallagher, T., Gilliland, G., Wang, L., and Bryan, P. (1995). The prosegment-subtilisin BPN' complex: crystal structure of a specific 'foldase'. *Structure*, **3**:907-914.
- Gentry, L. E. and Nash, B. W. (1990). The pro domain of pre-pro-transforming growth factor beta 1 when independently expressed is a functional binding protein for the mature growth factor. *Biochemistry*, **29**:6851-6857.
- Gray, A. M. and Mason, A. J. (1990). Requirement for activin A and transforming growth factor--beta 1 pro- regions in homodimer assembly. *Science*, **247**:1328-1330.
- Greene, L. A., Shooter, E. M., and Varon, S. (1969). Subunit interaction and enzymatic activity of mouse 7S nerve growth factor. *Biochemistry*, **8**:3735-3741.
- Grewal, S. S., York, R. D., and Stork, P. J. (1999). Extracellular-signal-regulated kinase signalling in neurons. *Curr. Opin. Neurobiol.*, **9**:544-553.

- Hasan, W., Pedchenko, T., Krizsan-Agbas, D., Baum, L., and Smith, P. G. (2003). Sympathetic neurons synthesize and secrete pro-nerve growth factor protein. *J. Neurobiol.*, **57**:38-53.
- He, X. L. and Garcia, K. C. (2004). Structure of nerve growth factor complexed with the shared neurotrophin receptor p75. *Science*, **304**:870-875.
- Hempstead, B. L. (2002). The many faces of p75NTR. *Curr. Opin. Neurobiol.*, **12**:260-267.
- Hempstead, B. L., Martin-Zanca, D., Kaplan, D. R., Parada, L. F., and Chao, M. V. (1991). High-affinity NGF binding requires coexpression of the trk proto-oncogene and the low-affinity NGF receptor. *Nature*, **350**:678-683.
- Hermey, G., Keat, S. J., Madsen, P., Jacobsen, C., Petersen, C. M., and Gliemann, J. (2003). Characterization of sorCS1, an alternatively spliced receptor with completely different cytoplasmic domains that mediate different trafficking in cells. *J. Biol. Chem.*, **278**:7390-7396.
- Hidaka, Y., Ohno, M., Hemmasi, B., Hill, O., Forssmann, W. G., and Shimonishi, Y. (1998). In vitro disulfide-coupled folding of guanylyl cyclase-activating peptide and its precursor protein. *Biochemistry*, **37**:8498-8507.
- Hillger, F., Herr, G., Rudolph, R., and Schwarz, E. (2005). Biophysical comparison of BMP-2, ProBMP-2, and the free pro-peptide reveals stabilization of the pro-peptide by the mature growth factor. *J. Biol. Chem.*, **280**:14974-14980.
- Hohn, A., Leibrock, J., Bailey, K., and Barde, Y. A. (1990). Identification and characterization of a novel member of the nerve growth factor/brain-derived neurotrophic factor family. *Nature*, **344**:339-341.
- Ibanez, C. F., Ebendal, T., Barbany, G., Murray-Rust, J., Blundell, T. L., and Persson, H. (1992). Disruption of the low affinity receptor-binding site in NGF allows neuronal survival and differentiation by binding to the trk gene product. *Cell*, **69**:329-341.
- Ibanez, C. F., Ilag, L. L., Murray-Rust, J., and Persson, H. (1993). An extended surface of binding to Trk tyrosine kinase receptors in NGF and BDNF allows the engineering of a multifunctional pan-neurotrophin. *EMBO J.*, **12**:2281-2293.
- Ikemura, H., Takagi, H., and Inouye, M. (1987). Requirement of pro-sequence for the production of active subtilisin E in *Escherichia coli*. *J. Biol. Chem.*, **262**:7859-7864.
- Inouye, M. (1991). Intramolecular chaperone: the role of the pro-peptide in protein folding. *Enzyme*, **45**:314-321.
- Ip, N. Y., Ibanez, C. F., Nye, S. H., McClain, J., Jones, P. F., Gies, D. R., Belluscio, L., Le Beau, M. M., Espinosa, R., III, Squinto, S. P., Persson, H., and Yancopoulos, G. D. (1992). Mammalian neurotrophin-4: structure, chromosomal localization, tissue distribution, and receptor specificity. *Proc Natl Acad Sci USA*, **89**:3060-3064.
- Isackson, P. J. and Bradshaw, R. A. (1984). The alpha-subunit of mouse 7 S nerve growth factor is an inactive serine protease. *J. Biol. Chem.*, **259**:5380-5383.

- Jain, S. C., Shinde, U., Li, Y., Inouye, M., and Berman, H. M. (1998). The crystal structure of an autoprocessed Ser221Cys-subtilisin E-propeptide complex at 2.0 Å resolution. *J. Mol. Biol.*, **284**:137-144.
- Kahle, P., Burton, L. E., Schmelzer, C. H., and Hertel, C. (1992). The amino terminus of nerve growth factor is involved in the interaction with the receptor tyrosine kinase p140trkA. *J. Biol. Chem.*, **267**:22707-22710.
- Kaplan, D. R. and Miller, F. D. (1997). Signal transduction by the neurotrophin receptors. *Curr. Opin. Cell Biol.*, **9**:213-221.
- Kaplan, D. R. and Miller, F. D. (2000). Neurotrophin signal transduction in the nervous system. *Curr. Opin. Neurobiol.*, **10**:381-391.
- Klein, R., Jing, S. Q., Nanduri, V., O'Rourke, E., and Barbacid, M. (1991a). The trk proto-oncogene encodes a receptor for nerve growth factor. *Cell*, **65**:189-197.
- Klein, R., Nanduri, V., Jing, S. A., Lamballe, F., Tapley, P., Bryant, S., Cordon-Cardo, C., Jones, K. R., Reichardt, L. F., and Barbacid, M. (1991b). The trkB tyrosine protein kinase is a receptor for brain-derived neurotrophic factor and neurotrophin-3. *Cell*, **66**:395-403.
- Kliemann, M. (2001). Untersuchung zur Funktion der beiden C-terminalen Aminosäuren des NGF-Vorläuferproteins bei der in vitro Strukturbildung des Wachstumsfaktors. *Diplomarbeit* Martin-Luther-Universität Halle-Wittenberg.
- Kliemann, M., Rattenholl, A., Golbik, R., Balbach, J., Lilie, H., Rudolph, R., and Schwarz, E. (2004). The mature part of proNGF induces the structure of its pro-peptide. *FEBS Lett.*, **566**:207-212.
- Kullander, K., Kaplan, D., and Ebendal, T. (1997). Two restricted sites on the surface of the nerve growth factor molecule independently determine specific TrkA receptor binding and activation. *J Biol Chem*, **272**:9300-9307.
- LaLonde, J. M., Zhao, B., Janson, C. A., D'Alessio, K. J., McQueney, M. S., Orsini, M. J., Debouck, C. M., and Smith, W. W. (1999). The crystal structure of human procathepsin K. *Biochemistry*, **38**:862-869.
- Lamballe, F., Klein, R., and Barbacid, M. (1991). trkC, a new member of the trk family of tyrosine protein kinases, is a receptor for neurotrophin-3. *Cell*, **66**:967-979.
- Large, T. H., Bodary, S. C., Clegg, D. O., Weskamp, G., Otten, U., and Reichardt, L. F. (1986). Nerve growth factor gene expression in the developing rat brain. *Science*, **234**:352-355.
- Larsen, T. A., Olson, A. J., and Goodsell, D. S. (1998). Morphology of protein-protein interfaces. *Structure*, **6**:421-427.
- Le-Niculescu, H., Bonfoco, E., Kasuya, Y., Claret, F. X., Green, D. R., and Karin, M. (1999). Withdrawal of survival factors results in activation of the JNK pathway in neuronal cells leading to Fas ligand induction and cell death. *Mol. Cell Biol.*, **19**:751-763.

- Lee, R., Kermani, P., Teng, K. K., and Hempstead, B. L. (2001). Regulation of cell survival by secreted proneurotrophins. *Science*, **294**:1945-1948.
- Leibrock, J., Lottspeich, F., Hohn, A., Hofer, M., Hengerer, B., Masiakowski, P., Thoenen, H., and Barde, Y. A. (1989). Molecular cloning and expression of brain-derived neurotrophic factor. *Nature*, **341**:149-152.
- Levi-Montalcini, R. (1987). The nerve growth factor 35 years later. *Science*, **237**:1154-1162.
- Linggi, M. S., Burke, T. L., Williams, B. B., Harrington, A., Kraemer, R., Hempstead, B. L., Yoon, S. O., and Carter, B. D. (2005). Neurotrophin receptor interacting factor (NRIF) is an essential mediator of apoptotic signaling by the p75 neurotrophin receptor. *J. Biol. Chem.*, **280**:13801-13808.
- Lobos, E., Gebhardt, C., Kluge, A., and Spanel-Borowski, K. (2005). Expression of nerve growth factor isoforms in the rat uterus during pregnancy: accumulation of precursor proNGF. *Endocrinology*
- Lonze, B. E. and Ginty, D. D. (2002). Function and regulation of CREB family transcription factors in the nervous system. *Neuron*, **35**:605-623.
- Mahadeo, D., Kaplan, L., Chao, M. V., and Hempstead, B. L. (1994). High affinity nerve growth factor binding displays a faster rate of association than p140trk binding. Implications for multi-subunit polypeptide receptors. *J. Biol. Chem.*, **269**:6884-6891.
- Maisonpierre, P. C., Belluscio, L., Squinto, S., Ip, N. Y., Furth, M. E., Lindsay, R. M., and Yancopoulos, G. D. (1990). Neurotrophin-3: a neurotrophic factor related to NGF and BDNF. *Science*, **247**:1446-1451.
- Majdan, M., Walsh, G. S., Aloyz, R., and Miller, F. D. (2001). TrkA mediates developmental sympathetic neuron survival in vivo by silencing an ongoing p75NTR-mediated death signal. *J. Cell Biol.*, **155**:1275-1285.
- Mason, A. J., Evans, B. A., Cox, D. R., Shine, J., and Richards, R. I. (1983). Structure of mouse kallikrein gene family suggests a role in specific processing of biologically active peptides. *Nature*, **303**:300-307.
- McDonald, N. Q. and Hendrickson, W. A. (1993). A structural superfamily of growth factors containing a cystine knot motif. *Cell*, **73**:421-424.
- McDonald, N. Q., Lapatto, R., Murray-Rust, J., Gunning, J., Wlodawer, A., and Blundell, T. L. (1991). New protein fold revealed by a 2.3-Å resolution crystal structure of nerve growth factor. *Nature*, **354**:411-414.
- Miller, F. D. and Kaplan, D. R. (2001). Neurotrophin signalling pathways regulating neuronal apoptosis. *Cell Mol. Life Sci.*, **58**:1045-1053.
- Mobley, W. C., Schenker, A., and Shooter, E. M. (1976). Characterization and isolation of proteolytically modified nerve growth factor. *Biochemistry*, **15**:5543-5552.
- Moore, J. B., Jr., Mobley, W. C., and Shooter, E. M. (1974). Proteolytic modification of the beta nerve growth factor protein. *Biochemistry*, **13**:833-840.

- Mowla, S. J., Pareek, S., Farhadi, H. F., Petrecca, K., Fawcett, J. P., Seidah, N. G., Morris, S. J., Sossin, W. S., and Murphy, R. A. (1999). Differential sorting of nerve growth factor and brain-derived neurotrophic factor in hippocampal neurons. *J. Neurosci.*, **19**:2069-2080.
- Murphy, R. A., Chlumecky, V., Smillie, L. B., Carpenter, M., Nattriss, M., Anderson, J. K., Rhodes, J. A., Barker, P. A., Siminoski, K., Campenot, R. B., and Haskins, J. (1989). Isolation and characterization of a glycosylated form of beta nerve growth factor in mouse submandibular glands. *J. Biol. Chem.*, **264**:12502-12509.
- Narhi, L. O., Rosenfeld, R., Talvenheimo, J., Prestrelski, S. J., Arakawa, T., Lary, J. W., Kolvenbach, C. G., Hecht, R., Boone, T., Miller, J. A., and Yphantis, D. A. (1993). Comparison of the biophysical characteristics of human brain-derived neurotrophic factor, neurotrophin-3, and nerve growth factor. *J. Biol. Chem.*, **268**:13309-13317.
- Neri, D., Billeter, M., Wider, G., and Wuthrich, K. (1992). NMR determination of residual structure in a urea-denatured protein, the 434-repressor. *Science*, **257**:1559-1563.
- Nilsson, A. S., Fainzilber, M., Falck, P., and Ibanez, C. F. (1998). Neurotrophin-7: a novel member of the neurotrophin family from the zebrafish. *FEBS Lett.*, **424**:285-290.
- Nykjaer, A., Lee, R., Teng, K. K., Jansen, P., Madsen, P., Nielsen, M. S., Jacobsen, C., Kliemannel, M., Schwarz, E., Willnow, T. E., Hempstead, B. L., and Petersen, C. M. (2004). Sortilin is essential for proNGF-induced neuronal cell death. *Nature*, **427**:843-848.
- O'Connell, L., Hongo, J. A., Presta, L. G., and Tsoulfas, P. (2000). TrkA amino acids controlling specificity for nerve growth factor. *J. Biol. Chem.*, **275**:7870-7877.
- Oefner, C., D'Arcy, A., Winkler, F. K., Eggimann, B., and Hosang, M. (1992). Crystal structure of human platelet-derived growth factor BB. *EMBO J.*, **11**:3921-3926.
- Ohta, Y., Hojo, H., Aimoto, S., Kobayashi, T., Zhu, X., Jordan, F., and Inouye, M. (1991). Pro-peptide as an intramolecular chaperone: renaturation of denatured subtilisin E with a synthetic pro-peptide [corrected]. *Mol. Microbiol.*, **5**:1507-1510.
- Pallaghy, P. K., Nielsen, K. J., Craik, D. J., and Norton, R. S. (1994). A common structural motif incorporating a cystine knot and a triple-stranded beta-sheet in toxic and inhibitory polypeptides. *Protein Sci.*, **3**:1833-1839.
- Pantazis, N. J. (1983). Nerve growth factor synthesized by mouse fibroblast cells in culture: absence of alpha and gamma subunits. *Biochemistry*, **22**:4264-4271.
- Petersen, C. M., Nielsen, M. S., Nykjaer, A., Jacobsen, L., Tommerup, N., Rasmussen, H. H., Roigaard, H., Gliemann, J., Madsen, P., and Moestrup, S. K. (1997). Molecular identification of a novel candidate sorting receptor purified from human brain by receptor-associated protein affinity chromatography. *J. Biol. Chem.*, **272**:3599-3605.
- Philo, J., Talvenheimo, J., Wen, J., Rosenfeld, R., Welcher, A., and Arakawa, T. (1994). Interactions of Neurotrophin-3 (Nt-3), Brain-Derived Neurotrophic Factor (Bdnf), and the Nt-3 Bdnf Heterodimer with the Extracellular Domains of the Trkb and Trkc Receptors. *J. Biol. Chem.*, **269**:27840-27846.

- Philo, J. S., Rosenfeld, R., Arakawa, T., Wen, J., and Narhi, L. O. (1993). Refolding of Brain-Derived Neurotrophic Factor from Guanidine-Hydrochloride - Kinetic Trapping in A Collapsed Form Which Is Incompetent for Dimerization. *Biochemistry*, **32**:10812-10818.
- Podobnik, M., Kuhelj, R., Turk, V., and Turk, D. (1997). Crystal structure of the wild-type human procathepsin B at 2.5 Å resolution reveals the native active site of a papain-like cysteine protease zymogen. *J. Mol. Biol.*, **271**:774-788.
- Price-Carter, M., Gray, W. R., and Goldenberg, D. P. (1996a). Folding of omega-conotoxins. 1. Efficient disulfide-coupled folding of mature sequences in vitro. *Biochemistry*, **35**:15537-15546.
- Price-Carter, M., Gray, W. R., and Goldenberg, D. P. (1996b). Folding of omega-conotoxins. 2. Influence of precursor sequences and protein disulfide isomerase. *Biochemistry*, **35**:15547-15557.
- Radziejewski, C., Robinson, R. C., DiStefano, P. S., and Taylor, J. W. (1992). Dimeric structure and conformational stability of brain-derived neurotrophic factor and neurotrophin-3. *Biochemistry*, **31**:4431-4436.
- Ramer, M. S., Priestley, J. V., and McMahon, S. B. (2000). Functional regeneration of sensory axons into the adult spinal cord. *Nature*, **403**:312-316.
- Rattenholl, A. (2001). Untersuchung zur Pro-Sequenz-vermittelten Faltung von rekombinanten, humanen Nervenwachstumsfaktor. *Dissertation* Martin-Luther-Universität Halle-Wittenberg.
- Rattenholl, A., Lilie, H., Grossmann, A., Stern, A., Schwarz, E., and Rudolph, R. (2001a). The pro-sequence facilitates folding of human nerve growth factor from *Escherichia coli* inclusion bodies. *Eur. J. Biochem.*, **268**:3296-3303.
- Rattenholl, A., Ruoppolo, M., Flagiello, A., Monti, M., Vinci, F., Marino, G., Lilie, H., Schwarz, E., and Rudolph, R. (2001b). Pro-sequence assisted folding and disulfide bond formation of human nerve growth factor. *J. Mol. Biol.*, **305**:523-533.
- Reinshagen, M., Geerling, I., Eysselein, V. E., Adler, G., Huff, K. R., Moore, G. P., and Lakshmanan, J. (2000). Commercial recombinant human beta-nerve growth factor and adult rat dorsal root ganglia contain an identical molecular species of nerve growth factor prohormone. *J. Neurochem.*, **74**:2127-2133.
- Rosenthal, A., Goeddel, D. V., Nguyen, T., Lewis, M., Shih, A., Laramee, G. R., Nikolics, K., and Winslow, J. W. (1990). Primary structure and biological activity of a novel human neurotrophic factor. *Neuron*, **4**:767-773.
- Roux, P. P. and Barker, P. A. (2002). Neurotrophin signaling through the p75 neurotrophin receptor. *Prog. Neurobiol.*, **67**:203-233.
- Rudolph, R. and Lilie, H. (1996). In vitro folding of inclusion body proteins. *FASEB J.*, **10**:49-56.

- Ruvinov, S., Wang, L., Ruan, B., Almog, O., Gilliland, G. L., Eisenstein, E., and Bryan, P. N. (1997). Engineering the independent folding of the subtilisin BPN' prodomain: analysis of two-state folding versus protein stability. *Biochemistry*, **36**:10414-10421.
- Ryden, M., Hempstead, B., and Ibanez, C. F. (1997). Differential modulation of neuron survival during development by nerve growth factor binding to the p75 neurotrophin receptor. *J. Biol. Chem.*, **272**:16322-16328.
- Saab-Rincon, G., Gualfetti, P. J., and Matthews, C. R. (1996). Mutagenic and thermodynamic analyses of residual structure in the alpha subunit of tryptophan synthase. *Biochemistry*, **35**:1988-1994.
- Schlunegger, M. P. and Grutter, M. G. (1992). An unusual feature revealed by the crystal structure at 2.2 Å resolution of human transforming growth factor-beta 2. *Nature*, **358**:430-434.
- Schmid, F. X. and Baldwin, R. L. (1979). The rate of interconversion between the two unfolded forms of ribonuclease A does not depend on guanidinium chloride concentration. *J. Mol. Biol.*, **133**:285-287.
- Seidah, N. G., Benjannet, S., Pareek, S., Savaria, D., Hamelin, J., Goulet, B., Laliberte, J., Lazure, C., Chretien, M., and Murphy, R. A. (1996). Cellular processing of the nerve growth factor precursor by the mammalian pro-protein convertases. *Biochem. J.*, **314** (Pt 3):951-960.
- Selby, M. J., Edwards, R., Sharp, F., and Rutter, W. J. (1987). Mouse nerve growth factor gene: structure and expression. *Mol. Cell Biol.*, **7**:3057-3064.
- Shinde, U. and Inouye, M. (2000). Intramolecular chaperones: polypeptide extensions that modulate protein folding. *Semin. Cell Dev. Biol.*, **11**:35-44.
- Shinde, U., Li, Y., Chatterjee, S., and Inouye, M. (1993). Folding pathway mediated by an intramolecular chaperone. *Proc. Natl. Acad. Sci. USA*, **90**:6924-6928.
- Silen, J. L. and Agard, D. A. (1989). The alpha-lytic protease pro-region does not require a physical linkage to activate the protease domain in vivo. *Nature*, **341**:462-464.
- Silverman, R. E. and Bradshaw, R. A. (1982). Nerve growth factor: subunit interactions in the mouse submaxillary gland 7S complex. *J. Neurosci. Res.*, **8**:127-136.
- Sorenson, P., Winther, J. R., Kaarsholm, N. C., and Poulsen, F. M. (1993). The pro region required for folding of carboxypeptidase Y is a partially folded domain with little regular structural core. *Biochemistry*, **32**:12160-12166.
- Steiner, D. F. and Clark, J. L. (1968). The spontaneous reoxidation of reduced beef and rat proinsulins. *Proc. Natl. Acad. Sci. USA*, **60**:622-629.
- Taylor, J. M., Cohen, S., and Mitchell, W. M. (1970). Epidermal growth factor: high and low molecular weight forms. *Proc. Natl. Acad. Sci. USA*, **67**:164-171.
- Teng, H. K., Teng, K. K., Lee, R., Wright, S., Tevar, S., Almeida, R. D., Kermani, P., Torkin, R., Chen, Z. Y., Lee, F. S., Kraemer, R. T., Nykjaer, A., and Hempstead, B. L. (2005).

- ProBDNF induces neuronal apoptosis via activation of a receptor complex of p75^{NTR} and sortilin. *J. Neurosci.*, **25**:5455-5463.
- Timm, D. E., de Haset, P. L., and Neet, K. E. (1994). Comparative equilibrium denaturation studies of the neurotrophins: nerve growth factor, brain-derived neurotrophic factor, neurotrophin 3, and neurotrophin 4/5. *Biochemistry*, **33**:4667-4676.
- Timm, D. E. and Neet, K. E. (1992). Equilibrium denaturation studies of mouse β -nerve growth factor. *Protein Sci.*, **1**:236-244.
- Tuszynski, M. H. (2002). Growth-factor gene therapy for neurodegenerative disorders. *Lancet Neurol.*, **1**:51-57.
- Tuszynski, M. H., Thal, L., Pay, M., Salmon, D. P., HS, U., Bakay, R., Patel, P., Blesch, A., Vahlsing, H. L., Ho, G., Tong, G., Potkin, S. G., Fallon, J., Hansen, L., Mufson, E. J., Kordower, J. H., Gall, C., and Conner, J. (2005). A phase 1 clinical trial of nerve growth factor gene therapy for Alzheimer disease. *Nat. Med.*, **11**:551-555.
- Ullrich, A., Gray, A., Berman, C., and Dull, T. J. (1983). Human β -nerve growth factor gene sequence highly homologous to that of mouse. *Nature*, **303**:821-825.
- Urfer, R., Tsoulfas, P., Soppet, D., Escandon, E., Parada, L. F., and Presta, L. G. (1994). The binding epitopes of neurotrophin-3 to its receptors trkC and gp75 and the design of a multifunctional human neurotrophin. *EMBO J.*, **13**:5896-5909.
- Varon, S., Nomura, J., and Shooter, E. M. (1967). Subunit structure of a high-molecular-weight form of the nerve growth factor from mouse submaxillary gland. *Proc. Natl. Acad. Sci. USA*, **57**:1782-1789.
- Varon, S., Normura, J., and Shooter, E. M. (1968). Reversible dissociation of the mouse nerve growth factor protein into different subunits. *Biochemistry*, **7**:1296-1303.
- Voorberg, J., Fontijn, R., van Mourik, J. A., and Pannekoek, H. (1990). Domains involved in multimer assembly of von willebrand factor (vWF): multimerization is independent of dimerization. *EMBO J.*, **9**:797-803.
- Weissman, J. S. and Kim, P. S. (1992). The pro region of BPTI facilitates folding. *Cell*, **71**:841-851.
- Wentzel, A., Christmann, A., Kratzner, R., and Kolmar, H. (1999). Sequence requirements of the GPNG beta-turn of the Ecballium elaterium trypsin inhibitor II explored by combinatorial library screening. *J. Biol. Chem.*, **274**:21037-21043.
- Whittemore, S. R., Ebendal, T., Larkfors, L., Olson, L., Seiger, A., Stromberg, I., and Persson, H. (1986). Development and regional expression of beta nerve growth factor messenger RNA and protein in the rat central nervous system. *Proc. Natl. Acad. Sci. USA*, **83**:817-821.
- Wiesmann, C., Ultsch, M. H., Bass, S. H., and de Vos, A. M. (1999). Crystal structure of nerve growth factor in complex with the ligand-binding domain of the TrkA receptor. *Nature*, **401**:184-188.

- Winther, J. R. and Sorensen, P. (1991). Propeptide of carboxypeptidase Y provides a chaperone-like function as well as inhibition of the enzymatic activity. *Proc. Natl. Acad. Sci. USA*, **88**:9330-9334.
- Winther, J. R., Sorensen, P., and Kielland-Brandt, M. C. (1994). Refolding of a carboxypeptidase Y folding intermediate in vitro by low-affinity binding of the proregion. *J. Biol. Chem.*, **269**:22007-22013.
- Wise, R. J., Pittman, D. D., Handin, R. I., Kaufman, R. J., and Orkin, S. H. (1988). The propeptide of von Willebrand factor independently mediates the assembly of von Willebrand multimers. *Cell*, **52**:229-236.
- Woo, S. B. and Neet, K. E. (1996). Characterization of histidine residues essential for receptor binding and activity of nerve growth factor. *J. Biol. Chem.*, **271**:24433-24441.
- Woo, S. B., Timm, D. E., and Neet, K. E. (1995). Alteration of NH₂-terminal residues of nerve growth factor affects activity and Trk binding without affecting stability or conformation. *J. Biol. Chem.*, **270**:6278-6285.
- Yamashita, T., Fujitani, M., Hata, K., Mimura, F., and Yamagishi, S. (2005). Diverse functions of the p75 neurotrophin receptor. *Anat. Sci. Int.*, **80**:37-41.
- Yardley, G., Relf, B., Lakshmanan, J., Reinshagen, M., and Moore, G. P. (2000). Expression of nerve growth factor mRNA and its translation products in the anagen hair follicle. *Exp. Dermatol.*, **9**:283-289.
- Yoon, S. O., Casaccia-Bonofil, P., Carter, B., and Chao, M. V. (1998). Competitive signaling between TrkA and p75 nerve growth factor receptors determines cell survival. *J. Neurosci.*, **18**:3273-3281.
- Zhang, Z. and Smith, D. L. (1993). Determination of amide hydrogen exchange by mass spectrometry: a new tool for protein structure elucidation. *Protein Sci.*, **2**:522-531.

Abkürzungen

Å	Angström
CD	Cirkulardichroismus
Cys	Cystein
Da	Dalton
<i>E. coli</i>	<i>Escherichia coli</i>
GdnHCl/GdmCl	Guanidin Hydrochlorid
GSH	reduziertes Glutathion
GSSG	oxidiertes Glutathion
HSQC	<i>heteronuclear single-quantum correlation</i>
IB	<i>inclusion body</i>
IPTG	Isopropyl- β -thiogalactopyranosid
K_D	Gleichgewichts-Dissoziationskonstante
M	Molar
MALDI-TOF	<i>matrix assisted laser desorption ionization-time of flight</i>
NMR	<i>nuclear magnetic resonance</i>
NGF	Nervenwachstumsfaktor; <i>nerve growth factor</i>
NT	Neurotrophin
pdb	Proteindatenbank (www.pdb.org)
RP-HPLC	<i>reverse phase-HPLC</i>
Trk	Tyrosinkinase
UV	Ultraviolett