

7. Literaturverzeichnis

- Ali, S. und Taylor, W.C. (2001a) The 3' non-coding region of a C4 photosynthesis gene increases transgene expression when combined with heterologous promoters. *Plant Mol Biol* (46) 325-34.
- Ali, S. und Taylor, W.C. (2001b) Quantitative regulation of the Flaveria Me1 gene is controlled by the 3'-untranslated region and sequences near the amino terminus. *Plant Mol Biol* (46) 251-61.
- Allen, R.S., Millgate, A.G., Chitty, J.A., Thisleton, J., Miller, J.A., Fist, A.J., Gerlach, W.L. und Larkin, P.J. (2004) RNAi-mediated replacement of morphine with the nonnarcotic alkaloid reticuline in opium poppy. *Nat Biotechnol* (22) 1559-66.
- Bauer, W. und Zenk, M.H. (1989) Formation of both methylenedioxy groups in the alkaloid (*S*)-stylophine is catalyzed by cytochrome P-450 enzymes. *Tetrah Lett* (30) 5257-60.
- Bauer, W. und Zenk, M.H. (1991) Two methylenedioxy bridge forming cytochrome P-450 dependent enzymes are involved in (*S*)-stylophine biosynthesis. *Phytochemistry* (30) 2953-61.
- Bernáth, J. und Tetenyi, P. (1982) Production characteristics of *Papaver somniferum* L. cultivars of different origin and vegetation cycles. *Bull Narc* (34) 113-27.
- Beyer, H. und Walter, W. (1991) *Lehrbuch der organischen Chemie*. S. Hirzel Verlag, Stuttgart, 22. Auflage.
- Bird, D.A., Franceschi, V.R. und Facchini, P.J. (2003) A tale of three cell types: alkaloid biosynthesis is localized to sieve elements in opium poppy. *Plant Cell* (15) 2626-35.
- Bradford, M.M. (1976) A rapid and sensitive method for the quantitation of protein utilizing the protein-dye binding. *Anal Biochem* (72) 248-54.
- Burbulis, I.E. und Winkel-Shirley, B. (1999) Interactions among enzymes of the Arabidopsis flavonoid biosynthetic pathway. *PNAS USA* (96) 12929-34.
- Chitty, J.A., Allen, R.S., Fist, A.J. und Larkin, P.J. (2003) Genetic transformation in commercial Tasmanian cultivars of opium poppy, *Papaver somniferum*, and movement of transgenic pollen in the field. *Func Plant Biol* (30) 1045-58.
- Choi, K.B., Morishige, T. und Sato, F. (2001) Purification and characterization of coclaurine *N*-methyltransferase from cultured *Coptis japonica* cells. *Phytochemistry* (56) 649-55.
- Choi, K.B., Morishige, T., Shitan, N., Yazaki, K. und Sato, F. (2002) Molecular cloning and characterization of coclaurine *N*-methyltransferase from cultured cells of *Coptis japonica*. *J Biol Chem* (277) 830-5.
- Decker, G., Wanner, G., Zenk, M.H. und Lottspeich, F. (2000) Characterization of proteins in latex of the opium poppy (*Papaver somniferum*) using two-dimensional gel electrophoresis and microsequencing. *Electrophoresis* (21) 3500-16.
- De-Eknamkul, W. und Zenk, M.H. (1992) Purification and properties of 1,2-dehydroreticuline reductase from *Papaver somniferum* seedlings. *Phytochemistry* (31) 813-21.
- Dittrich, H. und Kutchan, T.M. (1991) Molecular cloning, expression, and induction of berberine bridge enzyme, an enzyme essential to the formation of benzophenanthridine alkaloids in the response of plants to pathogenic attack. *PNAS USA* (88) 9969-73.
- Facchini, P.J. und De Luca, V. (1994) Differential and tissue-specific expression of a gene family for tyrosine/dopa decarboxylase in opium poppy. *J Biol Chem* (269) 26684-90.
- Facchini, P.J. und De Luca, V. (1995) Phloem-Specific Expression of Tyrosine/Dopa Decarboxylase Genes and the Biosynthesis of Isoquinoline Alkaloids in Opium Poppy. *Plant Cell* (07) 1811-21.
- Facchini, P.J., Penzes, C., Johnson, A.G. und Bull, D. (1996) Molecular characterization of berberine bridge enzyme genes from opium poppy. *Plant Physiol* (112) 1669-77.
- Facchini, P.J. (2001) ALKALOID BIOSYNTHESIS IN PLANTS: Biochemistry, Cell Biology, Molecular Regulation, and Metabolic Engineering Applications. *Annu Rev Plant Physiol Plant Mol Biol* (52) 29-66.
- Facchini, P.J. und Park, S.U. (2003) Developmental and inducible accumulation of gene transcripts involved in alkaloid biosynthesis in opium poppy. *Phytochemistry* (64) 177-86.
- Frenzel, T. und Zenk, M.H. (1990a) Purification and characterization of three isoforms of *S*-adenosyl-L-methionine: (*R,S*)-tetrahydrobenzylisoquinoline-*N*-methyltransferase from *Berberis koetianeana* cell cultures. *Phytochemistry* (29) 3491-97.
- Frenzel, T. und Zenk, M.H. (1990b) *S*-adenosyl-L-methionine: 3'-hydroxy-*N*-methyl-(*S*)-coclaurine-4'-*O*-methyl transferase, a regio- and stereoselective enzyme of the (*S*)-reticuline pathway. *Phytochemistry* (29) 3505-11.
- Frick, S. und Kutchan, T.M. (1999) Molecular cloning and functional expression of *O*-methyltransferases

- common to isoquinoline alkaloid and phenylpropanoid biosynthesis. *Plant J* (17) 329-39.
- Frick, S., Chitty, J.A., Kramell, R., Schmidt, J., Allen, R.S., Larkin, P.J. und Kutchan, T.M. (2004) Transformation of opium poppy (*Papaver somniferum* L.) with antisense berberine bridge enzyme gene (*anti-bbe*) via somatic embryogenesis results in an altered ratio of alkaloids in latex but not in roots. *Transgenic Res* (13) 607-13.
- Frick, S., Kramell, R., Larkin, P.J. und Kutchan, T.M. (2005a) Studying morphine biosynthesis using transgenic opium poppy (*Papaver somniferum* L.). *Acta Horticulturae* (680) 37-43.
- Frick, S., Kramell, R., Schmidt, J., Fist, A.J. und Kutchan, T.M. (2005b) Comparative qualitative and quantitative determination of alkaloids in narcotic and condiment *Papaver somniferum* cultivars. *J Nat Prod* (68) 666-73.
- Frick, S., Kramell, R. und Kutchan, T.M. (im Druck) Metabolic engineering of a morphine biosynthetic P450 in opium poppy surpasses breeding. *Metabolic Engineering*.
- Garfinkel, D.J. und Nester, E.W. (1980) *Agrobacterium tumefaciens* mutants affected in crown gall tumorigenesis and octopine catabolism. *J Bacteriol* (144) 732-43.
- Gerardy, R. und Zenk, M.H. (1992) Formation of salutaridine from (*R*)-reticuline by a membrane-bound cytochrome P-450 enzyme from *Papaver somniferum*. *Phytochemistry* (32) 79-86.
- Gerardy, R. und Zenk, M.H. (1993) Purification and characterization of salutaridine: NADPH 7-oxidoreductase from *Papaver somniferum*. *Phytochemistry* (34) 125-32.
- Gleave, A.P. (1992) A versatile binary vector system with a T-DNA organisational structure conducive to efficient integration of cloned DNA into the plant genome. *Plant Mol Biol* (20) 1203-7.
- Gollwitzer, J., Lenz, R., Hampp, N. und Zenk, M.H. (1993) The transformation of neopinone to codeinone in morphine biosynthesis proceeds non-enzymatically. *Tetrah Lett* (34) 5703-06.
- Grothe, T., Lenz, R. und Kutchan, T.M. (2001) Molecular characterization of the salutaridinol 7-*O*-acetyltransferase involved in morphine biosynthesis in opium poppy *Papaver somniferum*. *J Biol Chem* (276) 30717-23.
- Haas, B., Volfovsky, N., Town, C., Troukhan, M., Alexandrov, N., Feldmann, K., Flavell, R., White, O. und Salzberg, S. (2002) Full-length messenger RNA sequences greatly improve genome annotation. *Genome Biol* (03) RESEARCH0029. Epub 2002 May 30.
- Hammer, K. und Fritsch, R. (1977) Zur Frage nach der Ursprungsart des Kulturmahns (*Papaver somniferum* L.). *Kulturpflanze* (25) 113-24.
- Hanahan, D. (1983) Studies on transformation of *Escherichia coli* with plasmids. *J Mol Biol* (166) 557-80.
- Hanahan, D., Jessee, J. und Bloom, F.R. (1991) Plasmid transformation of *Escherichia coli* and other bacteria. *Methods Enzymol* (204) 63-113.
- Hellens, R., Mullineaux, P. und Klee, H. (2000) Technical Focus: a guide to *Agrobacterium* binary Ti vectors. *Trends Plant Sci* (05) 446-51.
- Hibi, N., Higashiguchi, S., Hashimoto, T. und Yamada, Y. (1994) Gene expression in tobacco low-nicotine mutants. *Plant Cell* (06) 723-35.
- Hileman, L.C., Drea, S., Martino, G., Litt, A. und Irish, V.F. (2005) Virus-induced gene silencing is an effective tool for assaying gene function in the basal eudicot species *Papaver somniferum* (opium poppy). *Plant J* (44) 334-41.
- Hirata, K., Poeaknapo, C., Schmidt, J. und Zenk, M.H. (2004) 1,2-Dehydroreticuline synthase, the branch point enzyme opening the morphinan biosynthetic pathway. *Phytochemistry* (65) 1039-46.
- Huang, F.C. und Kutchan, T.M. (2000) Distribution of morphinan and benzo[*c*]phenanthridine alkaloid gene transcript accumulation in *Papaver somniferum*. *Phytochemistry* (53) 555-64.
- Hutvagner, G. und Zamore, P.D. (2002) RNAi: nature abhors a double-strand. *Curr Opin Genet Dev* (12) 225-32.
- Jez, J.M., Ferrer, J.L., Bowman, M.E., Dixon, R.A. und Noel, J.P. (2000) Dissection of malonyl-coenzyme A decarboxylation from polyketide formation in the reaction mechanism of a plant polyketide synthase. *Biochemistry* (39) 890-902.
- Jorgensen, R.A., Cluster, P.D., English, J., Que, Q. und Napoli, C.A. (1996) Chalcone synthase cosuppression phenotypes in petunia flowers: comparison of sense vs. antisense constructs and single-copy vs. complex T-DNA sequences. *Plant Mol Biol* (31) 957-73.
- Joshi, C.P. und Chiang, V.L. (1998) Conserved sequence motifs in plant S-adenosyl-L-methionine-dependent methyltransferases. *Plant Mol Biol* (37) 663-74.
- Jurna, I. (2003) Sertürner und Morphin - eine historische Vignette. In *Der Schmerz*. Springer Verlag, Berlin/Heidelberg, Vol. 17, pp. 280-83.

- Kassem, M.A. und Jacquin, A. (2001) Somatic Embryogenesis, Rhizogenesis, and Morphinan Alkaloids Production in Two Species of Opium Poppy. *J Biomed Biotechnol* (01) 70-78.
- Kato, M., Mizuno, K., Crozier, A., Fujimura, T. und Ashihara, H. (2000) Caffeine synthase gene from tea leaves. *Nature* (406) 956-7.
- Kempe, K. (2003) Herstellung von Vektoren zur "RNA interference" für Gene der Benzylisochinolin-Biosynthese aus *Papaver somniferum* L. *Diplomarbeit, Technische Universität Bergakademie Freiberg*.
- Klahre, U., Crete, P., Leuenberger, S.A., Iglesias, V.A. und Meins, F., Jr. (2002) High molecular weight RNAs and small interfering RNAs induce systemic posttranscriptional gene silencing in plants. *PNAS USA* (99) 11981-6.
- Kodaira, H. und Spector, S. (1988) Transformation of thebaine to oripavine, codeine, and morphine by rat liver, kidney, and brain microsomes. *PNAS USA* (85) 1267-71.
- Kristensen, C., Morant, M., Olsen, C.E., Ekstrom, C.T., Galbraith, D.W., Moller, B.L. und Bak, S. (2005) Metabolic engineering of dhurrin in transgenic Arabidopsis plants with marginal inadvertent effects on the metabolome and transcriptome. *PNAS USA* (102) 1779-84.
- Ku, M.S., Agarie, S., Nomura, M., Fukayama, H., Tsuchida, H., Ono, K., Hirose, S., Toki, S., Miyao, M. und Matsuoka, M. (1999) High-level expression of maize phosphoenolpyruvate carboxylase in transgenic rice plants. *Nat Biotechnol* (17) 76-80.
- Kutchan, T.M. und Dittrich, H. (1995) Characterization and mechanism of the berberine bridge enzyme, a covalently flavinylated oxidase of benzophenanthridine alkaloid biosynthesis in plants. *J Biol Chem* (270) 24475-81.
- Kutchan, T.M. (1998) Molecular genetics of plant alkaloid biosynthesis. In Cordell, G.A. (ed.), *The Alkaloids*. Academic Press, San Diego, Vol. 50, pp. 257-316.
- Laemmli, U.K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* (227) 680-5.
- Larkin, P.J., Chitty, J.A. und Brettell, R.I.S. (1999) Methods for plant transformation and regeneration. International patent publication number WO 99/34663.
- Lazo, G.R., Stein, P.A. und Ludwig, R.A. (1991) A DNA transformation-competent Arabidopsis genomic library in Agrobacterium. *Biotechnology (NY)* 963-7.
- Le Flem-Bonhomme, V., Laurain-Mattar, D. und Fliniaux, M.A. (2004) Hairy root induction of *Papaver somniferum* var. album, a difficult-to-transform plant, by *A rhizogenes* LBA 9402. *Planta* (218) 890-3.
- Lee, J. und Pedersen, H. (2001) Stable genetic transformation of *Eschscholzia californica* expressing synthetic green fluorescent proteins. *Biotechnol Prog* (17) 247-51.
- Lenz, R. und Zenk, M.H. (1994) Closure of the oxide bridge in morphine biosynthesis. *Tetrah Lett* (35) 3897-900.
- Lenz, R. und Zenk, M.H. (1995a) Acetyl coenzyme A:salutaridinol-7-O-acetyltransferase from *Papaver somniferum* plant cell cultures. The enzyme catalyzing the formation of thebaine in morphine biosynthesis. *J Biol Chem* (270) 31091-6.
- Lenz, R. und Zenk, M.H. (1995b) Purification and properties of codeinone reductase (NADPH) from *Papaver somniferum* cell cultures and differentiated plants. *Eur J Biochem* (233) 132-9.
- Lenz, R. und Zenk, M.H. (1995c) Stereoselective reduction of codeinone, the penultimate enzymic step during morphine biosynthesis in *Papaver somniferum*. *Tetrah Lett* (36) 2449-52.
- Liscombe, D.K. und Facchini, P.J. (nicht veröffentlicht) Molecular cloning and characterization of S-adenosyl-L-methionine:(S)-tetrahydroprotoberberine-cis-N-methyltransferase from opium poppy. *unveröffentlicht*.
- Loeffler, S., Deus-Neumann, B. und Zenk, M.H. (1995) S-adenosyl-L-methionine:(S)-coclaurine-N-methyltransferase from *Tinospora cordifolia*. *Phytochemistry* (38) 1387-95.
- MacLeod, B.P. und Facchini, P.J. (2005) Methods for regeneration and transformation in *Eschscholzia californica* – a model plant to investigate alkaloid biosynthesis. In Vasquez-Flota, V.M.L.-V.a.F. (ed.), *Plant Cell Culture Protocols, Second Edition*. Humana Press, Totowa, NJ, USA.
- Mahmoud, S.S. und Croteau, R.B. (2001) Metabolic engineering of essential oil yield and composition in mint by altering expression of deoxyxylulose phosphate reductoisomerase and menthofuran synthase. *PNAS USA* (98) 8915-20.
- Miller, J.H. (1972) *Experiments in molecular genetics*. Cold Spring Harbor Press, Cold Spring Harbor, N.Y.

- Millgate, A.G., Pogson, B.J., Wilson, I.W., Kutchan, T.M., Zenk, M.H., Gerlach, W.L., Fist, A.J. und Larkin, P.J. (2004) Analgesia: morphine-pathway block in top1 poppies. *Nature* (431) 413-4.
- Morimoto, S., Suemori, K., Moriwaki, J., Taura, F., Tanaka, H., Aso, M., Tanaka, M., Suemune, H., Shimohigashi, Y. und Shoyama, Y. (2001) Morphine metabolism in the opium poppy and its possible physiological function. Biochemical characterization of the morphine metabolite, bismorphine. *J Biol Chem* (276) 38179-84.
- Morishige, T., Tsujita, T., Yamada, Y. und Sato, F. (2000) Molecular characterization of the S-adenosyl-L-methionine:3'-hydroxy-N-methylcoclaurine 4'-O-methyltransferase involved in isoquinoline alkaloid biosynthesis in *Coptis japonica*. *J Biol Chem* (275) 23398-405.
- Mullis, K.B. und Faloona, F.A. (1987) Specific synthesis of DNA in vitro via a polymerase-catalyzed chain reaction. *Methods Enzymol* (155) 335-50.
- Novak, B.H., Hudlicky, T., Reed, J.W., Mulzer, J. und Trauner, D. (2000) Morphine Synthesis and Biosynthesis: An Update. *Curr Org Chem* (04) 343-62.
- Ogita, S., Uefuji, H., Yamaguchi, Y., Koizumi, N. und Sano, H. (2003) Producing decaffeinated coffee plants. *Nature* (423) 823.
- O'Keefe, B.R. und Beecher, C. (1994) Isolation and Characterization of S-Adenosyl-L-Methionine:Tetrahydroberberine-cis-N-Methyltransferase from Suspension Cultures of *Sanguinaria canadensis* L. *Plant Physiol* (105) 395-403.
- Ounaron, A., Decker, G., Schmidt, J., Lottspeich, F. und Kutchan, T.M. (2003) (R,S)-Reticuline 7-O-methyltransferase and (R,S)-norcoclaurine 6-O-methyltransferase of *Papaver somniferum* - cDNA cloning and characterization of methyl transfer enzymes of alkaloid biosynthesis in opium poppy. *Plant J* (36) 808-19.
- Page, J.E. (2005) Silencing nature's narcotics: metabolic engineering of the opium poppy. *Trends Biotechnol* (23) 331-33.
- Panicot, M., Minguet, E., Ferrando, A., Alcazar, R., Blazquez, M., Carbonell, J., Altabella, T., Koncz, C. und Tiburcio, A. (2002) A polyamine metabolon involving aminopropyl transferase complexes in Arabidopsis. *Plant Cell* (14) 2539-51.
- Park, S.U., Johnson, A.G., Penzes-Yost, C. und Facchini, P.J. (1999) Analysis of promoters from tyrosine/dihydroxyphenylalanine decarboxylase and berberine bridge enzyme genes involved in benzyloquinoline alkaloid biosynthesis in opium poppy. *Plant Mol Biol* (40) 121-31.
- Park, S.U. und Facchini, P.J. (2000) Agrobacterium rhizogenes-mediated transformation of opium poppy, *Papaver somniferum* L., and California poppy, *Eschscholzia californica* Cham., root cultures. *J Exp Bot* (51) 1005-16.
- Park, S.U. und Facchini, P.J. (2001) Somatic embryogenesis from embryogenic cell suspension cultures of California poppy, *Eschscholzia californica* Cham. *In Vitro Cell Devel Biol - Plant* (37) 35-39.
- Park, S.U., Yu, M. und Facchini, P.J. (2002) Antisense RNA-mediated suppression of benzophenanthridine alkaloid biosynthesis in transgenic cell cultures of California poppy. *Plant Physiol* (128) 696-706.
- Park, S.U., Yu, M. und Facchini, P.J. (2003) Modulation of berberine bridge enzyme levels in transgenic root cultures of California poppy alters the accumulation of benzophenanthridine alkaloids. *Plant Mol Biol* (51) 153-64.
- Pauli, H.H. und Kutchan, T.M. (1998) Molecular cloning and functional heterologous expression of two alleles encoding (S)-N-methylcoclaurine 3'-hydroxylase (CYP80B1), a new methyl jasmonate-inducible cytochrome P-450-dependent mono-oxygenase of benzyloquinoline alkaloid biosynthesis. *Plant J* (13) 793-801.
- Rice, K.C. (1980) Synthetic opium alkaloids and derivatives. A short total synthesis of (+)-dihydrothebainone, (+)-dihydrocodeinone, and (+)-nordihydrocodeinone as an approach to a practical synthesis of morphine, codeine, and congeners. *J Org Chem* (45) 3135-37.
- Riechers, D.E. und Timko, M.P. (1999) Structure and expression of the gene family encoding putrescine N-methyltransferase in *Nicotiana tabacum*: new clues to the evolutionary origin of cultivated tobacco. *Plant Mol Biol* (41) 387-401.
- Rosco, A., Pauli, H.H., Priesner, W. und Kutchan, T.M. (1997) Cloning and heterologous expression of NADPH-cytochrome P450 reductases from the Papaveraceae. *Arch Biochem Biophys* (348) 369-77.
- Rueffer, M., El-Shagi, H., Nagakura, N. und Zenk, M.H. (1981) (S)-Norlaudanosoline Synthase: The first enzyme in the benzylochinoline biosynthetic pathway. *FEBS letters* (129) 5-9.

- Rueffer, M., Nagakura, N. und Zenk, M.H. (1983) Partial purification and properties of S-adenosylmethionine:(R), (S)-norlaudanolin-6-O-methyltransferase from *Argemone platyceras* cell cultures. *Planta Med* (49) 131–37.
- Rueffer, M. und Zenk, M.H. (1986) S-adenosyl-L-methionine: (S) -7,8,13, 14-tetrahydroberberine-N-methyltransferase, a branch point enzyme in the biosynthesis of benzophenanthridine and protopine alkaloids. *Tetrah Lett* (27) 5603-04.
- Rueffer, M., Zumstein, G. und Zenk, M.H. (1990) Partial purification of S-adenosyl-L-methionine:(S)-tetrahydroprotoberberine-cis N-methyltransferase from suspension cultured cells of *Eschscholtzia* and *Corydalis*. *Phytochemistry* (29) 3727-33.
- Sala, F., Manuela Rigano, M., Barbante, A., Basso, B., Walmsley, A.M. und Castiglione, S. (2003) Vaccine antigen production in transgenic plants: strategies, gene constructs and perspectives. *Vaccine* (21) 803-08.
- Samanani, N. und Facchini, P.J. (2002) Purification and characterization of norcoclaurine synthase. The first committed enzyme in benzyloquinoline alkaloid biosynthesis in plants. *J Biol Chem* (277) 33878-83.
- Samanani, N., Liscombe, D.K. und Facchini, P.J. (2004) Molecular cloning and characterization of norcoclaurine synthase, an enzyme catalyzing the first committed step in benzyloquinoline alkaloid biosynthesis. *Plant J* (40) 302-13.
- Samanani, N., Park, S.U. und Facchini, P.J. (2005) Cell type-specific localization of transcripts encoding nine consecutive enzymes involved in protoberberine alkaloid biosynthesis. *Plant Cell* (17) 915-26.
- Sambrook, J., Fritsch, E.F. und Maniatis, T. (1989) *Molecular cloning: a laboratory manual*. Cold Spring Harbor Press, Cold Spring Harbor, New York.
- Sanger, F., Nicklen, S. und Coulson, A.R. (1977) DNA sequencing with chain-terminating inhibitors. *PNAS USA* (74) 5463-7.
- Sato, F., Tsujita, T., Katagiri, Y., Yoshida, S. und Yamada, Y. (1994) Purification and characterization of S-adenosyl-L-methionine: norcoclaurine 6-O-methyltransferase from cultured *Coptis japonica* cells. *Eur J Biochem* (225) 125-31.
- Sato, F., Hashimoto, T., Hachiya, A., Tamura, K., Choi, K.B., Morishige, T., Fujimoto, H. und Yamada, Y. (2001) Metabolic engineering of plant alkaloid biosynthesis. *PNAS USA* (98) 367-72.
- Satyanarayana, K.V., Kumar, V., Chandrashekar, A. und Ravishankar, G.A. (2005) Isolation of promoter for N-methyltransferase gene associated with caffeine biosynthesis in *Coffea canephora*. *J Biotechnol* (119) 20-25.
- Schünmann, P.H.D., Llewellyn, D.J., Surin, B., Boevink, P., Feyter, R.C.D. und Waterhouse, P.M. (2003) A suite of novel promoters and terminators for plant biotechnology. *Func Plant Biol* (30) 443-52.
- Sertürner, F. (1806) Darstellung der reinen Mohnsäure (Opiumsäure) nebst einer Untersuchung des Opiums mit vorzüglicher Hinsicht auf einen darin neu entdeckten Stoff und die dahin gehörigen Bemerkungen. *J Pharm (Leipzig)* (14) 33–37.
- Southern, E.M. (1975) Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J Mol Biol* (98) 503-17.
- Steffens, P., Nagakura, N. und Zenk, M.H. (1985) Purification and characterization of the berberine bridge enzyme from *Berberis beaniana* cell cultures. *Phytochemistry* (24) 2577-83.
- Stenzel, O., Teuber, M. und Dräger, B. (2006) Putrescine N-methyltransferase in *Solanum tuberosum* L., a calystegine-forming plant. *Planta* (223) 200-12.
- Szabo, B., Lakatos, A., Koszegi, T., Katay, G. und L, B. (2005) Thin-layer chromatography-densitometry and liquid chromatography analysis of alkaloids in leaves of *Papaver somniferum* under stress conditions. *J AOAC Int* (88) 1571-7.
- Takeshita, N., Fujiwara, H., Mimura, H., Fitch, J.H., Yamada, Y. und Sato, F. (1995) Molecular cloning and characterization of S-adenosyl-L-methionine:scoulerine-9-O-methyltransferase from cultured cells of *Coptis japonica*. *Plant Cell Physiol* (36) 29-36.
- Tanahashi, T. und Zenk, M.H. (1990) New hydroxylated benzo[c]phenanthridine alkaloids from *Eschscholtzia californica* cell suspension cultures. *J Nat Prod* (53) 579-86.
- Uefuji, H., Ogita, S., Yamaguchi, Y., Koizumi, N. und Sano, H. (2003) Molecular cloning and functional characterization of three distinct N-methyltransferases involved in the caffeine biosynthetic pathway in coffee plants. *Plant Physiol* (132) 372-80.

- Unterlinner, B., Lenz, R. und Kutchan, T.M. (1999) Molecular cloning and functional expression of codeinone reductase: the penultimate enzyme in morphine biosynthesis in the opium poppy *Papaver somniferum*. *Plant J* (18) 465-75.
- Waterhouse, P.M., Graham, M.W. und Wang, M.B. (1998) Virus resistance and gene silencing in plants can be induced by simultaneous expression of sense and antisense RNA. *PNAS USA* (95) 13959-64.
- Weid, M., Ziegler, J. und Kutchan, T.M. (2004) The roles of latex and the vascular bundle in morphine biosynthesis in the opium poppy, *Papaver somniferum*. *PNAS USA* (101) 13957-62.
- Wesley, S.V., Helliwell, C.A., Smith, N.A., Wang, M.B., Rouse, D.T., Liu, Q., Gooding, P.S., Singh, S.P., Abbott, D., Stoutjesdijk, P.A., Robinson, S.P., Gleave, A.P., Green, A.G. und Waterhouse, P.M. (2001) Construct design for efficient, effective and high-throughput gene silencing in plants. *Plant J* (27) 581-90.
- Wesley, S.V., Helliwell, C., Wang, M.B. und Waterhouse, P. (2004) Posttranscriptional gene silencing in plants. *Methods Mol Biol* (265) 117-29.
- Winkel, B.S.J. (2004) METABOLIC CHANNELING IN PLANTS. *Ann Rev Pl Biol* (55) 85-107.
- Ye, X., Al-Babili, S., Klöti, A., Zhang, J., Lucca, P., Beyer, P. und Potrykus, I. (2000) Engineering the provitamin A (β -carotene) biosynthetic pathway into (carotenoid-free) rice endosperm. *Science* (287) 303-5.
- Zhang, L., Ding, R., Chai, Y., Bonfill, M., Moyano, E., Oksman-Caldentey, K.M., Xu, T., Pi, Y., Wang, Z., Zhang, H., Kai, G., Liao, Z., Sun, X. und Tang, K. (2004) Engineering tropane biosynthetic pathway in *Hyoscyamus niger* hairy root cultures. *PNAS USA* (101) 6786-91.
- Ziegler, J., Diaz-Chavez, M.L., Kramell, R., Ammer, C. und Kutchan, T.M. (2005) Comparative macroarray analysis of morphine containing *Papaver somniferum* and eight morphine free *Papaver* species identifies an *O*-methyltransferase involved in benzyloquinoline biosynthesis. *Planta* (222) 458-71.
- Ziegler, J., Voigtlander, S., Schmidt, J., Kramell, R., Miersch, O., Ammer, C., Gesell, A. und Kutchan, T.M. (2006) Comparative transcript and alkaloid profiling in *Papaver* species identifies a short chain dehydrogenase/reductase involved in morphine biosynthesis. *Plant J* (im Druck, Epub 13. September).
- Zubieta, C., Ross, J.R., Koscheski, P., Yang, Y., Pichersky, E. und Noel, J.P. (2003) Structural basis for substrate recognition in the salicylic acid carboxyl methyltransferase family. *Plant Cell* (15) 1704-16.