

## 6 References

- Adams, D.F.; Farwell, S.O.; Robinson, E.; Pack, M.R. and Bamesberger, W.L. (1981): Biogenic sulfur source strengths. *Environ. Sci. Technol.* 15 (12), 1493-1498.
- Aiyuk, S.; Amoako J.; Raskin, L.; Van Haandel A.; and Verstraete, W. (2004): Removal of carbon and nutrients from domestic wastewater using a low investment, integrated treatment concept. *Water Res.* 38 (13), 248-256.
- Akagi, J.M. (1995): Respiratory sulfate reduction in: (Barton, L.L., Ed.), *Sulfate-Reducing Bacteria*, pp. 889-111, Plenum Press, NY.
- Allam, A.I. and Hollis, J.P. (1972): Sulphide inhibition of oxidase in rice roots. *Phytopathology.* 62, 634-639.
- Allen, E.R. and Phatak, S. (1993): Control of organosulfur compound emissions using biofiltration: methyl mercaptan. In: *Proc. 86<sup>th</sup> Air and Waste Management Association Annual Meeting and Exhibition.* Denver, Colorado, USA.
- Al-Malack, M.H.; Anderson, G.K. and Almasi, A. (1998): Treatment of anoxic pond effluent using crossflow microfiltration. *Water Res.* 12 (32), 3738-37446.
- Annachhatre, A.P. and Suktrakoolvait, S. (2001): Biological sulfide oxidation in a fluidized bed reactor. *Environ. Technol.* 22 (6), 661-672.
- Armstrong, W. (1971): Radial oxygen losses from intact rice roots as affected by distance from the apex, respiration and water logging. *Physiologia Plantarum.* 25, 192-197.
- Armstrong, W. (1979): Aeration in higher plants. *Adv. Bot. Res.* 7, 226-332.
- Armstrong, J. and Armstrong, W. (1988): *Phragmites australis* – A preliminary study of soil oxidizing sites and internal gas transport pathways. *New Phytologist.* 108, 373-382.

- Armstrong, W.; Armstrong, J. and Beckett, P.M. (1990): Measurements and modelling of oxygen release from roots of *Phragmites australis*. In: Cooper, P.F., Findlater, B.C. (Eds.), *Constructed Wetlands in water Pollution Control*. Pergamon Press. Oxford, UK, pp. 41-51.
- Armstrong, J.; Armstrong, W. and Beckett, P.M. (1992): *Phragmites australis*: Venturi- and humidity-induced convections enhance rhizome aeration and rhizosphere oxidation. *New Phytol.* 120, 197-207.
- Armstrong, W.; Braendle, R. and Jackson, M.B. (1994): Mechanisms of flood tolerance in plants. *Acta Bot. Neerl.* 43, 307-358.
- Armstrong, J.; AlfreinZobayed, F. and Armstrong, W. (1996a): *Phragmites* die-back: Sulphide and acetic acid induced bud and root death, lignification, and blockage within aeration and vascular systems. *New Phytol.* 134 (4), 601-614.
- Armstrong, J.; Armstrong, W.; Armstrong, I.B. and Pittaway, G.R. (1996b): Senescence and phytotoxin, insect, fungal and mechanical damage: factors reducing convective gas-flow in *Phragmites australis*. *Aquat. Bot.* 54, 211-216.
- Armstrong, J.; Armstrong, W. and Van der Putten, W.H. (1996c): *Phragmites* die-back: bud and root death, blockages within the aeration and vascular systems and the possible role of phytotoxins. *New Phytol.* 133, 399-414.
- Armstrong, J. and Armstrong, W. (1999): *Phragmites* die-back: toxic effect of propionic, butyric and caproic acids in relation to pH. *New Phytol.* 142, 201-218.
- Asai, S.; Konishi, Y. And Yabu, T. (1990): Kinetics of absorption of hydrogen sulfide into aqueous ferric sulfate solutions. *AICHE J.* 36, 1331-1338.
- ASTM. (1994): D503089 Standard test method for density of soil and rock in place by the water replacement method in a test pit. American society for testing and materials, pp. 1006-1018.

- Baas Becking, L.G.M.; Kaplan, I.R. and Moore, D. (1960): Limits of the natural environment in terms of pH and oxidation-reduction potential. *J. Geol.* 68, 243-284.
- Bahlo, K.E. and Wach, F.G. (1990): Purification of domestic sewage with and without faeces by vertical intermittent filtration in reed and rush beds. In *Constructed wetlands in water pollution control* (ed. P.F. Cooper and B.C. Finlander), pp. 215-221. Oxford, U.K: Pergamon Press.
- Balck, G.R. and Hartge, K.H. (1986): Particle density. In Klute, A. (ed.) *Methods of soil analysis part 1. Physical and mineralogical methods*, 2<sup>nd</sup> ed. Pp. 377-382. Am. Society of Agronomy, Madison, WI.
- Balistrieri, L.S.; Brewer, P.G. and Murray, J.W. (1981): *Deep-Sea res.* 28A, 101-121.
- Barber, D.A. and Martin, J.K. (1976): The release of organic substances by cereal roots into soil. *New Phytol.* 76, 69-80.
- Barko, J.W., Gunnison, D. and Carpenter, S.R. (1991): Sediment interactions with submersed macrophytes growth and community dynamics. *Aqua.Bot.* 41, 41-65.
- Bart, D. and Hartman, J.M. (2000): Environmental determinants of *Phragmites australis* expansion in a New Jersey salt marsh: an experimental approach. *Oikos.* 89, 69-69.
- Batal, W.H.; Wildeman, T.R.; Laudon, L.S. and Mohdnoordin, N. (1989): Bacteriological tests from the constructed wetland of the Big Five Tunnel, Idaho Springs, Colorado. In: *Constructed Wetlands for Wastewater Treatment* (ed. D.A. Hammer), 550-557. Lewis Publishers.
- Batchelor, B. and Lawrence, A.W. (1978a): Autotrophic denitrification using elemental sulfur. *J. WPCF* 50, 1986-2001.
- Batchelor, B. and Lawrence, A.W. (1978b): A kinetic model for autotrophic denitrification using elemental sulfur. *Water Res.* 12, 1075-1084.

- Bayley, S.E., and Schindler, D.W. (1986): Source of Alkalinity in Precambrian Shield Watersheds under Natural Conditions and after Fire or Acidification in Hutchinson T. and Meema K. (eds.), Effects of Acid Deposition on Forests, Wetlands and Agricultural Ecosystems, Springer-Verlag, New York.
- Bayley, M.L., Davidson, L., and Headley, T.R. (2003): Nitrogen removal from domestic effluent using subsurface flow constructed wetlands: Influence of depth, hydraulic residence time and pre-nitrification. *Water Sci. Technol.* 48 (5), 175-182.
- Beadle, C.L. (1982): Plant growth analysis. In: Coombs, J., Hall, D.O. (Eds.), Techniques in bioproductivity and photosynthesis. Pergamon Press, New York City, NY, pp. 20-24.
- Beccari, M.; Pascino, R.; Ramadori, R. and Tandoi, V. (1980): Inhibition effects of nitrification by typical compounds in coke plant wastewaters. *Environ. Technol.* 1, 245-252.
- Bentzen, G.; Smith, A.T.; Bennet, D.; Webster, N.J.; Reinholt, F.; Sletholt, E. and Hobson, J. (1995): Controlled dosing of nitrate for prevention of H<sub>2</sub>S in a sewer network and the effect on the subsequent treatment process. *Water Sci. Technol.* 31 (7), 293-302.
- Berner, R.A. (1984): Sedimentary pyrite formation: an update. *Geochim. Cosmochim. Acta* 48, 605-615.
- Bezbaruah, A.N. and Zhang, T.C. (2004): pH, redox, and oxygen microprofiles in the rhizosphere of Bulrush (*Scirpus validus*) in a constructed wetland treating municipal wastewater. *Biotechnol. Bioeng.* 88 (1), 60-70.
- Bhatia, S.P. (1978): Organosulfur emissions from industrial sources. In: Nriagu J.O. (ed.), Sulphur in the Environment, part I, John Wiley and Sons, Canada.
- Blaabjerg, V. and Finster, K. (1998): Sulphate reduction associated with roots and rhizomes of the marine macrophytes *Zostera marine*. *Aquat. Microb. Ecol.* 15 (3), 311-314.

- Boon, A.G. (1995): Septicity in sewers: causes, consequences and containment. *Water Sci. Technol.* 31 (7), 237-253.
- Borum, J.; Pedersen, O.; Greve, T.M.; Frankovich, T.A.; Zieman, J.C.; Fourqurean, J.W. and Madden, C.J. (2005): The potential role of plant oxygen and sulphide dynamics in die-off events of the tropical seagrass, *Thalassia testudinum*, in Florida Bay. *J. Ecol.* 93, 148-158.
- Bostick, B.C.; Hansel, C.M.; La Force, M.J. and Fendorf, S. (2001): Seasonal fluctuations in zinc speciation within a contaminated wetland. *Environ. Sci. Technol.* 35, 3823-3829.
- Boudou, J.P.; Chehimi, M.; Broniek, E.; Siemieniowska T. and Bimer, J. (2003): Adsorption of H<sub>2</sub>S or SO<sub>2</sub> on an activated carbon cloth modified by ammonia treatment. *Carbon* 41 (10), 1999-2007.
- Boulègue, J. and Michard, G. (1979): Sulfur speciations and redox processes in reducing environments. In: *Chemical modelling in aqueous systems*, ed. E.A. Jenne. Am. Chem. Soc. Symp. Ser. 93, pp. 25-50. Washington, DC.
- Boulègue, J.; Lord, C.J. and Church, T.M. (1982): Sulfur speciations and associated trace metals (Fe, Cu) in the pore waters of Great Marsh, Delaware. *Geochim. Cosmochim. Acta* 46, 453-464.
- Bowker, R.P.G.; Smith, J.M. and Webster, N.A. (1985): EPA-625/185/018, USEPA. Cincinnati, Ohio.
- Börner, T. (1992): Factors influencing the efficiency of constructed wetlands. Verein zur Förderung der Industrie für Wasserversorgung, Abwasserbeseitigung und Raumplanung (WAR, Vol. 58), Darmstadt, Germany (In German).
- Brix, H. (1987): Treatment of wastewater in the rhizosphere of wetland plants- the Root-Zone Method. *Wat. Sci. Technol.* 19, 107-118.
- Brix, H. and Schierup, H. (1989): The use of aquatic macrophytes in water-pollution control. *Ambio.* 18, 100-1007.

- Brix, H. (1994): Functions of macrophytes in constructed wetlands. *Water Sci. Technol.* 29, 71-78.
- Brodie, G.A., Hammer, D.A. and Tomljanovich, D.A. (1988): Constructed wetlands for acid drainage control in the Tennessee Valley. In: *Proc. Mine Drainage and Surface Mine Reclam. Vol. 1: Mine Water and Mine Waste.* 325-331, U.S. Dept. Interior, Bur. Mines Info. Circ. IC-9183.
- Brodie, G.A.; Hammer, D.A. and Tomljanovich, D.A. (1989a): Treatment of acid mine drainage with a constructed wetland at the Tennessee Valley Authority 950 coal mine. In: *Constructed Wetlands for Wastewater Treatment* (Ed. D.A. Hammer), 201-209. Lewis Publishers.
- Brodie, A.G.; Hammer, D.A. and Tomljanovich, D.A. (1989b): Constructed wetlands for treatment of ash pond seepage. In: *Constructed Wetlands for Wastewater Treatment* (Ed. D.A. Hammer), 211-219. Lewis Publishers.
- Brix, H. (1987): Treatment of wastewater in the rhizosphere of wetland plants- the Root-Zone Method. *Water Sci. Technol.* 19, 107-118.
- Brodie, G.A. (1991): Achieving compliance with staged, aerobic, constructed wetlands to treat acid main drainage. In: *Conf. Mine Drainage and Surface Mine Reclam.* (1), 325-331.
- Cadena, F.P. and Peters, R.W. (1998): Evaluation of chemical oxidizer for hydrogen sulphide control. *J. Water Pollut. Control Fed.* 60, 1259-1263.
- Cadenhead, P. and Sublette, K.L. (1990) : Oxidation of hydrogen sulfide by *Thiobacilli*. *Biotechnol. Bioeng.*, 35 (11): 1150-1154
- Caffrey, J.M. and Kemp, W.M. (1991): Seasonal and spatial patterns of oxygen production, respiration and root-rhizome release in *Potamogeton perfoliatus* L. and *Zostera marina* L. *Aquat. Bot.* 40, 109-128.
- Calabrese, J.P.; Bhumbla, D.K.; Sexstone, A.J.; Bissonnette, G.K. and Sencindiver, J.C. (1990): In *Proc. Min Reclam Conf.* (Eds. J.J. Skousen, J.C. Sencindiver and D.E. Samuel), 351. W. Virginia University, Morgantown, W.V.

- Carlson, P.R.; Yarbrow, L.A. and Barber, T.R. (1994): Relationship of sediment sulfide to mortality of *Thalassia testudinum*, Florida bay. Bulletin of Marine Science. 54, 733-746.
- Carlson, D.A. and Leiser, C.P. (1966): Soil beds for the control of sewage odors. J. WPCF, 829-840.
- CDMB, 2003. Evaluation of the process of biooxidation of H<sub>2</sub>S in the effluent of UASB for the odor control.
- Chanton, J.P. and Dacey, J.W.H. (1991): Effects of vegetation on methane flux, reservoirs, and carbon isotopic composition. In: Sharkey, T.; Holland, E. and Mooney, H. (Eds.), Trace Gas emissions by plants. Academic Press, San Diego, CA, USA. 65-92.
- Chambers, R. M.; Mozdzer, T. J. and Ambrose, J.C. (1998): Effects of salinity and sulfide on the distribution of *Phragmites australis* and *Spartina alterniflora* in a tidal. Aquat. Botany. 62(3), 161-169.
- Chambers, L.A. and Trudinger, P.A. (1975). Are thiosulfate and trithionate intermediates in dissimilatory sulphate reduction? J. Bacteriol. 123 (1), 36-40.
- Cheremisinoff, P.N. (1988): Industrial Odour Control. Butterworth - Heinemann Ltd. Oxford.
- Chernicharo, C.A. de L. (1997): Reactores Anaeróbios. UFMG. Belo Horizonte. MG. 246p.
- Cho, K.S.; Hirai, M. and Shoda, M. (1992): Degradation of hydrogen sulfide by *Xanthomonas* sp. Strain DY44 isolated from Peat. Appl. Environ. Microbiol. 58 (4), 1183-1189.
- Choi, J.H.; Park, S.S. and Jaffé, P.R. (2005): The effect of emergent macrophytes on the dynamics of sulphur species and trace metals in wetland sediments. Environmental Pollution 140, 286-293.
- Čížková-Kočnalová, H.; Kvet, J. and Thompson, K. (1992): Carbon starvation: a key to read decline in eutrophic lakes. Aquat. Bot. 43, 105-113.

- Claus, G. and Kutzner, H.J. (1985): Autotrophic denitrification by *Thiobacillus denitrificans* in a packed reactor. Appl. Microbiol. Biotechnol. 22, 289-296.
- Colmer, T.D.; Gibberd, M.R.; Wiengweera, A. and Tinh, T.K. (1998): The barrier to radial oxygen loss from roots of rice *Oryza sativa* L. is induced by growth in stagnant solution. J. Experimental Botany. 51, 1-54.
- Connell, E.L.; Colmer, T.D. and Walker, D.I. (1999): Radial oxygen loss from intact roots of *Halophila ovalis* as a function of distance behind the root tip and shoot illumination. Aquat. Botany. 63, 219-228.
- Cork D.J. and Ma, S. (1982): Acid-gas bioconversion favors sulfur production. Biotechnology and Bioengineering Symp. 12: 285-290.
- Cooper, P.F.; Job, G.D.; Green, M.B. and Shutes, R.B.E. (1996): Reed beds and constructed wetlands for wastewater treatment. Medmenham. Marlow, UK: WRC Publications.
- Cooper, P.E. and de Maeseneer, J. (1996): Hybrid systems- what is the best arrangement for the vertical and horizontal flow stage? In: IEWQ specialist group on the use of Macrophytes for Water Pollution Control Newsletter, No.15, 8-13.
- Cullimore, D.R. (1991): Practical manual for groundwater microbiology. Chelsea, MI: Lewis Publ.
- Dacey, J.W. (1980): Internal winds in the water-lilies: adaptation for life in anaerobic sediments. Science 210, 1017-1019.
- Di Toro, D.M.; Mahony, J.D.; Hansen, D.J.; Scott, K.J.; Hicks, M.B.; Mayr, S.M. and Redmond, M.S. (1990): Toxicity of cadmium in sediments: the role of acid volatile sulfide. Environ. Toxicol. Chem. 9, 1487-1502.
- Dvorak, D.H.; Hedin, R.S.; Edenborn, H.M. and McIntire, P.E. (1991): Treatment of metal contaminated water using bacterial sulphate reduction: Results from pilot scale reactors. In: Proc. National Meeting Am. Soc. Surf. Mining and Reclam. 109-122.



- El-Shatnawi, M.K.J. and Makhadmeh, I.M. (2001): Ecophysiology of the plant-rizhosphere system. *J. Agron. Crop Sci.* 187, 1-9.
- Emerson, S.; Jacobs, L. and Tebo, B. (1983): The behavior of trace metals in marine anoxic waters: solubilities at the oxygen-hydrogen sulfide interface. In: Wong, C.S.; Boyle, E.; Bruland, K.W.; Burton, J.D. and Goldberg, E.D. (Eds.), *Trace metals in sea water*. Plenum Press, New York, NY, USA., 579-608.
- EPS. (1984): *Technical information for problem spills: Hydrogen Sulphide*, Ottawa, Canada.
- Eger, P. and Lapakko, K. (1989): Use of wetlands to remove nickel and cooper from mine drainage. In: *Constructed Wetlands for Wastewater Treatment* (Ed. D.A. Hammer), 780-787. Lewis Publishers.
- Eun-Ku, L.; Kwang-Deog, J.; Oh-Shim J. and Yong-Gun, S. (2005): Selective oxidation of hydrogen sulfide to elemental sulfur with Fe/MgO catalysts in a slurry reactor. *Bull. Korean Che. Soc.* 26 (2), 281-284.
- Fagan, M. and Walton, J.R. (1999): Headworks odour control with hydrogen peroxide: a review of recent studies. *US Peroxide.* 949, 661-6713.
- Feng, J.N and Hsieh, Y.P. (1998): Sulphate reduction in freshwater wetland soil and the effects of sulphate and substrate loading. *J. Environ. Qual.* 27 (4), 968-972.
- Fernandez-Polanco, F.; Martínez, B.; Olmedo, F., and García del Valle, J. (1996): Removal of H<sub>2</sub>S using a chemical scrubber and biological oxidation. In: *Proc. IV Workshop and Seminar of Anaerobic Wastewater Treatment in Latin America (in Spanish)*, November 19-22, Bucaramanga, Colombia.
- Ferrera, I.; Massana, R.; Casamayor, E.O.; Balague, V.; Sanchez, O.; Pedros-Alio, C. and Mas, J. (2004): High-diversity biofilm for the oxidation of sulfide containing effluents. *Appl. Microbiol. Biotechnol.* 64 (5), 726-734.
- Flere, J.M. and Zhang, T.C. (1999): Nitrate removal with sulfur-limestone autotrophic denitrification processes. *J. Environ. Eng.* 8 (125), 721-729.

- Flessa, H. (1991): Redoxprozesse in Böden in der Nähe von wachsenden und absterbenden Pflanzenwurzeln. PhD. Thesis, Technical University of München, Germany.
- Foresti, E. (2001): Perspectives on anaerobic treatment in developing countries. *Water Sci. Technol.* 44(8) 141-148.
- Frechen, F.B. (1988): Odour emissions and odour control at wastewater treatment plants in West Germany. *Water Sci. Technol.* 20, 261-266.
- Fürtig, K.; Rueeggsegger, A.; Brunold, C. and Braendle, R. (1996): Sulphide utilization and injuries in hypoxic roots and rhizomes of common reed (*Phragmites australis*). *Folia Geobot. Phytotax.* 31 (1), 143-151.
- García, J.; Aguirre, P.; Mujeriego, R.; Huang, Y.; Ortiz, L. and Bayona, J.M. (2004): Initial contaminant removal performance factors in horizontal flow reed beds used for treating urban wastewater. *Water Res.* 38, 1669-1678.
- Gayle, B.P.; Boardman, G.D.; Sherrad, J.H. and Benoit, R.E. (1989): Biological denitrification of water. *J. Environ. Eng. ASCE* 115, 930-943.
- Gearheart, R.A. (1992): Use of constructed wetlands to treat domestic wastewater, city of Arcata, California. *Water Sci. Technol.* 26 (7-8), 1625-1637.
- Geller, G. (1996): Horizontal flow system's for wastewater treatment: long term scientific and practical experiences; recommendation. In. *Proc. 5<sup>th</sup>. Int. Conf. Wetland Systems for Water Pollution Control*, chapter III/2. Universität für Bodenkultur Wien, Austria.
- Giraldo, E. and Zárate, E. (2001): Removal of hydrogen sulphide DBO from brackish water using vertical flow wetlands in a Caribbean environment, *Water Sci. Technol.* 44 (11-12), 361-367.
- Goldhaber, M.B. and Kaplan, I.R. (1974): The sulfur cycle. – In: Goldberg, E.D. (ed.) *The sea (Vol. 5), marine chemistry.* Wiley & Sons, New York, 569-665.

- Goldman, C.R. and Horne, A. J. (1983): Limnology. New York: McGraw-Hill.
- Goodman, J.L.; Moore, K.A. and Dennison, W.C. (1995): Photosynthetic response of eelgrass (*Zostera marina* L.) to light and sediment sulfide in a shallow barrier island lagoon. Aquatic Botany. 50, 37-47.
- Guckert, J.B.; Carr, G.J.; Johson, T.D.; Hamm, B.G.; Davidson, D.H. and Kumagi, Y. (1996): Community analysis by Biolog: curve of integration for statistical analysis of activated sludge microbial habitats. Journal of Microbiological Methods, 27: 183-197.
- Haberl, R.; Perfler, R. and Mayer, H. (1995): Constructed wetlands in Europe. Water Sci. Technol. 32 (3), 305-315.
- Hale, M.G. and Moore, L.D. (1979): Factors affecting root exudation II: 1970-1978. Adv. Agron. 31, 93-124.
- Hammack, R.W. and Hedin, R.S. (1989): Microbial sulfate reduction for the treatment of acid mine drainage: A laboratory study. In: Proc. Conf. "Reclamation, A Global Perspective", 673-680.
- Hammack, R.W. and Edenborn, H.M. (1991): The removal of nickel from mine waters using bacterial sulfate reduction. In: Proc. National Meeting Am. Soc. Surf. Mining and Reclam. 97-107.
- Hammer, D.A. and Bastian, R.K. (1989): Wetlands ecosystems: Natural water purifiers? In: Constructed Wetlands for Wastewater Treatment (ed. D.A. Hammer), 5-19. Lewis Publishers.
- Hanaki, K. and Wantawin, S. (1990): Nitrification at low-levels of dissolved-oxygen with and without organic loading in a suspended-growth reactor. Water Res. 24 (3), 297-302.
- Harkness, N. (1980): Chemistry of Septicity. J. Effluent Water Treatment. 20, 16-23.

- Hedin, R.S.; Hyman, D.M and Hammack, R.W. (1988): Implication of sulfate-reduction and pyrite formation processes for water quality in a constructed wetland: Preliminary observations. In: Proc. Conf. Mine Drainage and Surf. Mine Reclam. Vol. 1: Mine Water and Mine Waste, 382-388. U.S. Dept. Interior, Bur. Mines Info. Circ. IC-9183.
- Hedin, R.S.; Hammack, R.W. and Hyman, D.M. (1989): Potential importance of sulfate reduction processes in wetlands constructed to treat mine drainage. In: Constructed Wetlands for Wastewater Treatment (Ed. D.A. Hammer), 508-514. Lewis Publishers.
- Helder, W and de Vries, R.T.P. (1983): Estuarine nitrite maxima and nitrifying bacteria (Ems-Dollar estuary). *Neth. J. Sea Res.* 17, 1-18.
- Henze, M.; Harremos, P.; Jansen, J. C. and Arvin, E. (1995): *Wastewater Treatment*. Springer Verlag. Germany.
- Hiel; M.T. and Kerins, F.J. (1988): The Tracy wetlands: A case study of two passive mine drainage treatment system in Montana. In: Proc. Conf. Mine Drainage and Surf. Mine Reclam. Vol. 1: Mine Water and Mine Waste, 352-358. U.S. Dept. Interior, Bur. Mines Info. Circ. IC-9183.
- Hobson, J. (1995): The odour potential: a new tool for odour management. *Water Environ. Manage.* 9, 458-463.
- Holmer, M. and Bondgaard, E.J. (2001): Photosynthetic and growth response of eelgrass to low oxygen and high sulfide concentration during hypoxic events. *Aquatic Botany.* 70, 29-38.
- Holmer, M.; Frederiksen, M.S. and Mollegaard, H. (2005): Sulphur accumulation in eelgrass (*Zostera marina*) and effect of sulphur on eelgrass growth. *Aquatic Botany.* 81 (4), 367-379.
- Howarth, R.W. and Jorgensen, B.B. (1984): Formation of  $^{35}\text{S}$ -labelled elemental sulfur and pyrite in coastal marine sediment (Limfjorden and Kysing Fjord, Denmark) during short-term  $^{35}\text{SO}_4^{2-}$  reduction measurements. *Geochim. Cosmochim. Acta* 48. 1807-1818.

- Huerta-Diaz, M.A.; Tessier, A. And Carignan, R. (1998): Geochemistry of trace metals associated with reduced sulphur in freshwater sediments. *Applied Geochemistry*. 13, 213-233.
- Hutchinson, G.E. (1975): A treatise on limnology. III. Limnological Botany. John Wiley, New York: 660 pp.
- Hwang, Y.; Matsuo, T.; Hanaki, K. and Suzuki, N. (1995): Identification and quantification of sulphur and nitrogen containing odorous compounds in wastewater. *Water Res.* 29 (2), 711-718.
- Jackson, L. M. and J.E. Myers., 2002. Evaluation of Subsurface Flow Wetlands vs. Free-water Surface wetlands Treating NPR-3 produced Water – Yea no. 2. 1-15. United States Department of Energy / Rocky Mountain Oilfield testing Centre.
- Jaffé, P.R. ; Wang, S.; Kallin, P.L. and Smith, S.L. (2001): The dynamics of arsenic in saturated porous media: fate and transport modelling for deep-water sediments. In: Hellman, R. and Wood, S.A. (Eds.) *Water Rock Interactions, Ore Deposits, and Environmental Geochemistry; A tribute to David Crerar*. The Geochemical Society Special Publication No 7.
- Jordao, E.P. and Volschan, I. (2004): Cost-effective solutions for sewage treatment on developing countries – the case of Brazil. *Water Sci. Technol.* 50 (7), 237-242.
- Jorgensen, B.B. (1990): A thiosulfate shunt in the sulfur cycle of marine sediments. *Science* 249, 152-154.
- Jorgensen, B.B.; Isksen, M.F. and Jannasch, H.W. (1992): Bacterial sulfate reduction above 100 °C in deep-sea hydrothermal vent sediments. *Science*. 258, 1756-1757.
- Justin, P. and Kelly, D.P. (1978): Metabolic changes in *Thiobacillus denitrificans* accompanying the transition from aerobic to anaerobic growth in continuous chemostat culture. *J. Gen. Microbiol.* 107, 131-137.

- Kadlec, R.H. (1987): Northern natural wetland water treatment systems. In: Reddy, K.R. and Smith, W.H., editors. Aquatic plants for water treatment and resource recover. Orlando, USA: Magnolia Publishing. 83-98.
- Kadlec, R.H. and Knight, R.L. (1996): Treatment wetlands. CRC Press/Lewis Publishers, Boca Raton.
- Kadlec, R.H.; Knight, R.L.; Vymazal, J.; Brix, H.; Cooper, P. and Harberl, R. (2000): Constructed Wetlands for Pollution Control. Report no. 8, IWA Publishing, London.
- Kalogo, Y. and Verstraete, W. (2001): Physical and biological performance of self-inoculated UASB treating raw domestic sewage. J. Environ. Eng. –ASCE. 127 (2), 179-183.
- Kappelmeyer, U., Wießner, A., Kusch, P., and Kästner, M. (2002): Operation of a universal test unit for planted soil filter – Planted fixed bed reactor. Eng. Life Sci. 2, 311-315.
- Karamenko, L.Y. (1969): Biogenic factors in the formation of sedimentary ores. Intl. Geol. Rev. 11, 271-281.
- Kaseva, M.E.; Mbwette, T.S. and Katima, J.H. (2002): Domestic sewage treatment in a pilot plant composed of septic tank and constructed wetlands-a case study in Dar es Salam, Tanzania. In: Presented at the Eighth IWA Conference on Wetland Systems for water Pollution Control, Arusha, Tanzania.
- Kaseva, M.E. (2004): Performance of sub-surface flow constructed wetland in polishing pre-treated wastewater- a tropical case study. Water Res. 38 (3), 681-687.
- Kaiser, S.K., Guckert, J.B. and Gledhill, D.W. (1998): Comparison of activated sludge microbial communities using Biolog TM Microplates. Water Sci. Tech., 37 (4-5), 57-63.
- Kemp, W.M. and Murray, L. (1986): Oxygen release from roots of the submerged macrophytes *Potamogeton perfoliatus* L.: regulating factors and ecological implications. Aquat. Bot. 26, 271-283.

- Kim, S.Y. and Geary, P.M. (2001): The impact of biomass harvesting on phosphorous uptake by wetland plants. *Water Sci. Technol.* 44, 61-67.
- Klusman, R. and Machemer, S.D. (1991): Natural processes of acidity reduction and metal removal from acid mine drainage. In: *Geology in Coal Resource Utilizations* (ed. D.C. Peters), 513-540. Tech Books, Fairfax, VA.
- Knight, R.L. (1997): Wildlife habitat and public use benefits of treatment wetlands. *Water Sci. Technol.* 35 (5), 35-43.
- Knight, R.L.; Payne Jr., V.W.E.; Borer, R.E.; Clarke, Jr., R.A. and Pries, J.H. (2000): Constructed wetlands for livestock wastewater management. *Ecol. Eng.* 15, 41-55.
- Koch, M.S. and Erskine, J.M. (2001): Sulfide as a phytotoxin to the tropical seagrass *Thalassia testudinum*. interactions with light, salinity and temperature. *J. Exp. Mar. Biol. Ecol.* 266(1), 81-95.
- Koenig, A. and Liu, L.H. (1996): Autotrophic denitrification of landfill leachate using elemental sulphur. *Water Sci. Technol.* 34 (5-6), 469-476.
- Koenig, A. and Liu, L.H. (2001a): Microbial aspects of autotrophic denitrification of wastewater. In: Matsuo, T.; Hanaki, K.; Takizawa, S. and Satoh, H. (Eds.) *Advances in Water and Wastewater Treatment Technology*. Elsevier, The Netherlands, 217-226.
- Koenig, A. and Liu, L.H. (2001b): Kinetic model of autotrophic denitrification in sulfur packed-bed reactors. *Water Res.* 35 (8), 1969-1978.
- Končalová, H. (1990): Anatomical adaptations to waterlogging in roots of wetland graminoids: limitation and drawbacks. *Aquatic Botany.* 38, 127-134.
- Krause, G.H. and Weis, E. (1984): Chlorophyll fluorescence as a tool in plant physiology. II. Interpretation of fluorescence signals. *Photosynthesis Research.* 5, 139-157.

- Krause, G.H. and Weis, E. (1991): Chlorophyll fluorescence and photosynthesis: The basics. *Annual Review of Plant Physiology and Plant Molecular Biology*. 42, 313-349.
- Kuschik, P.; Wießner, A.; Kappelmeyer, U. and Weißbrodt, E. (2003): Annual cycle of nitrogen removal by pilot-scale subsurface horizontal flow in a constructed wetland under moderate climate. *Water Res.* 37, 4236-4242.
- Kutschera, L. and Lichtenegger, E. (1982): *Wurzelatlas Mitteleuropäischer Grünlandpflanzen I: Monocotyledoneae*. Gustav Fischer, Stuttgart/New York.
- Laber, J.; Haberl, R. and Shrestha, R. (1999): Two stage constructed wetland for treatment hospital wastewater in Nepal. *Water Sci. Technol.* 40 (3), 317-324.
- Lamers, L.P.; Tomassen, H.B. and Roelofs, J.G. (1998): Sulfate induced eutrophication and phytotoxicity in freshwater wetlands. *Environ. Sci. Technol.* 32, 199-2005.
- Lang, M.E. and Jager, R.A. (1992): Odour control for municipal sludge composting. *Bio Cycle*. 76-85.
- Laanbroek, H.J. and Gerards, S. (1993): Competition for limiting amounts of oxygen between *Nitrosomonas europaea* and *Nitrobacter winogradsky* grown in mixed continuous culture. *Arch. Microbiol.* 159 (5), 453-459.
- Lee, R.W. (1999): Oxidation of sulphide by *Spartina alterniflora* roots. *Limnol. Oceanogr.* 44 (4), 1155-1159.
- Lefroy, R.D.B.; Chaitep, W. and Blair, G.J. (1994): Release of sulfur from rice residues under flooded and non-flooded soil conditions. *Aust. J. Agric. Res.* 45, 657-667.
- Lemon, E.; Braybrook, T.; Bis, G. and Rozema, L. (2003): Oxygen status in vertical flow wetlands. *Water Sci. Technol.* 32 (3), 103-109.
- Liebmann, H. (1962): *Handbuch der Frischwasser- und Abwasserbiologie*, Vol. 1 and 2. Oldenbourg, München.



- Liesack, W.; Schnell, S. and Revsbech, N.P. (2000): Microbiology of flooded rice paddies. *FEMS Microbiol. Rev.* 24 (5), 625-645.
- Machemer, S.D. and Wildeman, T.R. (1992): Adsorption compared with sulfide precipitation as metal removal processes from acid mine drainage in a constructed wetland. *J. Contam. Hydrol.* 9, 115-131.
- Mamta, T.; Tamama, H.A.A. and Jamila, H.A.A. (1995): Effect of aeration on generation and emission of hydrogen sulfide in wet wells of lifting stations. *Water, Air, and Soil Pollut.* 81, 385-399.
- Mandi, L.; Bouhoum, K. and Ouazzania, N. (1998): Application of constructed wetlands for domestic wastewater treatment in an arid climate. *Water Sci. Technol.* 38, 379-387.
- Martínez, C.P. and Zamorano, J.P. (1996): Practical experience on anaerobic treatment and odor control in a yeast factory. In: *Proc. IV Workshop and Seminar of Anaerobic wastewater Treatment in Latin America (in Spanish)*, November, 19-22, Bucaramanga, Colombia.
- Mashauri, D.A.; Senzia, M.A. and Mayo, A.W. (2003): Suitability of constructed wetlands and waste stabilisation ponds in wastewater treatment: nitrogen transformation and removal. *Physics and Chemistry of the Earth.* 28, 1117-1124.
- Mbuligwe, S.E. (2004): Comparative effectiveness of engineered wetland system in the treatment of anaerobically pre-treated domestic wastewater. *Ecological Engineering.* 23, 269-284.
- McGovern, J.E. and Clarkson, C.R. (1994): The development of Northumbrian water Limited's approach to odour abatement for wastewater facilities. *CIWEM, Symposium on Odour Control and Prevention in the Water Industry*, University of Newcastle upon Tyne.

- McIntire, P.E. and Edenborn, H.M. (1990): the use of bacterial sulfate reduction in the treatment of drainage from coal mines. In: Proc. Mining and Reclam. Conf. (Eds. J.J. Skousen, J.C. Sencindiver and D.E. Samuel), 409-415, W. Virginia University, Morgantown. W.V.
- McJannet, C.L.; Keddy, P.A. and Pick, F.R. (1995): Nitrogen and phosphorous tissue concentrations in 41 wetland plants: a comparison across habitats and functional groups. *Funct. Ecol.* 9, 231-238.
- Mendelssohn, I.A. and McKee, K.L. (1988): *Spartina alterniflora* die-back in Louisiana: time-course investigation of soil waterlogging effects. *J. Ecol.* 76, 509-521.
- Mendelssohn, I.A.; Kleiss, B.A. and Wakeley, J.S. (1995): Factors controlling the formation of oxidized root channels – a review. *Wetlands.* 15, 37-46.
- Metcalf and Eddy, (2005): *Wastewater Engineering Treatment and Reuse.* 4. ed. New York, McGraw-Hill Book. 1819p.
- Middelburg, J.J. and Van Der Nat, F.W.A. (1998): Effect of two common macrophytes on methane dynamics in freshwater sediments. *Biogeochemistry.* 43, 79-104.
- Moorhead, K.K. and Reddy, K.R. (1988): Oxygen transport through selected aquatic macrophytes. *J. Envir. Qual.* 17, 138-142.
- Moormann, H.; Kusch, P. and Stottmeister, U. (2002): The effect of rhizodeposition from helophytes on bacterial degradation of phenolic compounds. *Acta Biotechnol.* 22, 107-112.
- Morse, J.W.; Millero, F.J.; Cornwell, J.C. and Rikard, D. (1987): The chemistry of the hydrogen sulfide and iron sulfide systems in natural waters. *Earth Sci. Rev.* 24, 1-42.
- Mulder, A.; van de Graaf, A.A.; Robertson, L.A. and Kuenen, J.G. (1995): Anaerobic ammonium oxidation discovered in a denitrifying fluidized bed reactor. *FEMS Microbiol. Ecol.* 16 (3), 177-184.

- Müller, G. (1966) Die Verteilung von Eisenmonosulfid ( $\text{FeS} \times n \text{H}_2\text{O}$ ) und organischer Substanz in den Bodensedimenten des Bodensees- ein Beitrag zur Frage der Eutrophierung des Bodensees. Gas- u. Wass. Fach 107, 364-368.
- Ngo, V. (1987): Boosting pond performance with aquaculture. Ops. Forum 4,20-24.
- Noyola, A.; Morgan-Sagastume, J.M. and López-Hernández, J.E. (2006): Treatment of biogas produced in anaerobic reactors for domestic wastewater: odor control and energy/resource recovery. Environ. Sci. Technol. 5, 93-114.
- Ongcharit, C.; Shah, Y.T. and Sublette, J.L. (1990): Novel immobilized cell reactor for microbial oxidation of  $\text{H}_2\text{S}$ . Chem. Eng. Sci. 45, 2383-2389.
- Oranjewoud, J. (2000): Rapport Baggerplan Venkoelen. In: Opdracht van Waterschap Peel en Maasvallei and Zuiveringschap Limburg. Document Nr. 0589-44094.
- Ottova, V.; Balcarova, J. and Vymazal, J. (1997): Microbial characteristics of constructed wetlands. Water Sci. Technol. 35, 117-123.
- Overmann, J. and van Gemerden H. (2000): Microbial interactions involving sulfur bacteria: implications for the ecology and evolution of bacterial communities. FEMS Microbiol. Rev. 24 (5), 591-599.
- Paing, J.; Picot, B. and Sambuco, J. P. (2003): Emission of  $\text{H}_2\text{S}$  and mass balance of sulfur in anaerobic ponds. Water Sci. Technol. 48 (2), 227-234.
- Pedersen, O.; Binzer, T. and Borum, J. (2004): Sulphide intrusion in eelgrass (*Zostera marine* L.). Plant Cell Environm. 27 (5), 595-602.
- Perdomo, S., Fujita, M. and Furukawa, K. (1996): Oxygen transport through *Pistia stratiotes* L. In Pro. 5<sup>th</sup> Intern. Conf. Wetland systems for water Pollutions Control. Pp. VIII/4-1-VIII/4-8. Universität für Bodenkultur Wien. Vienna, Austria.
- Pettecrew, E.L. and Kalff, J. (1992): Water flow and clay retention in submerged macrophytes bed. Can J. Fish. Aquat. Sci. 49, 2483-2489.

- Pezeshki, S.R.; Delaune, R.D. and Lindau, C.W. (1998): Interaction among sediment anaerobiosis, nitrogen uptake and photosynthesis of *Spartina alterniflora*. *Physiol. Plant.* 74, 561-565.
- Platzer, C. and Netter, R. (1994): Factors affecting nitrogen removal in horizontal flow reed beds. *Water Sci. Technol.* 29, 319-324.
- Polanco, F.F., Polanco M.F., Fernandez, N., Urena, M.A., García, P.A., and Villaverde, S. (2001): New process for simultaneous removal of nitrogen and sulfur under anaerobic condition. *Water Res.* 35, 1111-1114.
- Polprasert, C. and Chatsanguthai, S. (1989): Sulfide production during anaerobic lagoon treatment of Tapioca wastewater. *Environment International.* 14, 563-566.
- Rands, M.B.; Cooper, D.E.; Woo, C.P.; Fletcher, G.C. and Rolfe, F.K. (1981): Compost filters for H<sub>2</sub>S removal from anaerobic digestion and rendering exhausts. *J. WPCF* 53, 185-189.
- Ranieri, E. (2003): Hydraulics of subsurface flow constructed wetlands in semi arid climates conditions. *Water Sci. Technol.* 47 (7-8), 49-55.
- Raven, J.A. and Scrimgeour, C.M. (1997): The influence of anoxia on plants of saline habitats with special reference to the sulphur cycle. *Annals of Botany.* 79, 79-86.
- Reddy, K.R. and Patrick, W.H. (1984): Nitrogen transformations and loss in flooded soils and sediments. *CRC Crit. Rev. Environ. Contr.* 13, 273-309.
- Reddy, K.R.; Patrick, W.H. and Lindau, C.W. (1989): Nitrification-denitrification at the plant root-sediment interface in wetlands. *Limnol. Oceanogr.* 34, 1003-1013.
- Rethmeier, J., Rabenstein, A., Langer, M. and Fischer, U. (1997): Detection of traces of oxidized and reduced sulfur compounds in small samples by combination of different high-performance liquid chromatography methods, *Journal of chromatography A* 1997, 760, 295-302.

- Reynolds, J.S.; Macheimer, S.D.; Wildeman, T.R.; Updegraff, D.M. and Coher, R. (1991): Determination of the rate of sulfide production in a constructed wetland receiving acid mine drainage. In: Proc. National Meeting Am. Soc. Surf. Mining and Reclam. 175-182.
- Rheinheimer, G. (1981): Aquatic microbiology. New York: Wiley-Interscience.
- Richards, P.W. and Clapham, A.R. (1941): *Juncus effusus* L. J. Ecol. 29, 375-380.
- Rovira, A.D. (1965): Interactions between plant roots and soil microorganisms. A. rev. Microbiol. 19, 241-266.
- Rovira, A.D. (1969): Plant root exudates. Bot. Rev. 35, 35-57.
- Sand-Jensen, K.; Prahl, C. and Stokhol, H. (1982): Oxygen release from roots of submerged aquatic macrophytes. Oikos 38, 349-354.
- Schippers, J.C.; Kruithof, J.C.; Mulder, F.G. and van Lieshout, J.W. (1987): Removal of nitrates by slow sulfur limestone filtration. Aqua 5, 274-280.
- Seidel, K. (1966): Reinigung von Gewässern durch höhere Pflanzen. Naturwissenschaften. 53, 289-297.
- Seitzinger, S.P. (1990): Denitrification in aquatic sediments. In: Denitrification in soil and sediment, Revsbech, N.P. and Sorensen, J. (Eds), 301-322.
- Sicher, R.C. (1998): Yellowing and photosynthetic decline of barley primary leaves in response to atmospheric CO<sub>2</sub> enrichment. Physiol Plant. 103, 193-200.
- Sicher, R.C. (1999): Photosystem-II activity is decreased by yellowing of barley primary leaves during growth in elevated carbon dioxide. Int. J. Plant Sci. 160, 849-854.
- Sikora, F.J.; Zim, T., Behrend, J.L.; Steinberg, S.L. and Coonrod, H.S. (1995): Ammonium removal in constructed wetlands with recirculating subsurface flow: removal rates and mechanisms. Water Sci. Technol. 32 (3), 193-202.

- Simpson, S.L.; Apte, S.C. and Batley, G.E. (1998): Effect on short-term resuspension events on trace metal speciation in polluted anoxic sediments. *Environ. Sci. Technol.* 32, 620-625.
- Smet, E. and Van Langenhove, H. (1998): Abatement of volatile organic sulphur compounds in odorous emissions from the bio-industry. *Biodegradation.* 9, 273-284.
- Smith, I.D.; Bis, G.N.; Lemon, E.R. and Rozema, L.R. (1996): A thermal analysis of a sub-surface, vertical flow constructed wetland. In *Pro. 5th Internat. Conf. Wetland System for Water Pollution Control*, pp. III/4-1-III/4-8. Vienna, Austria: Universität für Bodenkultur Wien.
- Smolders, A.J.P.; Nijboer, R.C. and Roelofs, J.G.M. (1995a): Prevention of sulphide accumulation and phosphate mobilization by the addition of iron (II) chloride to a reduced sediment: An enclosure experiment. *Freshwater Biology.* 343, 559-568.
- Smolders, A.J.P.; Roelofs, J.G.M. and Den Hartog, C. (1995): Internal eutrophication of aquatic ecosystems: mechanisms and possible remedies. *Acta Botanica Gallica.* 142, 707-717.
- Smolder, A.J.P. and Roelofs, J.G.M. (1996): The roles of internal iron hydroxide precipitation, sulphide toxicity and oxidizing ability in the survival of *Stratiotes aloides* roots at different iron concentrations in sediment pore water. *New Phytologist.* 1332, 253-260.
- Somes, N.L.G.; Breen, P.F. and Wong, T.H.F. (1996): Integrated hydrology and botanical design of stormwater control wetlands. In *Pro. 5th Internat. Conf. Wetland System for Water Pollution Control*, pp. III/4-1-III/4-8. Vienna, Austria: Universität für Bodenkultur Wien.
- Sorrell, B.K. and Armstrong, W. (1994): Effect of external oxygen demand on radial oxygen loss by juncos roots in titanium citrate solutions. *Plant Cell Environ.* 22, 1587-1593.

- Sorrell, B.K. (1999): On the difficulties of measuring oxygen release by root systems of wetland plants. *J. Ecol.* 82, 177-183.
- Sousa, J.T.; Van Haandel, A.; Lima, E.P.C. and Guimaraes, A.V.A. (2003): Performance of constructed wetland systems treating anaerobic effluents. *Water Sci. Technol.* 48 (6), 295-299.
- Stark, L.R.; Kolbash, R.L.; Webster, H.J.; Stevens, S.E.Jr.; Dionis, K.A. and Murphy, E.R. (1988): The Simco # 4 wetland: Biological patterns and performance of a wetland receiving mine drainage. In: Proc. Conf. Mine Drainage and Surface Mine Reclam. Vol. 1: Mine Water and Mine Waste, 332-344. U.S. Dept. Interior, Bur. Mines Info. Circ. IC-9183.
- Stillings, L.L.; Gryta, J.J. and Ronning, T.A. (1988): Iron and manganese removal in a *Typha*-dominated wetland during ten months following its construction. In: Proc. Conf. Mine Drainage and Surf. Mine Reclam. Vol. 1: Mine Water and Mine Waste, 317-324. U.S. Dept. Interior, Bur. Mines Info. Circ. IC-9183.
- Stottmeister, U.; Wießner, A.; Kusch, P.; Kappelmeyer, U.; Kästner, M.; Bederski, O.; Müller, R.A. and H. Moormann. (2003): Effects of plants and microorganisms in constructed wetlands for wastewater treatment. *Biotechnol. Adv.* 22 (1-2), 93-117.
- Sublette, K.L. and Sylvester, N.N. (1987): Oxidation of hydrogen sulfide by mixed cultures of *Thiobacillus denitrificans* and heterotrophs. *Biotechnol. Bioeng.* 29, 759-761.
- Takahashi, S.; Tamashiro, A.; Sakihama, Y.; Yamamoto, Y.; Kawamitsu, Y. and Yamasaki, H. (2002): High-susceptibility of photosynthesis to photoinhibition in the tropical plant *Ficus microcarpa* L. f. cv. Golden Leaves. *BMC Plant Biol.* 2 (2), 1-8.
- Tanner, C.C. (1996): Plants for constructed wetland treatment systems - a comparison of the growth and nutrient uptake of eight emergent species. *Ecological Engineering.* 7, 59-83.

- Tanner, C.C.; Adams, D.D. and Downes, M.T. (1997): Methane emissions from constructed wetlands treating agricultural wastewater. *J. Environ. Qual.* 26, 1056-1062.
- Tanner, C.C. (2002): Plants as ecosystem engineers in subsurface-flow treatment wetlands. *Water Sci. Technol.* 44 (11-12), 47-54.
- Thable, T.S. (1984): Einbau und Abbau von Stickstoffverbindungen aus Abwasser in der Wurzelraumanlage Othfresen, Dissertation, Gesamthochschule Kassel, Universität Hessen.
- Thomas, K.L.; Benstead, J.; Davies, K.L. and Lloyd, D. (1996): Role of wetland plants in the diurnal control of CH<sub>4</sub> and CO<sub>2</sub> fluxes in peat. *Soil Biol. Biochem.* 28, 17-23.
- Thurston, J.A.; Falabi, J.A.; Gerba, C.P.; Foster, K.E. and Karpiscak, M.M. (1996): Fate of indicator microorganisms, *Giardia*, and *Cryptosporidium* in two constructed wetlands. 5<sup>th</sup> International Conference on Wetland Systems for Water Pollution Control. Vienna, Poster 30-1.
- Tomlinson, E.J. and Bruce, A.M. (1979): Problems of septicity in biological treatment. Proc. Symposium: Septic Sewage – problems and solutions. The Institute of Water Pollution Control, Bournemouth, 84-95.
- Urban, N.R.; Brezonik, P.L.; Baker, L.A. and Sherman, L.A. (1994): Sulfate reduction and diffusion in sediments of little Rock Lake, Wisconsin. *Limnol. Oceanogr.* 39, 797-815.
- Urbanc-Bercic, O. and Bulc, T. (1994): Integrated constructed wetland for small communities. Proceedings of 4<sup>th</sup> Int. Conf. On wetlands systems for water pollution control. Guangzhou, P.R. China, 138-146.
- Van der Hoek, J.P.; Hijnen, W.A.M.; van Bennekom, C.A. and Mijnders, B.J. (1992): Biological nitrate from groundwater by sulphur/limestone denitrification. *J. Chem. Technol. Biotechnol.* 54, 197-200.



- Van der Welle, M.E.W.; Cuppens, M.; Lamers, L.P.M. and Roelofs, J.G.M. (2006): Detoxifying toxicants: interactions between sulphide and iron toxicity to freshwater plants. *Environ. Toxicol. and Chem.* 25 (6),1592-1597.
- Van der Welle, M.E.W.; Niggebrugge, K.; Lamers, L.P.M. and Roelofs, J.G.M. (2007): Differential responses of the freshwater wetland species *Juncus effusus* L. and *Caltha palustris* L. to iron supply in sulfidic environments. *Environ. Pollut.* 147, 222-230.
- Van Haandel, A.C. and Lettinga, G. (1994): Tratamento anaeróbio de esgotos: Um Manual Regiões de Clima Quente, EpGraf, Campina Grande-PB.
- Van Starckenburg, W. (1997): Anaerobic treatment of wastewater: State of the art. *Microbiology.* 66 (5), 588-596.
- Vartapetian, B.B. and Jackson, M.B. (1997): Plant adaptations to anaerobic stress. *Ann. Bot.* 79 (Suppl. A), 3-20.
- Vincent, A. and Hobson, J. (1988): Odour Control. CIWEM Monographs on Best Practice No. 2, Chartered Institution of Water and Environmental Management, Terence Dalton Publishing, London.
- Visser, E.J.W.; Colmer, T.D.; Blom, C.W.P.M. and Voeselek, L.A.C.J. (2000): changes in growth porosity and radial oxygen loss from adventitious roots of selected mono- and dicotyledonous wetland species with contrasting types of aerenchyma. *Plant Cell and Environ.* 23, 1237-1245.
- Vymazal, J. (2002): The use of sub-surface constructed wetlands for wastewater treatment in the Czech Republic: 10 years experience. *Ecol. Eng.* 18, 633-646.
- Walhauser and Puchelt, (1966): Sulfate reducing bacteria in sulfur springs and connate waters of Germany and Austria (in German). *Contrib. Min. Petrol.-Beitr. Mineral. Petrologie.* 13 (1), 12-30.
- Walz, H. GmbH. (2000): Portable fluorometer PAM-2000 and data acquisition software DA-2000. Handbook of operation, with examples of practical applications. 2<sup>nd</sup> Eds.

- Wang, F.Y. and Chapman, P.M. (1999): Biological implication of sulfide in sediment – A review focusing on sediment toxicity. *Environ. Toxicol. Chem.* 18, 2526-2532.
- Webster, T.S.; Devinny, J.S.; Torres, E.M., and Barrai, S.S. (1996): Biofiltration of odours, toxics and volatile organic compounds from publicly owned treatment works. *Environ. Prog.* 15, 141-147.
- Wetzel, R.G. (1990): Land-water interfaces: metabolic and limnological regulators. *Verhandlungen der International Vereinigung für Theoretische und Angewandte Limnologie.* 24, 6-24.
- Wiegant, W.M. (2001): Experiences and potential of anaerobic wastewater treatment in tropical regions. *Water Sci. Technol.* 44 (8), 107-113.
- Wieder, R.K. (1989): A survey of constructed wetlands for acid coal mine drainage treatment in the eastern United States. *Wetlands.* 9, 299-315.
- Wildeman, T.R. and Laudon, L.S. (1989): Use of wetlands for treatment of environmental problems in mining: Non coal mining applications. In: *Constructed Wetlands for Wastewater Treatment* (Ed. D.A. Hammer), 221-231. Lewis Publishers.
- Wildeman, T.R.; Macheimer, S.D.; Klusman, R.W.; Cohen, R.R. and Lemke, P.R. (1990): Metal removal efficiencies from acid mine drainage in the Big Five constructed wetland. In: *Proc. Mining and Reclam. Conf. Exhibition. Vol II*, (Eds. J.J. Skousen, J.C. Sencindiver and D.E Samuel), 417-424. W. Virginia University, Morgantown, W.V.
- Wind, T. and Conrad, R. (1995): Sulfur compounds, potential turnover of sulfate and thiosulfate, and number of sulfate-reducing bacterial in planted and unplanted paddy soil. *FEMS Microbiol. Ecol.* 18, 257-266.
- Wind, T. and Conrad, R. (1997): Localization of sulphate reduction in planted and unplanted rice field soil. *Biogeochemistry.* 37, 253-278.

- Wiessner, A.; Kappelmeyer, U.; Kusch, P.; Kästner, M. and Stottmeister, U. (2002): Abilities of helophyte species to release oxygen into rhizosphere with varying redox conditions in laboratory-scale hydroponic systems. *Int. J. Phytoremed.* 1, 1-15.
- Wiessner, A.; Kappelmeyer, U.; Kusch, P. and Kästner, M. (2005a): Sulphate reduction and the removal of carbon and ammonia in a laboratory-scale constructed wetland. *Water Res.* 39 (19), 4643-4650.
- Wiessner, A.; Kappelmeyer, U.; Kusch, P. and Kästner, M. (2005b): Influence of the redox condition dynamic on the removal efficiency of a laboratory-scale constructed wetland. *Water Res.* 39 (1), 248-256.
- Winter, M.I. (1985): Entfernung und Umsetzung von Schwefelverbindungen aus Abwasser in Wurzelraumkläranlagen. PhD thesis, Gesamthochschule Kassel.
- Witherspoon, J.; Allen, E.; and Quigley, C. (2004): Modelling to assist in wastewater collection system odour and corrosion potential evaluations. *Water Sci. Technol.* 50 (4), 177-183.
- Winter, M. and Kickuth, R. (1989a): Elimination of sulphur compounds from wastewater by the root zone process-I. Performance of a large-scale purification plant at a textile finishing industry. *Water Res.* 23 (5), 535-546.
- Winter, M. and Kickuth, R. (1989b): Elimination of sulphur compounds from wastewater by the root zone process-II. Mode of formation of sulphur deposits. *Water Res.* 23, 547-560.
- Wissing, F. (1995): Wasserreinigung mit Pflanzen. Ulmer-Verlag. Stuttgart. 207 S.
- Worrall, P.; Peberdy, K.J. and Millett, M.C. (1996): Constructed wetlands and nature conservation. In Proc. 5<sup>th</sup>. Intern. Confe. Wetland systems for water Pollution Control, pp. V/1-1-V/1-9. Vienna, Austria: Universität für Bodenkultur Wien.
- Yavitt, J.B. and Knapp, A.K. (1995): Methane emission to the atmosphere through emergent cattail (*Typha latifolia* L.) plants. *Tellus.* 47B, 521-534.

- Yongsiri, C.; Hvitved-Jacobse, T.; Vollertsen, J. and Tanaka, N. (2003): Introducing the emission process of hydrogen sulfide to a sewer process model (WATS). *Water Sci. Technol.* 47 (4), 85-92.
- Zagorchev, B.N. (1967) : *Analytical Chemistry*. Technika Press, Sofia, p. 568.
- Zartner-Nyilas, G. and Deutsch, P. (2004). *Grenzwerte-Kennzahlen zur Umweltbelastung in Deutschland und der EG. Tabellenwerk*. Ecomed Fachverlag, Landsberg.
- Zehnder, A.J.B. (1988): *Biology of anaerobic microorganisms*. New York. John Wiley and Sons.
- Zhang, T.C. and Lampe, D.G. (1999): Sulfur: Limestone autotrophic denitrification processes for treatment of nitrate-contaminated water: Batch experiments. *Water Res.* 33 (3), 599-608.