

6 Literaturverzeichnis

Arabidopsis Genome Initiative (2000). Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* **408**, 796-815.

Altschul, S. F., Gish, W., Miller, W., Myers, E. W. and Lipman, D. J. (1990). Basic local alignment search tool. *J Mol Biol* **215**, 403-10.

André, B., Hein, C., Grenson, M. and Jauniaux, J. C. (1993). Cloning and expression of the *UGA4* gene coding for the inducible GABA-specific transport protein of *Saccharomyces cerevisiae*. *Mol Gen Genet* **237**, 17-25.

Balconi, C., Rizzi, E., Manzocchi, L., Soave, C. and Motto, M. (1991). Analysis of *in vivo* and *in vitro* grown endosperms of high and low protein strains of maize. *Plant Science* **73**, 1.

Barran, L. R. and Singh, J. (1982). Leucine transport in cells isolated from cold-hardened and nonhardened winter rye. *Plant Physiol.* **69**, 793-797.

Barratt, D. H. P. (1982). Changes during development in the nitrogen, uncombined amino-acid and carbohydrate contents of cotyledons from different cultivars and lines of field bean (*Vicia-Faba* L). *Ann Bot* **49**, 761-768.

Berlin, J. and Mutert, U. (1978). Evidence for distinct amino acid transport systems in cultured tobacco cells. *Z. Naturforsch Teil C: 33*, 641-645.

Bernard, O., Cory, S., Gerondakis, S., Webb, E. and Adams, J. M. (1983). Sequence of the murine and human cellular myc oncogenes and two modes of myc transcription resulting from chromosome translocation in B lymphoid tumours. *Embo J* **2**, 2375-83.

Bick, J. A., Neelam, A., Hall, J. L. and Williams, L. E. (1998). Amino acid carriers of *Ricinus communis* expressed during seedling development: molecular cloning and expression analysis of two putative amino acid transporters, RcAAP1 and RcAAP2. *Plant Mol Biol* **36**, 377-85.

Borden, L. A. (1996). GABA transporter heterogeneity: pharmacology and cellular localization. *Neurochem Int* **29**, 335-56.

Borisjuk, L., Weber, H., Panitz, R., Manteuffel, R. and Wobus, U. (1995). Embryogenesis of *Vicia-faba* L - histodifferentiation and protein storage in relation to seed coat associated invertase. *Plant Physiol* **108**, 95-95.

Borisjuk, L., Rolletschek, H., Radchuk, R., Weschke, W., Wobus, U. and Weber, H. (2004). Seed development and differentiation: a role for metabolic regulation. *Plant Biol (Stuttg)* **6**, 375-86.

Bouché, N., Lacombe, B. and Fromm, H. (2003a). GABA signaling: a conserved and ubiquitous mechanism. *Trends in Cell Biology* **13**, 607.

Bouché, N., Fait, A., Bouchez, D., Moller, S. G. and Fromm, H. (2003b). Mitochondrial succinic-semialdehyde dehydrogenase of the gamma-aminobutyrate shunt is required to restrict levels of reactive oxygen intermediates in plants. *Proc Natl Acad Sci U S A* **100**, 6843-8.

- Bouché, N. and Fromm, H.** (2004). GABA in plants: just a metabolite? *Trends Plant Sci* **9**, 110.
- Bradford, M. M.** (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**, 248-54.
- Breitkreuz, K. E., Shelp, B. J., Fischer, W. N., Schwacke, R. and Rentsch, D.** (1999). Identification and characterization of GABA, proline and quaternary ammonium compound transporters from *Arabidopsis thaliana*. *FEBS Lett* **450**, 280-4.
- Buchanan-Wollaston, V.** (1997). The molecular biology of leaf senescence. *J. Exp. Bot.* **48**, 181-199.
- Buchanan-Wollaston, V., Earl, S., Harrison, E., Mathas, E., Navabpour, S., Page, T. and Pink, D.** (2003). The molecular analysis of leaf senescence - a genomics approach. *Plant Biotechnol J* **1**, 3-22.
- Carter, C., Pan, S., Zouhar, J., Avila, E. L., Girke, T. and Raikhel, N. V.** (2004). The vegetative vacuole proteome of *Arabidopsis thaliana* reveals predicted and unexpected proteins. *Plant Cell* **16**, 3285-3303.
- Chang, H.-C. and Bush, D. R.** (1997). Topology of NAT2, a prototypical example of a new family of amino acid transporters. *J. Biol. Chem.* **272**, 30552-30557.
- Chen, N. H., Reith, M. E. and Quick, M. W.** (2004). Synaptic uptake and beyond: the sodium- and chloride-dependent neurotransmitter transporter family SLC6. *Pflugers Arch* **447**, 519-31.
- Cheruel, J., Jullien, M. and Surdin-Kerjan, Y.** (1979). Amino acid uptake into cultivated mesophyll cells from *Asparagus officinalis* L. *Plant Physiol.* **63**, 621-626.
- Church, G. M. and Gilbert, W.** (1984). Genomic sequencing. *Proc Natl Acad Sci U S A* **81**, 1991-5.
- Clough, S. J. and Bent, A. F.** (1998). Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* **16**, 735-43.
- Cohen, S. N., Chang, A. C. and Hsu, L.** (1972). Nonchromosomal antibiotic resistance in bacteria: genetic transformation of *Escherichia coli* by R-factor DNA. *Proc Natl Acad Sci U S A* **69**, 2110-4.
- Coleman, S. T., Fang, T. K., Rovinsky, S. A., Turano, F. J. and Moye-Rowley, W. S.** (2001). Expression of a glutamate decarboxylase homologue is required for normal oxidative stress tolerance in *Saccharomyces cerevisiae*. *J Biol Chem* **276**, 244-50.
- Daum, G., Gasser, S. M. and Schatz, G.** (1982). Import of proteins into mitochondria. Energy-dependent, two-step processing of the intermembrane space enzyme cytochrome b2 by isolated yeast mitochondria. *J. Biol. Chem.* **257**, 13075-13080.
- De Block, M. a. D., D.** (1996). RNA-RNA *in situ* hybridization using DIG-labeled probes: the effect of high molecular weight polyvinyl alcohol on the alkaline phosphatase indoxyl-nitroblue tetrazolium reaction. *Non-radioactive in situ hybridization Application manual 2nd edn*, 141-145.
- De Jong, A., Koerselman-Kooij, J. W., Schuurmans, J. and Borstlap, A. C.** (1997). The mechanism of amino acid efflux from seed coats of developing pea seeds as revealed by uptake experiments. *Plant Physiol.* **114**, 731-736.

- Dietrich, D., Hammes, U., Thor, K., Suter-Grotemeyer, M., Fluckiger, R., Slusarenko, A. J., Ward, J. M. and Rentsch, D.** (2004). AtPTR1, a plasma membrane peptide transporter expressed during seed germination and in vascular tissue of *Arabidopsis*. *Plant J* **40**, 488-99.
- Dohmen, R. J., Strasser, A. W., Honer, C. B. and Hollenberg, C. P.** (1991). An efficient transformation procedure enabling long-term storage of competent cells of various yeast genera. *Yeast* **7**, 691-2.
- Evan, G. I., Lewis, G. K., Ramsay, G. and Bishop, J. M.** (1985). Isolation of monoclonal antibodies specific for human c-myc proto-oncogene product. *Mol Cell Biol* **5**, 3610-6.
- Fei, Y. J., Ganapathy, V. and Leibach, F. H.** (1998). Molecular and structural features of the proton-coupled oligopeptide transporter superfamily. *Prog Nucleic Acid Res Mol Biol* **58**, 239-61.
- Feller, U. and Keist, M.** (1986). Senescence and nitrogen metabolism in annual plants. In *Fundamental, ecological and agricultural aspects of nitrogen metabolism in higher plants*, eds. H. Lambers J. J. Neeterson and I. Stulen), pp. 219-234: Dordrecht: Martinus Nijhoff Publishers.
- Feller, U. and Fischer, A.** (1994). Nitrogen-metabolism in senescing leaves. *CRC Crit Rev Plant Sci* **13**, 241-273.
- Fischer, W. N., Kwart, M., Hummel, S. and Frommer, W. B.** (1995). Substrate specificity and expression profile of amino acid transporters (AAPs) in *Arabidopsis*. *J Biol Chem* **270**, 16315-20.
- Fischer, W. N., André, B., Rentsch, D., Krolkiewicz, S., Tegeder, M., Breitzkreuz, K. E. and Frommer, W. B.** (1998). Amino acid transport in plants. *Trends Plant Sci* **3**, 188-194.
- Fischer, W. N., Loo, D. D., Koch, W., Ludewig, U., Boorer, K. J., Tegeder, M., Rentsch, D., Wright, E. M. and Frommer, W. B.** (2002). Low and high affinity amino acid H⁺-cotransporters for cellular import of neutral and charged amino acids. *Plant J* **29**, 717-31.
- Frommer, W. B., Hummel, S. and Riesmeier, J. W.** (1993). Expression cloning in yeast of a cDNA encoding a broad specificity amino acid permease from *Arabidopsis thaliana*. *Proc Natl Acad Sci U S A* **90**, 5944-8.
- Frommer, W. B., Kwart, M., Hirner, B., Fischer, W. N., Hummel, S. and Ninnemann, O.** (1994). Transporters for nitrogenous compounds in plants. *Plant Mol Biol* **26**, 1651-70.
- Gamborg, O. L., Miller, R. A. and Ojima, K.** (1968). Nutrient requirements of suspension cultures of soybean root cells. *Experimental Cell Research* **50**, 151-&.
- Gibson, S. I.** (2004). Sugar and phytohormone response pathways: navigating a signalling network. *J. Exp. Bot.* **55**, 253-264.
- Gottesdiener, K. M., Karpinski, B. A., Lindsten, T., Strominger, J. L., Jones, N. H., Thompson, C. B. and Leiden, J. M.** (1988). Isolation and structural characterization of the human 4F2 heavy-chain gene, an inducible gene involved in T-lymphocyte activation. *Mol Cell Biol* **8**, 3809-19.
- Grallath, S., Weimar, T., Meyer, A., Gummy, C., Suter-Grotemeyer, M., Neuhaus, J. M. and Rentsch, D.** (2005). The AtProT family. Compatible solute transporters with similar substrate specificity but differential expression patterns. *Plant Physiol* **137**, 117-26.

- Greenwood, J. S., Helm, M. and Gietl, C.** (2005). Ricinosomes and endosperm transfer cell structure in programmed cell death of the nucellus during Ricinus seed development. *Proc Natl Acad Sci U S A* **102**, 2238-2243.
- Grenson, M., Muyldermans, F., Broman, K. and Vissers, S.** (1987). 4-Aminobutyric acid (GABA) uptake in Baker's yeast *Saccharomyces cerevisiae* is mediated by the general amino acid permease, the proline permease and a GABA-specific permease integrated into the GABA-catabolic pathway. *Life Sci Adv [Ser C]* **6**, 35-39.
- Guo, Y., Cai, Z. and Gan, S.** (2004). Transcriptome of Arabidopsis leaf senescence. *Plant, Cell & Environment* **27**, 521-549.
- Harrington, H. M. and Henke, R. R.** (1981). Amino acid transport into cultured tobacco cells: I. LYSINE TRANSPORT. *Plant Physiol.* **67**, 373-378.
- Heim, U., Weber, H., Bäumllein, H. and Wobus, U.** (1993). A sucrose-synthase gene of *Vicia faba* L.: expression pattern in developing seeds in relation to starch synthesis and metabolic regulation. *Planta* **191**, 394-401.
- Hein, C. and André, B.** (1997). A C-terminal di-leucine motif and nearby sequences are required for NH₄⁽⁺⁾-induced inactivation and degradation of the general amino acid permease, Gap1p, of *Saccharomyces cerevisiae*. *Mol Microbiol* **24**, 607-16.
- Higgins, C. F. and Payne, J. W.** (1978). Peptide transport by germinating barley embryos: Evidence for a single common carrier for di- and oligopeptides. *Planta* **138**, 217.
- Higgins, C. F. and Payne, J. W.** (1980). Transport and utilization of amino acids and peptides by higher plants. In *Microorganisms and Nitrogen Sources*, (ed. J. W. Payne), pp. 609-637: John Wiley & Sons.
- Higgins, C. F. and Payne, J. W.** (1982). Plant peptides. *Encyclop. Plant Physiol.* **14A**, 438-458.
- Higo, K., Ugawa, Y., Iwamoto, M. and Korenaga, T.** (1999). Plant cis-acting regulatory DNA elements (PLACE) database:1999. *Nucl. Acids Res.* **27**, 297-300.
- Hill, J.** (1980). The remobilization of nutrients from leaves. *Journal of Plant Nutrition* **2**, 407-444.
- Hilpert, K., Hansen, G., Wessner, H., Kuttner, G., Welfle, K., Seifert, M. and Hohne, W.** (2001). Anti-c-myc antibody 9E10: epitope key positions and variability characterized using peptide spot synthesis on cellulose. *Protein Eng* **14**, 803-6.
- Himmelblau, E. and Amasino, R. M.** (2001). Nutrients mobilized from leaves of *Arabidopsis thaliana* during leaf senescence. *Journal of Plant Physiology* **158**, 1317-1323.
- Hirner, B., Fischer, W. N., Rentsch, D., Kwart, M. and Frommer, W. B.** (1998). Developmental control of H⁺/amino acid permease gene expression during seed development of Arabidopsis. *Plant J* **14**, 535-44.
- Holsters, M., de Waele, D., Depicker, A., Messens, E., van Montagu, M. and Schell, J.** (1978). Transfection and transformation of *Agrobacterium tumefaciens*. *Mol Gen Genet* **163**, 181-7.
- Hood, E. E., Gelvin, S. B., Melchers, L. S. and Hoekema, A.** (1993). New Agrobacterium helper plasmids for gene-transfer to plants. *Transgenic Research* **2**, 208-218.

- Hörtensteiner, S. and Feller, U.** (2002). Nitrogen metabolism and remobilization during senescence. *J Exp Bot* **53**, 927-37.
- Hosie, A. H., Allaway, D., Galloway, C. S., Dunsby, H. A. and Poole, P. S.** (2002). *Rhizobium leguminosarum* has a second general amino acid permease with unusually broad substrate specificity and high similarity to branched-chain amino acid transporters (Bra/LIV) of the ABC family. *J Bacteriol* **184**, 4071-80.
- Hsu, L., Chiou, T., Chen, L. and Bush, D. R.** (1993). Cloning a plant amino acid transporter by functional complementation of a yeast amino acid transport mutant. *PNAS* **90**, 7441-7445.
- Huang, N. C., Chiang, C. S., Crawford, N. M. and Tsay, Y. F.** (1996). *CHL1* encodes a component of the low-affinity nitrate uptake system in Arabidopsis and shows cell type-specific expression in roots. *Plant Cell* **8**, 2183-91.
- Jack, D. L., Paulsen, I. T. and Saier, M. H.** (2000). The amino acid/polyamine/organocation (APC) superfamily of transporters specific for amino acids, polyamines and organocations. *Microbiology* **146** (Pt 8), 1797-814.
- Jauniaux, J. C., Vandenbol, M., Vissers, S., Broman, K. and Grenson, M.** (1987). Nitrogen catabolite regulation of proline permease in *Saccharomyces cerevisiae*. Cloning of the *PUT4* gene and study of *PUT4* RNA levels in wild-type and mutant strains. *Eur J Biochem* **164**, 601-6.
- Jefferson, R. A., Kavanagh, T. A. and Bevan, M. W.** (1987). GUS fusions: beta-glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *Embo J* **6**, 3901-7.
- Kamachi, K., Yamaya, T., Hayakawa, T., Mae, T. and Ojima, K.** (1992). Changes in cytosolic glutamine synthetase polypeptide and its mRNA in a leaf blade of rice plants during natural senescence. *Plant Physiol.* **98**, 1323-1329.
- Karim, S., Lundh, D., Holmstrom, K. O., Mandal, A. and Pirhonen, M.** (2005). Structural and functional characterization of AtPTR3, a stress-induced peptide transporter of Arabidopsis. *J Mol Model (Online)* **11**, 226-36.
- Kinnersley, A. M. and Turano, F. J.** (2000). Gamma aminobutyric acid (GABA) and plant responses to stress. *Crit. Rev. Plant Sci.* **19**, 479-509.
- Knudson, S.** (1999). Promoter 2.0: for the recognition of PolII promoter sequences. *Bioinformatics* **15**, 356-361.
- Koh, S., Wiles, A. M., Sharp, J. S., Naider, F. R., Becker, J. M. and Stacey, G.** (2002). An oligopeptide transporter gene family in Arabidopsis. *Plant Physiol* **128**, 21-9.
- Kumar, S. and Punekar, N. S.** (1997). The metabolism of 4-aminobutyrate (GABA) in fungi. *Mycol. Res.* **101**, 403-409.
- Kumudini, S., Hume, D. J. and Chu, G.** (2002). Genetic improvement in short-season soybeans: II. Nitrogen accumulation, remobilization, and partitioning. *Crop Sci* **42**, 141-145.
- Künne, C., Lange, M., Funke, T., Miehe, H., Thiel, T., Grosse, I. and Scholz, U.** (2005). CR-EST: a resource for crop ESTs. *Nucleic Acids Res* **33**, D619-21.
- Kwart, M., Hirner, B., Hummel, S. and Frommer, W. B.** (1993). Differential expression of two related amino acid transporters with differing substrate specificity in *Arabidopsis thaliana*. *The Plant Journal* **4**, 993-1002.

- Laemmli, U. K.** (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680-5.
- Lalonde, S., Wipf, D. and Frommer, W. B.** (2004). Transport mechanisms for organic forms of carbon and nitrogen between source and sink. *Annu Rev Plant Biol* **55**, 341-72.
- Lam, E. and Chua, N. H.** (1989). ASF-2: A factor that binds to the cauliflower mosaic virus 35S promoter and a conserved GATA motif in cab promoters. *Plant Cell* **1**, 1147-1156.
- Lee, Y.-H., Foster, J., Chen, J., Voll, L. M., Weber, A. P. M. and Tegeder, M.** (2007). AAP1 transports uncharged amino acids into roots of Arabidopsis. *The Plant Journal* **50**, 305-319.
- Lhullier-Soundélé, A., Munier-Jolain, N. and Ney, B.** (1999). Influence of nitrogen availability on seed nitrogen accumulation in pea. *Crop Sci* **39**, 1741-1748.
- Lohaus, G., Büker, M., Hußmann, M., Soave, C. and Heldt, H.-W.** (1998). Transport of amino acids with special emphasis on the synthesis and transport of asparagine in the Illinois Low Protein and Illinois High Protein strains of maize. *Planta* **205**, 181.
- Lohaus, G. and Moellers, C.** (2000). Phloem transport of amino acids in two *Brassica napus* L. genotypes and one *B. carinata* genotype in relation to their seed protein content. *Planta* **211**, 833.
- Martinez, D. E., Bartoli, C. G., Grbic, V. and Guamet, J. J.** (2007). Vacuolar cysteine proteases of wheat (*Triticum aestivum* L.) are common to leaf senescence induced by different factors. *J. Exp. Bot.* **58**, 1099-1107.
- Martinoia, E., Maeshima, M. and Neuhaus, H. E.** (2007). Vacuolar transporters and their essential role in plant metabolism. *J. Exp. Bot.* **58**, 83-102.
- Masclaux, C., Valadier, M. H., Brugiere, N., Morot-Gaudry, J. F. and Hirel, B.** (2000). Characterization of the sink/source transition in tobacco (*Nicotiana tabacum* L.) shoots in relation to nitrogen management and leaf senescence. *Planta* **211**, 510-8.
- McDaniel, C. N., Holterman, R. K., Bone, R. F. and Wozniak, P. M.** (1982). Amino acid transport in suspension-cultured plant cells: III. COMMON CARRIER SYSTEM FOR THE UPTAKE OF L-ARGININE, L-ASPARTIC ACID, L-HISTIDINE, L-LEUCINE, AND L-PHENYLALANINE. *Plant Physiol.* **69**, 246-249.
- Meyer, A., Eskandari, S., Grallath, S. and Rentsch, D.** (2006). AtGAT1, a high affinity transporter for gamma-aminobutyric acid in *Arabidopsis thaliana*. *J Biol Chem* **281**, 7197-204.
- Meyer, S., Melzer, M., Truernit, E., Hummer, C., Besenbeck, R., Stadler, R. and Sauer, N.** (2000). AtSUC3, a gene encoding a new Arabidopsis sucrose transporter, is expressed in cells adjacent to the vascular tissue and in a carpel cell layer. *Plant J* **24**, 869-82.
- Miranda, M., Borisjuk, L., Tewes, A., Heim, U., Sauer, N., Wobus, U. and Weber, H.** (2001). Amino acid permeases in developing seeds of *Vicia faba* L.: expression precedes storage protein synthesis and is regulated by amino acid supply. *Plant J* **28**, 61-71.
- Miranda, M., Borisjuk, L., Tewes, A., Dietrich, D., Rentsch, D., Weber, H. and Wobus, U.** (2003). Peptide and amino acid transporters are differentially regulated during seed development and germination in faba bean. *Plant Physiol* **132**, 1950-60.
- Montamat, F., Maurousset, L., Tegeder, M., Frommer, W. and Delrot, S.** (1999). Cloning and expression of amino acid transporters from broad bean. *Plant Mol Biol* **41**, 259-68.

- Mossé, J.** (1990). Nitrogen To protein conversion factor for 10 cereals and 6 legumes or oilseeds - a reappraisal of its definition and determination - variation according to species and to seed protein-content. *J. Agric. Food Chem.* **38**, 18-24.
- Murashige, T. and Skoog, F.** (1962). A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiologia Plantarum* **15**, 473-&.
- Nagy, J. I. and Maliga, P.** (1976). Callus induction and plant regeneration from mesophyll protoplasts of *Nicotiana sylvestris*. *Zeitschrift Fur Pflanzenphysiologie* **78**, 453-455.
- Nishiguchi, R., Takanami, M. and Oka, A.** (1987). Characterization and sequence determination of the replicator region in the hairy-root-inducing plasmid pRI44b. *Mol Gen Genet* **206**, 1-8.
- Noodén, L. D., Guiamet, J. J. and John, I.** (1997). Senescence mechanisms. *Physiologia Plantarum* **101**, 746-753.
- Offler, C. E., Nerlich, S. M. and Patrick, J. W.** (1989). Pathway of photosynthate transfer in the developing seed of *Vicia faba* L. Transfer in relation to seed anatomy. *J Exp Bot* **40**, 769-780.
- Offler, C. E., Liet, E. and Sutton, E. G.** (1997). Transfer cell induction in cotyledons of *Vicia faba* L. *Protoplasma* **200**, 51-64.
- Offler, C. E., McCurdy, D. W., Patrick, J. W. and Talbot, M. J.** (2003). Transfer cells: cells specialized for a special purpose. *Annu Rev Plant Biol* **54**, 431-54.
- Okumoto, S., Schmidt, R., Tegeder, M., Fischer, W. N., Rentsch, D., Frommer, W. B. and Koch, W.** (2002). High affinity amino acid transporters specifically expressed in xylem parenchyma and developing seeds of Arabidopsis. *J Biol Chem* **277**, 45338-46.
- Okumoto, S., Koch, W., Tegeder, M., Fischer, W. N., Biehl, A., Leister, D., Stierhof, Y. D. and Frommer, W. B.** (2004). Root phloem-specific expression of the plasma membrane amino acid proton co-transporter AAP3. *J Exp Bot* **55**, 2155-68.
- Owens, D. F. and Kriegstein, A. R.** (2002). Is there more to GABA than synaptic inhibition? *Nat Rev Neurosci* **3**, 715-27.
- Palanivelu, R., Brass, L., Edlund, A. F. and Preuss, D.** (2003). Pollen tube growth and guidance is regulated by POP2, an Arabidopsis gene that controls GABA levels. *Cell* **114**, 47-59.
- Parker, J. M., Guo, D. and Hodges, R. S.** (1986). New hydrophilicity scale derived from high-performance liquid chromatography peptide retention data: correlation of predicted surface residues with antigenicity and X-ray-derived accessible sites. *Biochemistry* **25**, 5425-32.
- Radchuk, V., Borisjuk, L., Radchuk, R., Steinbiss, H.-H., Rolletschek, H., Broeders, S. and Wobus, U.** (2006). *Jekyll* encodes a novel protein involved in the sexual reproduction of barley. *Plant Cell* **18**, 1652-1666.
- Ranki, H., Mendezlozano, J. and Sopanen, T.** (1994). 3 carboxypeptidases occurring in the starchy endosperm of germinating barley-grain have different sites of synthesis. *Physiol. Plant.* **91**, 90-96.
- Rentsch, D., Laloi, M., Rouhara, I., Schmelzer, E., Delrot, S. and Frommer, W. B.** (1995). *NTR1* encodes a high affinity oligopeptide transporter in Arabidopsis. *FEBS Lett* **370**, 264-8.

- Rentsch, D., Boorer, K. J. and Frommer, W. B.** (1998). Structure and function of plasma membrane amino acid, oligopeptide and sucrose transporters from higher plants. *J Membr Biol* **162**, 177-90.
- Robinson, S. P. and Beevers, H.** (1981). Amino acid transport in germinating castor bean seedlings. *Plant Physiol.* **68**, 560-566.
- Rolletschek, H., Hajirezaei, M. R., Wobus, U. and Weber, H.** (2002). Antisense-inhibition of ADP-glucose pyrophosphorylase in *Vicia narbonensis* seeds increases soluble sugars and leads to higher water and nitrogen uptake. *Planta* **214**, 954-64.
- Rolletschek, H., Hosein, F., Miranda, M., Heim, U., Götz, K. P., Schlereth, A., Borisjuk, L., Saalbach, I., Wobus, U. and Weber, H.** (2005). Ectopic expression of an amino acid transporter (*VfAAP1*) in seeds of *Vicia narbonensis* and pea increases storage proteins. *Plant Physiol* **137**, 1236-49.
- Rombauts, S., Dehais, P., Van Montagu, M. and Rouze, P.** (1999). PlantCARE, a plant cis-acting regulatory element database. *Nucleic Acids Res* **27**, 295-6.
- Sagan, M., Ney, B. and Duc, G.** (1993). Plant symbiotic mutants as a tool to analyse nitrogen nutrition and yield relationship in field-growth peas (*Pisum sativum L.*). *Plant and Soil* **153**, 33-45.
- Saharan, K. and Khetarpaul, N.** (1994). Protein quality traits of vegetable and field peas: varietal differences. *Plant Foods Hum Nutr* **45**, 11-22.
- Salon, C., Munier-Jolain, N. G., Duc, G., Voisin, A. S., Grandgirard, D., Larmure, A., Emery, R. J. N. and Ney, B.** (2001). Grain legume seed filling in relation to nitrogen acquisition: A review and prospects with particular reference to pea. *Agronomie* **21**, 539-552.
- Sambrook, J., Fritsch, E. F. and Maniatis, T.** (1989). Molecular Cloning: a laboratory manual. Cold Spring Harbour: Cold Spring Harbour Laboratory Press.
- Schroeder, H. E., Schotz, A. H., Wardley-Richardson, T., Spencer, D. and Higgins, T.** (1993). Transformation and regeneration of two cultivars of pea (*Pisum sativum L.*). *Plant Physiol* **101**, 751-757.
- Schwacke, R., Schneider, A., van der Graaff, E., Fischer, K., Catoni, E., Desimone, M., Frommer, W. B., Flügge, U. I. and Kunze, R.** (2003). ARAMEMNON, a novel database for Arabidopsis integral membrane proteins. *Plant Physiol* **131**, 16-26.
- Shahmuradov, I. A., Gammerman, A. J., Hancock, J. M., Bramley, P. M. and Solovyev, V. V.** (2003). PlantProm: a database of plant promoter sequences. *Nucl. Acids Res.* **31**, 114-117.
- Shaibe, E., Metzger, E. and Halpern, Y. S.** (1985). Metabolic pathway for the utilization of L-arginine, L-ornithine, agmatine, and putrescine as nitrogen sources in *Escherichia coli* K-12. *J Bacteriol* **163**, 933-7.
- Shelp, B. J., Bown, A. W. and McLean, M. D.** (1999). Metabolism and functions of gamma-aminobutyric acid. *Trends Plant Sci* **4**, 446-452.
- Shewry, P. R., Napier, J. A. and Tatham, A. S.** (1995). Seed storage proteins: structures and biosynthesis. *Plant Cell* **7**, 945-56.
- Shimaoka, T., Ohnishi, M., Sazuka, T., Mitsuhashi, N., Hara-Nishimura, I., Shimazaki, K.-I., Maeshima, M., Yokota, A., Tomizawa, K.-I. and Mimura, T.** (2004). Isolation of intact

vacuoles and proteomic analysis of tonoplast from suspension-cultured cells of *Arabidopsis thaliana*. *Plant Cell Physiol.* **45**, 672-683.

Simpson, S. D., Nakashima, K., Narusaka, Y., Seki, M., Shinozaki, K. and Yamaguchi-Shinozaki, K. (2003). Two different novel cis-acting elements of *erd1*, a *clpA* homologous Arabidopsis gene function in induction by dehydration stress and dark-induced senescence. *The Plant Journal* **33**, 259-270.

Smart, C. M. (1994). Gene expression during leaf senescence. *New Phytol* **126**, 419-448.

Snedden, W. A. and Fromm, H. (1999). Regulation of the gamma-aminobutyrate-synthesizing enzyme, glutamate decarboxylase, by calcium-calmodulin: a mechanism for rapid activation in response to stress. In *Plant Responses to Environmental Stresses: From Phytohormones to Genome Reorganization*, (ed. H. R. Lerner), pp. 549-574.

Soldal, T. and Nissen, P. (1978). Multiphasic uptake of amino-acids by barley roots. *Physiol. Plant.* **43**, 181-188.

Solomon, P. S. and Oliver, R. P. (2002). Evidence that gamma-aminobutyric acid is a major nitrogen source during *Cladosporium fulvum* infection of tomato. *Planta* **214**, 414-20.

Song, W., Steiner, H. Y., Zhang, L., Naider, F., Stacey, G. and Becker, J. M. (1996). Cloning of a second Arabidopsis peptide transport gene. *Plant Physiol* **110**, 171-8.

Song, W., Koh, S., Czako, M., Marton, L., Drenkard, E., Becker, J. M. and Stacey, G. (1997). Antisense expression of the peptide transport gene *AtPTR2-B* delays flowering and arrests seed development in transgenic Arabidopsis plants. *Plant Physiol.* **114**, 927-935.

Soudry, E., Ulitzur, S. and Gepstein, S. (2005). Accumulation and remobilization of amino acids during senescence of detached and attached leaves: *in planta* analysis of tryptophan levels by recombinant luminescent bacteria. *J. Exp. Bot.* **56**, 695-702.

Stacey, G., Koh, S., Granger, C. and Becker, J. M. (2002). Peptide transport in plants. *Trends Plant Sci* **7**, 257-63.

Stalberg, K., Ellerstom, M., Ezcurra, I., Ablov, S. and Rask, L. (1996). Disruption of an overlapping E-box/ABRE motif abolished high transcription of the *napA* storage-protein promoter in transgenic *Brassica napus* seeds. *Planta* **199**, 515-9.

Steiner, H.-Y., Naider, F. and Becker, J. M. (1995). The PTR family: a new group of peptide transporters. *Mol Microbiol* **16**, 825-834.

Steward, F. C., Thompson, J. F. and Dent, C. E. (1949). Gamma-aminobutyric acid: a constituent of the potato tuber? *Science* **110**, 439-440.

Streeter, J. G. (1985). Nitrate inhibition of legume nodule growth and activity: I. Long term studies with a continuous supply of nitrate. *Plant Physiol* **77**, 321-324.

Tanaka, J. and Fink, G. R. (1985). The histidine permease gene (*HIP1*) of *Saccharomyces cerevisiae*. *Gene* **38**, 205-14.

Tegeder, M., Wang, X. D., Frommer, W. B., Offler, C. E. and Patrick, J. W. (1999). Sucrose transport into developing seeds of *Pisum sativum* L. *Plant J* **18**, 151-61.

Tegeder, M., Offler, C. E., Frommer, W. B. and Patrick, J. W. (2000). Amino acid transporters are localized to transfer cells of developing pea seeds. *Plant Physiol.* **122**, 319-326.

- Tilsner, J., Kassner, N., Struck, C. and Lohaus, G.** (2005). Amino acid contents and transport in oilseed rape (*Brassica napus* L.) under different nitrogen conditions. *Planta* **221**, 328-38.
- Timmermans, M. C., Maliga, P., Vieira, J. and Messing, J.** (1990). The pFF plasmids: cassettes utilising CaMV sequences for expression of foreign genes in plants. *J Biotechnol* **14**, 333-44.
- Truernit, E. and Sauer, N.** (1995). The promoter of the *Arabidopsis thaliana* *Suc2* Sucrose-H⁺ symporter gene directs expression of beta-glucuronidase to the phloem - evidence for phloem loading and unloading by *Suc2*. *Planta* **196**, 564-570.
- Tsay, Y. F., Chiu, C. C., Tsai, C. B., Ho, C. H. and Hsu, P. K.** (2007). Nitrate transporters and peptide transporters. *FEBS Lett* **581**, 2290-300.
- Ueda, A., Shi, W., Sanmiya, K., Shono, M. and Takabe, T.** (2001). Functional analysis of salt-inducible proline transporter of barley roots. *Plant Cell Physiol.* **42**, 1282-1289.
- Uemura, T., Tomonari, Y., Kashiwagi, K. and Igarashi, K.** (2004). Uptake of GABA and putrescine by UGA4 on the vacuolar membrane in *Saccharomyces cerevisiae*. *Biochem Biophys Res Commun* **315**, 1082-7.
- Ulmasov, T., Hagen, G. and Guilfoyle, T. J.** (1999). Dimerization and DNA binding of auxin response factors. *The Plant Journal* **19**, 309-319.
- Van der Graaff, E., Schwacke, R., Schneider, A., Desimone, M., Flügge, U. I. and Kunze, R.** (2006). Transcription analysis of *Arabidopsis* membrane transporters and hormone pathways during developmental and induced leaf senescence. *Plant Physiol* **141**, 776-92.
- Warren, K. C., Coyne, K. J., Waite, J. H., Cary, S. C.** (1998). Use of methacrylate de-embedding protocols for *in situ* hybridization on semithin plastic sections with multiple detection strategies. *J. Histochem. Cytochem.* **46**, 149-155.
- Waterworth, W. M., West, C. E. and Bray, C. M.** (2000). The barley scutellar peptide transporter: biochemical characterization and localization to the plasma membrane. *J. Exp. Bot.* **51**, 1201-1209.
- Waterworth, W. M., West, C. E. and Bray, C. M.** (2001). The physiology and molecular biology of peptide transport in seeds. *Seed Sci Res* **11**, 275-284.
- Waterworth, W. M., Ashley, M. K., West, C. E., Sunderland, P. A. and Bray, C. M.** (2005). A role for phosphorylation in the regulation of the barley scutellar peptide transporter HvPTR1 by amino acids. *J Exp Bot* **56**, 1545-52.
- Weber, H., Borisjuk, L. and Wobus, U.** (1997a). Sugar import and metabolism during seed development. *Trends Plant Sci* **2**, 169-174.
- Weber, H., Borisjuk, L., Heim, U., Sauer, N. and Wobus, U.** (1997b). A role for sugar transporters during seed development: molecular characterization of a hexose and a sucrose carrier in fava bean seeds. *Plant Cell* **9**, 895-908.
- Weber, H., Heim, U., Golombek, S., Borisjuk, L. and Wobus, U.** (1998a). Assimilate uptake and the regulation of seed development. *Seed Sci Res* **8**, 331-345.
- Weber, H., Heim, U., Golombek, S., Borisjuk, L., Manteuffel, R. and Wobus, U.** (1998b). Expression of a yeast-derived invertase in developing cotyledons of *Vicia narbonensis* alters the carbohydrate state and affects storage functions. *The Plant Journal* **16**, 163-172.

- Weber, H., Rolletschek, H., Heim, U., Golombek, S., Gubatz, S. and Wobus, U.** (2000). Antisense-inhibition of ADP-glucose pyrophosphorylase in developing seeds of *Vicia narbonensis* moderately decreases starch but increases protein content and affects seed maturation. *The Plant Journal* **24**, 33-43.
- Weber, A. and Flügge, U.-I.** (2002). Interaction of cytosolic and plastidic nitrogen metabolism in plants. *J. Exp. Bot.* **53**, 865-874.
- Weber, H., Borisjuk, L. and Wobus, U.** (2005). Molecular physiology of legume seed development. *Annu Rev Plant Biol* **56**, 253-79.
- Weschke, W., Panitz, R., Sauer, N., Wang, Q., Neubohn, B., Weber, H. and Wobus, U.** (2000). Sucrose transport into barley seeds: molecular characterization of two transporters and implications for seed development and starch accumulation. *Plant J* **21**, 455-67.
- Weschke, W., Panitz, R., Gubatz, S., Wang, Q., Radchuk, R., Weber, H. and Wobus, U.** (2003). The role of invertases and hexose transporters in controlling sugar ratios in maternal and filial tissues of barley caryopses during early development. *Plant J* **33**, 395-411.
- West, C. E., Waterworth, W. M., Stephens, S. M., Smith, C. P. and Bray, C. M.** (1998). Cloning and functional characterisation of a peptide transporter expressed in the scutellum of barley grain during the early stages of germination. *Plant J* **15**, 221-9.
- Weston, K., Hall, J. L. and Williams, L. E.** (1994). Characterization of a glutamine/proton cotransporter from *Ricinus communis* roots using isolated plasma-membrane vesicles. *Physiologia Plantarum* **91**, 623-630.
- Wiedemuth, K., Muller, J., Kahlau, A., Amme, S., Mock, H. P., Grzam, A., Hell, R., Egle, K., Beschow, H. and Humbeck, K.** (2005). Successive maturation and senescence of individual leaves during barley whole plant ontogeny reveals temporal and spatial regulation of photosynthetic function in conjunction with C and N metabolism. *Journal of Plant Physiology* **162**, 1226-36.
- Wipf, D., Ludewig, U., Tegeder, M., Rentsch, D., Koch, W. and Frommer, W. B.** (2002). Conservation of amino acid transporters in fungi, plants and animals. *Trends Biochem Sci* **27**, 139-47.
- Wobus, U. and Weber, H.** (1999). Seed maturation: genetic programmes and control signals. *Curr Opin Plant Biol* **2**, 33.
- Wright, K. M., Roberts, A. G., Martens, H. J., Sauer, N. and Oparka, K. J.** (2003). Structural and functional vein maturation in developing tobacco leaves in relation to *AtSUC2* promoter activity. *Plant Physiol* **131**, 1555-65.
- Wu, H. and Cheung, A. Y.** (2000). Programmed cell death in plant reproduction. *Plant Mol Biol* **44**, 267.
- Yamada, K., Matsushima, R., Nishimura, M. and Hara-Nishimura, I.** (2001). A slow maturation of a cysteine protease with a granulin domain in the vacuoles of senescing *Arabidopsis* leaves. *Plant Physiol* **127**, 1626-34.
- Yamagata, M., Matsumoto, S. and Ae, N.** (2001). Possibility of direct acquisition of organic nitrogen by crops *In* N Ae, J Arihara, K Okada, A Srinivasan, eds. *Plant Nutrient Acquisition: New Perspectives*. Springer-Verlag, Tokyo, 399-420.

Yanisch-Perron, C., Vieira, J. and Messing, J. (1985). Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* **33**, 103-119.

Yeung, A. K., Basu, S. K., Wu, S. K., Chu, C., Okamoto, C. T., Hamm-Alvarez, S. F., von Grafenstein, H., Shen, W. C., Kim, K. J., Bolger, M. B. et al. (1995). Molecular identification of a role for tyrosine 167 in the function of the human intestinal proton-coupled dipeptide transporter (hPepT1). *Biochem Biophys Res Commun.* **250**, 103.

Yohannes, E., Barnhart, D. M. and Slonczewski, J. L. (2004). pH-dependent catabolic protein expression during anaerobic growth of *Escherichia coli* K-12. *J Bacteriol* **186**, 192-9.

Yoshida, S. (2003). Molecular regulation of leaf senescence. *Curr Opin Plant Biol* **6**, 79-84.

Youle, R. J. and Huang, A. H. C. (1981). Occurrence of low-molecular weight and high cysteine containing albumin storage proteins in oilseeds of diverse species. *Am. J. Bot.* **68**, 44-48.

Zakharov, A., Giersberg, M., Hosein, F., Melzer, M., Müntz, K. and Saalbach, I. (2004). Seed-specific promoters direct gene expression in non-seed tissue. *J. Exp. Bot.* **55**, 1463-1471.

Zupan, A. L., Trobec, S., Gaberc-Porekar, V. and Menart, V. (2004). High expression of green fluorescent protein in *Pichia pastoris* leads to formation of fluorescent particles. *Recombinant Proteins and Host Cell Physiology* **109**, 115.