

References

- Adam, Z. and Clarke, A. K. (2002). Cutting edge of chloroplast proteolysis. *Trends Plant Sci*, 7(10):451–456.
- Adam, Z. and Ostersetzer, O. (2001). Degradation of unassembled and damaged thylakoid proteins. *Biochem Soc Trans*, 29(Pt 4):427–430.
- Akita, M., Nielsen, E., and Keegstra, K. (1997). Identification of protein transport complexes in the chloroplastic envelope membranes via chemical cross-linking. *J Cell Biol*, 136(5):983–994.
- Akiyama, Y. and Maegawa, S. (2007). Sequence features of substrates required for cleavage by GlpG, an Escherichia coli rhomboid protease. *Mol Microbiol*, 64(4):1028–1037.
- Alami, M., Lüke, I., Deitermann, S., Eisner, G., Koch, H.-G., Brunner, J., and Müller, M. (2003). Differential interactions between a twin-arginine signal peptide and its translocase in Escherichia coli. *Mol Cell*, 12(4):937–946.
- Alder, N. N. and Theg, S. M. (2003a). Energetics of protein transport across biological membranes. a study of the thylakoid Δ ph-dependent/cptat pathway. *Cell*, 112(2):231–242.
- Alder, N. N. and Theg, S. M. (2003b). Energy use by biological protein transport pathways. *Trends Biochem Sci*, 28(8):442–451.
- Allen, S. C. H., Barrett, C. M. L., Ray, N., and Robinson, C. (2002). Essential cytoplasmic domains in the Escherichia coli TatC protein. *J Biol Chem*, 277(12):10362–10366.
- Andersson, H. and von Heijne, G. (1994). Membrane protein topology: effects of $\Delta\mu_{H^+}$ on the translocation of charged residues explain the 'positive inside' rule. *EMBO J*, 13(10):2267–2272.
- Arnon, D. I. (1949). Copper enzymes in isolated chloroplasts. Polyphenoloxidase in beta vulgaris. *Plant Physiol*, 24(1):1–15.

- Bageshwar, U. K. and Musser, S. M. (2007). Two electrical potential dependent steps are required for transport by the Escherichia coli Tat machinery. *J Cell Biol*, 179(1):87–99.
- Barrett, C. M. L., Ray, N., Thomas, J. D., Robinson, C., and Bolhuis, A. (2003). Quantitative export of a reporter protein, GFP, by the twin-arginine translocation pathway in Escherichia coli. *Biochem Biophys Res Commun*, 304(2):279–284.
- Becker, T., Hritz, J., Vogel, M., Caliebe, A., Bukau, B., Soll, J., and Schleiff, E. (2004). Toc12, a novel subunit of the intermembrane space preprotein translocon of chloroplasts. *Mol Biol Cell*, 15(11):5130–5144.
- Behrendt, J., Lindenstrauss, U., and Brüser, T. (2007). The TatBC complex formation suppresses a modular TatB-multimerization in Escherichia coli. *FEBS Lett*, 581(21):4085–4090.
- Behrendt, J., Standar, K., Lindenstrauss, U., and Brüser, T. (2004). Topological studies on the twin-arginine translocase component TatC. *FEMS Microbiol Lett*, 234(2):303–308.
- Berghöfer, J. (1998). *Charakterisierung von Komponenten der plastidären Proteintransport-maschinerie*. PhD thesis, Ludwig-Maximilians-Universität München.
- Berghöfer, J., Karnauchoy, I., Herrmann, R. G., and Klösgen, R. B. (1995). Isolation and characterization of a cDNA encoding the seca protein from spinach chloroplasts. evidence for azide resistance of sec-dependent protein translocation across thylakoid membranes in spinach. *J Biol Chem*, 270(31):18341–18346.
- Berghöfer, J. and Klösgen, R. B. (1996). Isolation and characterization of a cDNA encoding the secy protein from spinach chloroplasts (accession no. z54351) (pgr96-090). *Plant Physiol*, 112:863.
- Berghöfer, J. and Klösgen, R. B. (1999). Two distinct translocation intermediates can be distinguished during protein transport by the TAT (Delta pH) pathway across the thylakoid membrane. *FEBS Lett*, 460(2):328–332.
- Berks, B. C. (1996). A common export pathway for proteins binding complex redox cofactors? *Mol Microbiol*, 22(3):393–404.
- Berks, B. C., Palmer, T., and Sargent, F. (2003). The Tat protein translocation pathway and its role in microbial physiology. *Adv Microb Physiol*, 47:187–254.
- Berks, B. C., Sargent, F., and Palmer, T. (2000). The Tat protein export pathway. *Mol Microbiol*, 35(2):260–274.

- Birnboim, H. C. and Doly, J. (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res*, 7(6):1513–1523.
- Blaudeck, N., Kreutzenbeck, P., Freudl, R., and Sprenger, G. A. (2003). Genetic analysis of pathway specificity during posttranslational protein translocation across the *Escherichia coli* plasma membrane. *J Bacteriol*, 185(9):2811–2819.
- Blaudeck, N., Kreutzenbeck, P., Müller, M., Sprenger, G. A., and Freudl, R. (2005). Isolation and characterization of bifunctional *Escherichia coli* TatA mutant proteins that allow efficient Tat-dependent protein translocation in the absence of TatB. *J Biol Chem*, 280(5):3426–3432.
- Bogsch, E., Brink, S., and Robinson, C. (1997). Pathway specificity for a delta pH-dependent precursor thylakoid lumen protein is governed by a 'Sec-avoidance' motif in the transfer peptide and a 'Sec-incompatible' mature protein. *EMBO J*, 16(13):3851–3859.
- Bolhuis, A., Mathers, J. E., Thomas, J. D., Barrett, C. M., and Robinson, C. (2001). TatB and TatC form a functional and structural unit of the twin-arginine translocase from *Escherichia coli*. *J Biol Chem*, 276(23):20213–20219.
- Braun, N. A., Davis, A. W., and Theg, S. M. (2007). The chloroplast Tat pathway utilizes the transmembrane electric potential as an energy source. *Biophys J*, 93(6):1993–1998.
- Brink, S., Bogsch, E. G., Edwards, W. R., Hynds, P. J., and Robinson, C. (1998). Targeting of thylakoid proteins by the delta pH-driven twin-arginine translocation pathway requires a specific signal in the hydrophobic domain in conjunction with the twin-arginine motif. *FEBS Lett*, 434(3):425–430.
- Brüser, T. and Sanders, C. (2003). An alternative model of the twin arginine translocation system. *Microbiol Res*, 158(1):7–17.
- Buchanan, G., de Leeuw, E., Stanley, N. R., Wexler, M., Berks, B. C., Sargent, F., and Palmer, T. (2002). Functional complexity of the twin-arginine translocase TatC component revealed by site-directed mutagenesis. *Mol Microbiol*, 43(6):1457–1470.
- Chaddock, A. M., Mant, A., Karnauchov, I., Brink, S., Herrmann, R. G., Klösgen, R. B., and Robinson, C. (1995). A new type of signal peptide: central role of a twin-arginine motif in transfer signals for the delta pH-dependent thylakoidal protein translocase. *EMBO J*, 14(12):2715–2722.

- Chan, C. S., Zlomislic, M. R., Tieleman, D. P., and Turner, R. J. (2007). The TatA subunit of Escherichia coli twin-arginine translocase has an N-in topology. *Biochemistry*, 46(25):7396–7404.
- Chanal, A., Santini, C., and Wu, L. (1998). Potential receptor function of three homologous components, TatA, TatB and TatE, of the twin-arginine signal sequence-dependent metalloenzyme translocation pathway in Escherichia coli. *Mol Microbiol*, 30(3):674–676.
- Chen, L. and Tai, P. C. (1989). Effects of inhibitors of membrane signal peptide peptidase on protein translocation into membrane vesicles. *Arch Microbiol*, 153(1):90–94.
- Chupin, V., Killian, J. A., Breg, J., de Jongh, H. H., Boelens, R., Kaptein, R., and de Kruijff, B. (1995). PhoE signal peptide inserts into micelles as a dynamic helix-break-helix structure, which is modulated by the environment. A two-dimensional ¹H NMR study. *Biochemistry*, 34(36):11617–11624.
- Clark, S. A. and Theg, S. M. (1997). A folded protein can be transported across the chloroplast envelope and thylakoid membranes. *Mol Biol Cell*, 8(5):923–934.
- Clausmeyer, S., Klösgen, R. B., and Herrmann, R. G. (1993). Protein import into chloroplasts. The hydrophilic luminal proteins exhibit unexpected import and sorting specificities in spite of structurally conserved transit peptides. *J Biol Chem*, 268(19):13869–13876.
- Cline, K., Ettinger, W. F., and Theg, S. M. (1992). Protein-specific energy requirements for protein transport across or into thylakoid membranes. Two luminal proteins are transported in the absence of ATP. *J Biol Chem*, 267(4):2688–2696.
- Cline, K. and Mori, H. (2001). Thylakoid Delta pH-dependent precursor proteins bind to a cpTatC-Hcf106 complex before Tha4-dependent transport. *J Cell Biol*, 154(4):719–729.
- Creighton, A. M., Hulford, A., Mant, A., Robinson, D., and Robinson, C. (1995). A monomeric, tightly folded stromal intermediate on the delta pH-dependent thylakoidal protein transport pathway. *J Biol Chem*, 270(4):1663–1669.
- Cristobal, S., de Gier, J. W., Nielsen, H., and von Heijne, G. (1999). Competition between Sec- and TAT-dependent protein translocation in Escherichia coli. *EMBO J*, 18(11):2982–2990.

- Dabney-Smith, C., Mori, H., and Cline, K. (2003). Requirement of a Tha4-conserved transmembrane glutamate in thylakoid Tat translocase assembly revealed by biochemical complementation. *J Biol Chem*, 278(44):43027–43033.
- Dabney-Smith, C., Mori, H., and Cline, K. (2006). Oligomers of Tha4 organize at the thylakoid Tat translocase during protein transport. *J Biol Chem*, 281(9):5476–5483.
- de Leeuw, E., Granjon, T., Porcelli, I., Alami, M., Carr, S. B., Müller, M., Sargent, F., Palmer, T., and Berks, B. C. (2002). Oligomeric properties and signal peptide binding by *Escherichia coli* Tat protein transport complexes. *J Mol Biol*, 322(5):1135–1146.
- de Leeuw, E., Porcelli, I., Sargent, F., Palmer, T., and Berks, B. C. (2001). Membrane interactions and self-association of the TatA and TatB components of the twin-arginine translocation pathway. *FEBS Lett*, 506(2):143–148.
- DeLisa, M. P., Lee, P., Palmer, T., and Georgiou, G. (2004). Phage shock protein PspA of *Escherichia coli* relieves saturation of protein export via the Tat pathway. *J Bacteriol*, 186(2):366–373.
- DeLisa, M. P., Samuelson, P., Palmer, T., and Georgiou, G. (2002). Genetic analysis of the twin arginine translocator secretion pathway in bacteria. *J Biol Chem*, 277(33):29825–29831.
- DeLisa, M. P., Tullman, D., and Georgiou, G. (2003). Folding quality control in the export of proteins by the bacterial twin-arginine translocation pathway. *Proc Natl Acad Sci U S A*, 100(10):6115–6120.
- DiCola, A., Bailey, S., and Robinson, C. (2005). The thylakoid delta pH/delta psi are not required for the initial stages of Tat-dependent protein transport in tobacco protoplasts. *J Biol Chem*, 280(50):41165–41170.
- Dilks, K., Rose, R. W., Hartmann, E., and Pohlschröder, M. (2003). Prokaryotic utilization of the twin-arginine translocation pathway: a genomic survey. *J Bacteriol*, 185(4):1478–1483.
- Driessen, A. J., Manting, E. H., and van der Does, C. (2001). The structural basis of protein targeting and translocation in bacteria. *Nat Struct Biol*, 8(6):492–498.
- Dubini, A. and Sargent, F. (2003). Assembly of Tat-dependent [NiFe] hydrogenases: identification of precursor-binding accessory proteins. *FEBS Lett*, 549(1-3):141–146.
- Emanuelsson, O., Nielsen, H., Brunak, S., and von Heijne, G. (2000). Predicting subcellular localization of proteins based on their N-terminal amino acid sequence. *J Mol Biol*, 300(4):1005–1016.

- Endo, T., Kawakami, M., Goto, A., America, T., Weisbeek, P., and Nakai, M. (1994). Chloroplast protein import. Chloroplast envelopes and thylakoids have different abilities to unfold proteins. *Eur J Biochem*, 225(1):403–409.
- Finazzi, G., Chasen, C., Wollman, F.-A., and de Vitry, C. (2003). Thylakoid targeting of Tat passenger proteins shows no delta pH dependence in vivo. *EMBO J*, 22(4):807–815.
- Fincher, V., McCaffery, M., and Cline, K. (1998). Evidence for a loop mechanism of protein transport by the thylakoid Delta pH pathway. *FEBS Lett*, 423(1):66–70.
- Fisher, A. C., Kim, W., and DeLisa, M. P. (2006). Genetic selection for protein solubility enabled by the folding quality control feature of the twin-arginine translocation pathway. *Protein Sci*, 15(3):449–458.
- Frielingsdorf, S. and Klösgen, R. B. (2007). Prerequisites for terminal processing of thylakoidal Tat substrates. *J Biol Chem*, 282(33):24455–24462.
- Gasteiger, E., Gattiker, A., Hoogland, C., Ivanyi, I., Appel, R. D., and Bairoch, A. (2003). ExPASy: The proteomics server for in-depth protein knowledge and analysis. *Nucleic Acids Res*, 31(13):3784–3788.
- Glaser, S. M. and Cumsky, M. G. (1990). A synthetic presequence reversibly inhibits protein import into yeast mitochondria. *J Biol Chem*, 265(15):8808–8816.
- Gohlke, U., Pullan, L., McDevitt, C. A., Porcelli, I., de Leeuw, E., Palmer, T., Saibil, H. R., and Berks, B. C. (2005). The TatA component of the twin-arginine protein transport system forms channel complexes of variable diameter. *Proc Natl Acad Sci U S A*, 102(30):10482–10486.
- Gouffi, K., Gerard, F., Santini, C.-L., and Wu, L.-F. (2004). Dual topology of the Escherichia coli TatA protein. *J Biol Chem*, 279(12):11608–11615.
- Gouffi, K., Santini, C.-L., and Wu, L.-F. (2002). Topology determination and functional analysis of the Escherichia coli TatC protein. *FEBS Lett*, 525(1-3):65–70.
- Gould, S. B., Fan, E., Hempel, F., Maier, U.-G., and Klösgen, R. B. (2007). Translocation of a phycoerythrin alpha subunit across five biological membranes. *J Biol Chem*, 282(41):30295–30302.
- Gould, S. B., Sommer, M. S., Hadfi, K., Zauner, S., Kroth, P. G., and Maier, U.-G. (2006). Protein targeting into the complex plastid of cryptophytes. *J Mol Evol*, 62(6):674–681.

- Graubner, W., Schierhorn, A., and Brüser, T. (2007). DnaK plays a pivotal role in Tat targeting of CueO and functions beside SlyD as a general Tat signal binding chaperone. *J Biol Chem*, 282(10):7116–7124.
- Gutensohn, M., Fan, E., Frielingsdorf, S., Hanner, P., Hou, B., Hust, B., and Klösgen, R. B. (2006). Toc, Tic, Tat et al.: structure and function of protein transport machineries in chloroplasts. *J Plant Physiol*, 163(3):333–347.
- Hatzixanthis, K., Clarke, T. A., Oubrie, A., Richardson, D. J., Turner, R. J., and Sargent, F. (2005). Signal peptide-chaperone interactions on the twin-arginine protein transport pathway. *Proc Natl Acad Sci U S A*, 102(24):8460–8465.
- Heins, L., Mehrle, A., Hemmler, R., Wagner, R., Kchler, M., Hrmann, F., Sveshnikov, D., and Soll, J. (2002). The preprotein conducting channel at the inner envelope membrane of plastids. *EMBO J*, 21(11):2616–2625.
- Herrmann, R. G. (1996). *Photosynthesis research: aspects and perspectives*. Molecular genetics of photosynthesis. Oxford Univ. Press, Oxford, UK.
- Hessa, T., Kim, H., Bihlmaier, K., Lundin, C., Boekel, J., Andersson, H., Nilsson, I., White, S. H., and von Heijne, G. (2005). Recognition of transmembrane helices by the endoplasmic reticulum translocon. *Nature*, 433(7024):377–381.
- Hicks, M. G., de Leeuw, E., Porcelli, I., Buchanan, G., Berks, B. C., and Palmer, T. (2003). The Escherichia coli twin-arginine translocase: conserved residues of TatA and TatB family components involved in protein transport. *FEBS Lett*, 539(1-3):61–67.
- Hoffman, N. E. and Franklin, A. E. (1994). Evidence for a stromal GTP requirement for the integration of a chlorophyll a/b-binding polypeptide into thylakoid membranes. *Plant Physiol*, 105(1):295–304.
- Hou, B. (2005). *Mechanism of pH/TAT-dependent protein transport at the thylakoid membrane*. PhD thesis, Martin-Luther Universität Halle-Wittenberg.
- Hou, B., Frielingsdorf, S., and Klösgen, R. B. (2006). Unassisted membrane insertion as the initial step in Delta pH/Tat-dependent protein transport. *J Mol Biol*, 355(5):957–967.
- Hulford, A., Hazell, L., Mould, R. M., and Robinson, C. (1994). Two distinct mechanisms for the translocation of proteins across the thylakoid membrane, one requiring the presence of a stromal protein factor and nucleotide triphosphates. *J Biol Chem*, 269(5):3251–3256.

- Hynds, P. J., Robinson, D., and Robinson, C. (1998). The sec-independent twin-arginine translocation system can transport both tightly folded and malfolded proteins across the thylakoid membrane. *J Biol Chem*, 273(52):34868–34874.
- Ifuku, K., Nakatsu, T., Kato, H., and Sato, F. (2004). Crystal structure of the PsbP protein of photosystem ii from *Nicotiana tabacum*. *EMBO Rep*, 5(4):362–367.
- Initiative, T. A. G. (2000). Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature*, 408(6814):796–815.
- Izard, J. W., Doughty, M. B., and Kendall, D. A. (1995). Physical and conformational properties of synthetic idealized signal sequences parallel their biological function. *Biochemistry*, 34(31):9904–9912.
- Ize, B., Gerard, F., and Wu, L.-F. (2002). In vivo assessment of the Tat signal peptide specificity in *Escherichia coli*. *Arch Microbiol*, 178(6):548–553.
- Jack, R. L., Buchanan, G., Dubini, A., Hatzixanthis, K., Palmer, T., and Sargent, F. (2004). Coordinating assembly and export of complex bacterial proteins. *EMBO J*, 23(20):3962–3972.
- Jackson-Constan, D., Akita, M., and Keegstra, K. (2001). Molecular chaperones involved in chloroplast protein import. *Biochim Biophys Acta*, 1541(1-2):102–113.
- Jakob, M., K. S. H. P. G. M. S. A. and Klösgen, R. B. (2008). Biochemical characterization of the twin arginine translocation (Tat) machinery from *Arabidopsis Thaliana*. (*Submitted*).
- Jarvis, P. and Robinson, C. (2004). Mechanisms of protein import and routing in chloroplasts. *Curr Biol*, 14(24):R1064–R1077.
- Joly, J. C. and Wickner, W. (1993). The SecA and SecY subunits of translocase are the nearest neighbors of a translocating preprotein, shielding it from phospholipids. *EMBO J*, 12(1):255–263.
- Jongbloed, J. D., Martin, U., Antelmann, H., Hecker, M., Tjalsma, H., Venema, G., Bron, S., van Dijl, J. M., and Müller, J. (2000). TatC is a specificity determinant for protein secretion via the twin-arginine translocation pathway. *J Biol Chem*, 275(52):41350–41357.
- Karnauchov, I. (1998). *Mechanisms of intrachloroplast sorting of integral thylakoid membrane protein*. PhD thesis, Ludwig-Maximilians-Universität München.

- Karnauchov, I., Cai, D., Schmidt, I., Herrmann, R. G., and Klösgen, R. B. (1994). The thylakoid translocation of subunit 3 of photosystem I, the *psaF* gene product, depends on a bipartite transit peptide and proceeds along an azide-sensitive pathway. *J Biol Chem*, 269(52):32871–32878.
- Karnauchov, I., Herrmann, R. G., and Klösgen, R. B. (1997). Transmembrane topology of the Rieske Fe/S protein of the cytochrome b6/f complex from spinach chloroplasts. *FEBS Lett*, 408(2):206–210.
- Keegstra, K. and Cline, K. (1999). Protein import and routing systems of chloroplasts. *Plant Cell*, 11(4):557–570.
- Kessler, F. and Blobel, G. (1996). Interaction of the protein import and folding machineries of the chloroplast. *Proc Natl Acad Sci U S A*, 93(15):7684–7689.
- Kessler, F. and Schnell, D. J. (2004). Chloroplast protein import: solve the GTPase riddle for entry. *Trends Cell Biol*, 14(7):334–338.
- Ki, J. J., Kawarasaki, Y., Gam, J., Harvey, B. R., Iverson, B. L., and Georgiou, G. (2004). A periplasmic fluorescent reporter protein and its application in high-throughput membrane protein topology analysis. *J Mol Biol*, 341(4):901–909.
- Kim, S. J., Robinson, C., and Mant, A. (1998). Sec/SRP-independent insertion of two thylakoid membrane proteins bearing cleavable signal peptides. *FEBS Lett*, 424(1-2):105–108.
- Kinch, L. N., Ginalski, K., and Grishin, N. V. (2006). Site-2 protease regulated intramembrane proteolysis: sequence homologs suggest an ancient signaling cascade. *Protein Sci*, 15(1):84–93.
- Kipping, M., Lilie, H., Lindenstrauss, U., Andreesen, J. R., Griesinger, C., Carlomagno, T., and Brüser, T. (2003). Structural studies on a twin-arginine signal sequence. *FEBS Lett*, 550(1-3):18–22.
- Kirwin, P. M., Elderfield, P. D., and Robinson, C. (1987). Transport of proteins into chloroplasts. Partial purification of a thylakoidal processing peptidase involved in plastocyanin biogenesis. *J Biol Chem*, 262(34):16386–16390.
- Klösgen, R. B., Brock, I. W., Herrmann, R. G., and Robinson, C. (1992). Proton gradient-driven import of the 16 kDa oxygen-evolving complex protein as the full precursor protein by isolated thylakoids. *Plant Mol Biol*, 18(5):1031–1034.
- Kogata, N., Nishio, K., Hirohashi, T., Kikuchi, S., and Nakai, M. (1999). Involvement of a chloroplast homologue of the signal recognition particle receptor protein, FtsY, in protein targeting to thylakoids. *FEBS Lett*, 447(2-3):329–333.

- Kouranov, A., Chen, X., Fuks, B., and Schnell, D. J. (1998). Tic20 and Tic22 are new components of the protein import apparatus at the chloroplast inner envelope membrane. *J Cell Biol*, 143(4):991–1002.
- Kouranov, A. and Schnell, D. J. (1997). Analysis of the interactions of preproteins with the import machinery over the course of protein import into chloroplasts. *J Cell Biol*, 139(7):1677–1685.
- Kuhn, A., Kiefer, D., Khne, C., Zhu, H. Y., Tschantz, W. R., and Dalbey, R. E. (1994). Evidence for a loop-like insertion mechanism of pro-omp a into the inner membrane of escherichia coli. *Eur J Biochem*, 226(3):891–897.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227(5259):680–685.
- Langer, T. (2000). AAA proteases: cellular machines for degrading membrane proteins. *Trends Biochem Sci*, 25(5):247–251.
- Lee, P. A., Buchanan, G., Stanley, N. R., Berks, B. C., and Palmer, T. (2002). Truncation analysis of TatA and TatB defines the minimal functional units required for protein translocation. *J Bacteriol*, 184(21):5871–5879.
- Lemberg, M. K. and Martoglio, B. (2002). Requirements for signal peptide peptidase-catalyzed intramembrane proteolysis. *Mol Cell*, 10(4):735–744.
- Lemberg, M. K. and Martoglio, B. (2004). On the mechanism of SPP-catalysed intramembrane proteolysis: conformational control of peptide bond hydrolysis in the plane of the membrane. *FEBS Lett*, 564(3):213–218.
- Lill, R., Cunningham, K., Brundage, L. A., Ito, K., Oliver, D., and Wickner, W. (1989). SecA protein hydrolyzes ATP and is an essential component of the protein translocation ATPase of Escherichia coli. *EMBO J*, 8(3):961–966.
- Lindahl, M., Spetea, C., Hundal, T., Oppenheim, A. B., Adam, Z., and Andersson, B. (2000). The thylakoid ftsh protease plays a role in the light-induced turnover of the photosystem ii d1 protein. *Plant Cell*, 12(3):419–431.
- Lindahl, M., Tabak, S., Cseke, L., Pichersky, E., Andersson, B., and Adam, Z. (1996). Identification, characterization, and molecular cloning of a homologue of the bacterial FtsH protease in chloroplasts of higher plants. *J Biol Chem*, 271(46):29329–29334.
- Lorkovic, Z. J., Schröder, W. P., Pakrasi, H. B., Irrgang, K. D., Herrmann, R. G., and Oelmüller, R. (1995). Molecular characterization of PsbW, a nuclear-encoded component of the photosystem II reaction center complex in spinach. *Proc Natl Acad Sci U S A*, 92(19):8930–8934.

- Lu, Y. and Beavis, A. D. (1997). Effect of leader peptides on the permeability of mitochondria. *J Biol Chem*, 272(21):13555–13561.
- Lyko, F., Martoglio, B., Jungnickel, B., Rapoport, T. A., and Dobberstein, B. (1995). Signal sequence processing in rough microsomes. *J Biol Chem*, 270(34):19873–19878.
- Maduke, M. and Roise, D. (1993). Import of a mitochondrial presequence into protein-free phospholipid vesicles. *Science*, 260(5106):364–367.
- Maillard, J., Spronk, C. A. E. M., Buchanan, G., Lyall, V., Richardson, D. J., Palmer, T., Vuister, G. W., and Sargent, F. (2007). Structural diversity in twin-arginine signal peptide-binding proteins. *Proc Natl Acad Sci U S A*, 104(40):15641–15646.
- Mangels, D., Mathers, J., Bolhuis, A., and Robinson, C. (2005). The core TatABC complex of the twin-arginine translocase in *Escherichia coli*: TatC drives assembly whereas TatA is essential for stability. *J Mol Biol*, 345(2):415–423.
- Mant, A., Schmidt, I., Herrmann, R. G., Robinson, C., and Klösgen, R. B. (1995). Sec-dependent thylakoid protein translocation. Delta pH requirement is dictated by passenger protein and ATP concentration. *J Biol Chem*, 270(40):23275–23281.
- Manting, E. H. and Driessen, A. J. (2000). *Escherichia coli* translocase: the unravelling of a molecular machine. *Mol Microbiol*, 37(2):226–238.
- Marques, J. P., Dudeck, I., and Klösgen, R. B. (2003). Targeting of EGFP chimeras within chloroplasts. *Mol Genet Genomics*, 269(3):381–387.
- Marques, J. P., Schattat, M. H., Hause, G., Dudeck, I., and Klösgen, R. B. (2004). In vivo transport of folded EGFP by the Delta pH/TAT-dependent pathway in chloroplasts of *Arabidopsis thaliana*. *J Exp Bot*, 55(403):1697–1706.
- Matsuzaki, K., Murase, O., Fujii, N., and Miyajima, K. (1996). An antimicrobial peptide, magainin 2, induced rapid flip-flop of phospholipids coupled with pore formation and peptide translocation. *Biochemistry*, 35(35):11361–11368.
- May, T. and Soll, J. (2000). 14-3-3 proteins form a guidance complex with chloroplast precursor proteins in plants. *Plant Cell*, 12(1):53–64.
- Michl, D., Karnauchov, I., Berghöfer, J., Herrmann, R. G., and Klösgen, R. B. (1999). Phylogenetic transfer of organelle genes to the nucleus can lead to new mechanisms of protein integration into membranes. *Plant J*, 17(1):31–40.
- Michl, D., Robinson, C., Shackleton, J. B., Herrmann, R. G., and Klösgen, R. B. (1994). Targeting of proteins to the thylakoids by bipartite presequences: CFoII is imported by a novel, third pathway. *EMBO J*, 13(6):1310–1317.

- Miguel, M. S., Marrington, R., Rodger, P. M., Rodger, A., and Robinson, C. (2003). An *Escherichia coli* twin-arginine signal peptide switches between helical and unstructured conformations depending on the hydrophobicity of the environment. *Eur J Biochem*, 270(16):3345–3352.
- Miyao, M., F. Y. and Murata, N. (1988). Partial degradation of the 23-kDa protein of the Photosystem II complex of spinach. *Biochim. Biophys. Acta*, 936:465474.
- Molik, S. (2005). *Das plastidäre Rieske Fe/S-Protein: Analyse des Transport- und Assemblierungsprozesses*. PhD thesis, Martin-Luther Universität Halle-Wittenberg.
- Molik, S., Karnauchoy, I., Weidlich, C., Herrmann, R. G., and Klösgen, R. B. (2001). The Rieske Fe/S protein of the cytochrome b6/f complex in chloroplasts: missing link in the evolution of protein transport pathways in chloroplasts? *J Biol Chem*, 276(46):42761–42766.
- Monne, M., Hessa, T., Thissen, L., and von Heijne, G. (2005). Competition between neighboring topogenic signals during membrane protein insertion into the ER. *FEBS J*, 272(1):28–36.
- Moore, M., Harrison, M. S., Peterson, E. C., and Henry, R. (2000). Chloroplast Oxa1p homolog Albino3 is required for post-translational integration of the light harvesting chlorophyllII-binding protein into thylakoid membranes. *J Biol Chem*, 275(3):1529–1532.
- Mori, H. and Cline, K. (2001). Post-translational protein translocation into thylakoids by the Sec and Delta pH-dependent pathways. *Biochim Biophys Acta*, 1541(1-2):80–90.
- Mori, H. and Cline, K. (2002). A twin arginine signal peptide and the pH gradient trigger reversible assembly of the thylakoid [Delta]pH/Tat translocase. *J Cell Biol*, 157(2):205–210.
- Motohashi, R., Nagata, N., Ito, T., Takahashi, S., Hobo, T., Yoshida, S., and Shinozaki, K. (2001). An essential role of a TatC homologue of a Delta pH-dependent protein transporter in thylakoid membrane formation during chloroplast development in *Arabidopsis thaliana*. *Proc Natl Acad Sci U S A*, 98(18):10499–10504.
- Mould, R. M. and Robinson, C. (1991). A proton gradient is required for the transport of two lumenal oxygen-evolving proteins across the thylakoid membrane. *J Biol Chem*, 266(19):12189–12193.

- Mould, R. M., Shackleton, J. B., and Robinson, C. (1991). Transport of proteins into chloroplasts. requirements for the efficient import of two luminal oxygen-evolving complex proteins into isolated thylakoids. *J Biol Chem*, 266(26):17286–17289.
- Müller, M. and Klösgen, R. B. (2005). The Tat pathway in bacteria and chloroplasts (review). *Mol Membr Biol*, 22(1-2):113–121.
- Musser, S. M. and Theg, S. M. (2000). Characterization of the early steps of OE17 precursor transport by the thylakoid Delta pH/Tat machinery. *Eur J Biochem*, 267(9):2588–2598.
- Nakai, M., Goto, A., Nohara, T., Sugita, D., and Endo, T. (1994). Identification of the SecA protein homolog in pea chloroplasts and its possible involvement in thylakoidal protein transport. *J Biol Chem*, 269(50):31338–31341.
- Natale, P., Brüser, T., and Driessen, A. J. M. (2007). Sec- and Tat-mediated protein secretion across the bacterial cytoplasmic membrane-Distinct translocases and mechanisms. *Biochim Biophys Acta [Epub ahead of print]*.
- Nicolay, K., Laterveer, F. D., and van Heerde, W. L. (1994). Effects of amphipathic peptides, including presequences, on the functional integrity of rat liver mitochondrial membranes. *J Bioenerg Biomembr*, 26(3):327–334.
- Nielsen, E., Akita, M., Davila-Aponte, J., and Keegstra, K. (1997). Stable association of chloroplastic precursors with protein translocation complexes that contain proteins from both envelope membranes and a stromal hsp100 molecular chaperone. *EMBO J*, 16(5):935–946.
- Nohara, T., Asai, T., Nakai, M., Sugiura, M., and Endo, T. (1996). Cytochrome f encoded by the chloroplast genome is imported into thylakoids via the SecA-dependent pathway. *Biochem Biophys Res Commun*, 224(2):474–478.
- Novak, P. and Dev, I. K. (1988). Degradation of a signal peptide by protease IV and oligopeptidase A. *J Bacteriol*, 170(11):5067–5075.
- Oates, J., Barrett, C. M. L., Barnett, J. P., Byrne, K. G., Bolhuis, A., and Robinson, C. (2005). The Escherichia coli twin-arginine translocation apparatus incorporates a distinct form of TatABC complex, spectrum of modular TatA complexes and minor TatAB complex. *J Mol Biol*, 346(1):295–305.
- Oates, J., Mathers, J., Mangels, D., Kühlbrandt, W., Robinson, C., and Model, K. (2003). Consensus structural features of purified bacterial TatABC complexes. *J Mol Biol*, 330(2):277–286.

- Oresnik, I. J., Ladner, C. L., and Turner, R. J. (2001). Identification of a twin-arginine leader-binding protein. *Mol Microbiol*, 40(2):323–331.
- Palmer, T., Sargent, F., and Berks, B. C. (2005). Export of complex cofactor-containing proteins by the bacterial Tat pathway. *Trends Microbiol*, 13(4):175–180.
- Pasch, J. C., Nickelsen, J., and Schünemann, D. (2005). The yeast split-ubiquitin system to study chloroplast membrane protein interactions. *Appl Microbiol Biotechnol*, 69(4):440–447.
- Perez-Rodriguez, R., Fisher, A. C., Perlmutter, J. D., Hicks, M. G., Chanal, A., Santini, C.-L., Wu, L.-F., Palmer, T., and DeLisa, M. P. (2007). An essential role for the DnaK molecular chaperone in stabilizing over-expressed substrate proteins of the bacterial twin-arginine translocation pathway. *J Mol Biol*, 367(3):715–730.
- Pfanner, N., Rassow, J., van der Klei, I. J., and Neupert, W. (1992). A dynamic model of the mitochondrial protein import machinery. *Cell*, 68(6):999–1002.
- Porcelli, I., de Leeuw, E., Wallis, R., van den Brink-van der Laan, E., de Kruijff, B., Wallace, B. A., Palmer, T., and Berks, B. C. (2002). Characterization and membrane assembly of the TatA component of the Escherichia coli twin-arginine protein transport system. *Biochemistry*, 41(46):13690–13697.
- Preiss, S., Schrader, S., and Johanningmeier, U. (2001). Rapid, atp-dependent degradation of a truncated d1 protein in the chloroplast. *Eur J Biochem*, 268(16):4562–4569.
- Qbadou, S., Becker, T., Bionda, T., Reger, K., Ruprecht, M., Soll, J., and Schleiff, E. (2007). Toc64—a preprotein-receptor at the outer membrane with bipartite function. *J Mol Biol*, 367(5):1330–1346.
- Qbadou, S., Becker, T., Mirus, O., Tews, I., Soll, J., and Schleiff, E. (2006). The molecular chaperone Hsp90 delivers precursor proteins to the chloroplast import receptor Toc64. *EMBO J*, 25(9):1836–1847.
- Rawlings, N. D., Morton, F. R., and Barrett, A. J. (2006). MEROPS: the peptidase database. *Nucleic Acids Res*, 34(Database issue):D270–D272.
- Richter, S. and Lamppa, G. K. (1999). Stromal processing peptidase binds transit peptides and initiates their ATP-dependent turnover in chloroplasts. *J Cell Biol*, 147(1):33–44.
- Richter, S. and Lamppa, G. K. (2002). Determinants for removal and degradation of transit peptides of chloroplast precursor proteins. *J Biol Chem*, 277(46):43888–43894.

- Rizo, J., Blanco, F. J., Kobe, B., Bruch, M. D., and Gierasch, L. M. (1993). Conformational behavior of Escherichia coli OmpA signal peptides in membrane mimetic environments. *Biochemistry*, 32(18):4881–4894.
- Robinson, C. and Bolhuis, A. (2004). Tat-dependent protein targeting in prokaryotes and chloroplasts. *Biochim Biophys Acta*, 1694(1-3):135–147.
- Sambrook, J., F. E. F. M. T. (1989). *Molecular Cloning: A Laboratory Manual*[M]. 2nd ed. New York: Cold Spring Harbor Laboratory Press.
- Sanger, F., Nicklen, S., and Coulson, A. R. (1977). DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci U S A*, 74(12):5463–5467.
- Santini, C. L., Ize, B., Chanal, A., Müller, M., Giordano, G., and Wu, L. F. (1998). A novel sec-independent periplasmic protein translocation pathway in Escherichia coli. *EMBO J*, 17(1):101–112.
- Sargent, F., Berks, B. C., and Palmer, T. (2006). Pathfinders and trailblazers: a prokaryotic targeting system for transport of folded proteins. *FEMS Microbiol Lett*, 254(2):198–207.
- Sargent, F., Gohlke, U., de Leeuw, E., Stanley, N. R., Palmer, T., Saibil, H. R., and Berks, B. C. (2001). Purified components of the Escherichia coli Tat protein transport system form a double-layered ring structure. *Eur J Biochem*, 268(12):3361–3367.
- Sargent, F., Stanley, N. R., Berks, B. C., and Palmer, T. (1999). Sec-independent protein translocation in Escherichia coli. A distinct and pivotal role for the TatB protein. *J Biol Chem*, 274(51):36073–36082.
- Schägger, H., Cramer, W. A., and von Jagow, G. (1994). Analysis of molecular masses and oligomeric states of protein complexes by blue native electrophoresis and isolation of membrane protein complexes by two-dimensional native electrophoresis. *Anal Biochem*, 217(2):220–230.
- Schägger, H. and von Jagow, G. (1991). Blue native electrophoresis for isolation of membrane protein complexes in enzymatically active form. *Anal Biochem*, 199(2):223–231.
- Schatz, G. and Dobberstein, B. (1996). Common principles of protein translocation across membranes. *Science*, 271(5255):1519–1526.
- Schleiff, E., Jelic, M., and Soll, J. (2003). A GTP-driven motor moves proteins across the outer envelope of chloroplasts. *Proc Natl Acad Sci U S A*, 100(8):4604–4609.

- Schleiff, E. and Klösgen, R. B. (2001). Without a little help from 'my' friends: direct insertion of proteins into chloroplast membranes? *Biochim Biophys Acta*, 1541(1-2):22–33.
- Schnell, D. J., Kessler, F., and Blobel, G. (1994). Isolation of components of the chloroplast protein import machinery. *Science*, 266(5187):1007–1012.
- Schülke, N., Sepuri, N. B., and Pain, D. (1997). In vivo zippering of inner and outer mitochondrial membranes by a stable translocation intermediate. *Proc Natl Acad Sci U S A*, 94(14):7314–7319.
- Schünemann, D. (2004). Structure and function of the chloroplast signal recognition particle. *Curr Genet*, 44(6):295–304.
- Settles, A. M., Yonetani, A., Baron, A., Bush, D. R., Cline, K., and Martienssen, R. (1997). Sec-independent protein translocation by the maize Hcf106 protein. *Science*, 278(5342):1467–1470.
- Shackleton, J. B. and Robinson, C. (1991). Transport of proteins into chloroplasts. The thylakoidal processing peptidase is a signal-type peptidase with stringent substrate requirements at the -3 and -1 positions. *J Biol Chem*, 266(19):12152–12156.
- Shanmugham, A., Sang, H. W. W. F., Bollen, Y. J. M., and Lill, H. (2006). Membrane binding of twin arginine preproteins as an early step in translocation. *Biochemistry*, 45(7):2243–2249.
- Soll, J. and Schleiff, E. (2004). Protein import into chloroplasts. *Nat Rev Mol Cell Biol*, 5(3):198–208.
- Stanley, N. R., Palmer, T., and Berks, B. C. (2000). The twin arginine consensus motif of Tat signal peptides is involved in Sec-independent protein targeting in *Escherichia coli*. *J Biol Chem*, 275(16):11591–11596.
- Stengel, A., Soll, J., and Bölter, B. (2007). Protein import into chloroplasts: new aspects of a well-known topic. *Biol Chem*, 388(8):765–772.
- Summer, E. J., Mori, H., Settles, A. M., and Cline, K. (2000). The thylakoid delta pH-dependent pathway machinery facilitates RR-independent N-tail protein integration. *J Biol Chem*, 275(31):23483–23490.
- Sun, C., Rusch, S. L., Kim, J., and Kendall, D. A. (2007). Chloroplast SecA and *Escherichia coli* SecA have distinct lipid and signal peptide preferences. *J Bacteriol*, 189(3):1171–1175.

- Sundberg, E., Slagter, J. G., Fridborg, I., Cleary, S. P., Robinson, C., and Coupland, G. (1997). ALBINO3, an Arabidopsis nuclear gene essential for chloroplast differentiation, encodes a chloroplast protein that shows homology to proteins present in bacterial membranes and yeast mitochondria. *Plant Cell*, 9(5):717–730.
- Theg, S. M., Cline, K., Finazzi, G., and Wollman, F.-A. (2005). The energetics of the chloroplast Tat protein transport pathway revisited. *Trends Plant Sci*, 10(4):153–154.
- Thompson, S. J., Kim, S. J., and Robinson, C. (1998). Sec-independent insertion of thylakoid membrane proteins. Analysis of insertion forces and identification of a loop intermediate involving the signal peptide. *J Biol Chem*, 273(30):18979–18983.
- van Dijl, J. M., Braun, P. G., Robinson, C., Quax, W. J., Antelmann, H., Hecker, M., Müller, J., Tjalsma, H., Bron, S., and Jongbloed, J. D. H. (2002). Functional genomic analysis of the *Bacillus subtilis* Tat pathway for protein secretion. *J Biotechnol*, 98(2-3):243–254.
- van Klompenburg, W. and de Kruijff, B. (1998). The role of anionic lipids in protein insertion and translocation in bacterial membranes. *J Membr Biol*, 162(1):1–7.
- VanderVere, P. S., Bennett, T. M., Oblong, J. E., and Lamppa, G. K. (1995). A chloroplast processing enzyme involved in precursor maturation shares a zinc-binding motif with a recently recognized family of metalloendopeptidases. *Proc Natl Acad Sci U S A*, 92(16):7177–7181.
- Vestweber, D. and Schatz, G. (1988). A chimeric mitochondrial precursor protein with internal disulfide bridges blocks import of authentic precursors into mitochondria and allows quantitation of import sites. *J Cell Biol*, 107(6 Pt 1):2037–2043.
- von Heijne, G. (1998). Life and death of a signal peptide. *Nature*, 396(6707):111, 113.
- Walker, M. B., Roy, L. M., Coleman, E., Voelker, R., and Barkan, A. (1999). The maize *tha4* gene functions in sec-independent protein transport in chloroplasts and is related to *hcf106*, *tatA*, and *tatB*. *J Cell Biol*, 147(2):267–276.
- Weihofen, A., Binns, K., Lemberg, M. K., Ashman, K., and Martoglio, B. (2002). Identification of signal peptide peptidase, a presenilin-type aspartic protease. *Science*, 296(5576):2215–2218.
- Wieprecht, T., Apostolov, O., Beyermann, M., and Seelig, J. (2000). Interaction of a mitochondrial presequence with lipid membranes: role of helix formation for membrane binding and perturbation. *Biochemistry*, 39(50):15297–15305.

- Wolfe, M. S. and Kopan, R. (2004). Intramembrane proteolysis: theme and variations. *Science*, 305(5687):1119–1123.
- Yahr, T. L. and Wickner, W. T. (2001). Functional reconstitution of bacterial Tat translocation in vitro. *EMBO J*, 20(10):2472–2479.
- Yang, F., Moss, L. G., and Phillips, G. N. (1996). The molecular structure of green fluorescent protein. *Nat Biotechnol*, 14(10):1246–1251.
- Ye, J., Dav, U. P., Grishin, N. V., Goldstein, J. L., and Brown, M. S. (2000). Asparagine-proline sequence within membrane-spanning segment of SREBP triggers intramembrane cleavage by site-2 protease. *Proc Natl Acad Sci U S A*, 97(10):5123–5128.
- Yen, M.-R., Tseng, Y.-H., Nguyen, E. H., Wu, L.-F., and Saier, M. H. (2002). Sequence and phylogenetic analyses of the twin-arginine targeting (Tat) protein export system. *Arch Microbiol*, 177(6):441–450.
- Yuan, J. and Cline, K. (1994). Plastocyanin and the 33-kDa subunit of the oxygen-evolving complex are transported into thylakoids with similar requirements as predicted from pathway specificity. *J Biol Chem*, 269(28):18463–18467.
- Yuan, J., Kight, A., Goforth, R. L., Moore, M., Peterson, E. C., Sakon, J., and Henry, R. (2002). ATP stimulates signal recognition particle (SRP)/FtsY-supported protein integration in chloroplasts. *J Biol Chem*, 277(35):32400–32404.
- Zardeneta, G. and Horowitz, P. M. (1992). Analysis of the perturbation of phospholipid model membranes by rhodanese and its presequence. *J Biol Chem*, 267(34):24193–24198.