

7 Literaturverzeichnis

- 1 Abbott KC, Glanton CW, Trespalacios FC, Oliver DK, Ortiz MI, Agodoa LY: Body mass index, dialysis modality and survival: Analysis of the United States Renal Data System Dialysis Morbidity and Mortality Wave II Study. *Kidney Int* 65 (2004) 597-605
- 2 Anker SD, Egerer KR, Volk HD, Kox WJ, Pool-Wilson PA, Coats AJ: Elevated soluble CD 14 receptors and altered cytokines in chronic heart failure. *Am J Cardiol* 79 (1997) 1426-1430
- 3 Anker SD, Volterrani M, Egerer KR, Felton CV, Kox WJ, Poole-Wilson PA, Coats AJ: Tumour necrosis factor alpha as a predictor of impaired peak leg blood flow in patients with chronic heart failure. *QJM* 91 (1998) 199-203
- 4 Beddhu S, Pappas LM, Ramkumar N, Samore M: Effects of body size and body composition on survival in hemodialysis patients. *J Am Soc Nephrol* 14 (2003) 2366-2372
- 5 Behring-Nephelometer: Bedienungsanleitung-Version 2.0
- 6 Bergström J: Metabolic acidosis and nutrition in dialysis patients. *Blood Purif* 13 (1995) 361-367
- 7 Berneis K, Iseli-Schaub J, Garbani E, Meier R, Kiss D: Effects of intradialytic parenteral nutrition in chronic haemodialysis patients with malnutrition: A pilot study *Wien. Klin Wochenschr* 111/21 (1999) 876-881
- 8 Bernstein LH, Leukhardt-Fairfield CJ, Pleban W et al: Usefulness of data on albumin and prealbumin concentrations in determining effectiveness of nutritional support. *Clin Chem* 35 (1989) 271-274
- 9 Bethea M, Forman DT: β_2 -Mikroglobulin: its significance and clinical usefulness. *Ann Clin Lab Sci* 20 (1990) 163-168

- 10 Boege F: Harnproteine. In: Thomas L (Hrsg): Labor und Diagnose. TH-Books Verlagsgesellschaft, Frankfurt, 1998, S. 393-410
- 11 Brandtzaeg P, Kierulf P, Gaustad P, et al: Plasma endotoxin as a predictor of multiple organ failure and death in systemic meningococcal disease. *J Infect Dis* 159 (1989) 195-204
- 12 Busse R: Gefäßsystem und Kreislaufregulation. In: Schmidt RF, Thews G (Hrsg): Physiologie des Menschen. Springer, Berlin, Heidelberg, New York (usw.), 1995, S. 498-561
- 13 Casl MT, Bulatovic G, Orlic P, Salbljar-Matovinovic SM: The diagnostic capacity of serum amyloid A protein for early recognition of kidney allograft rejection. *Nephrol Dial Transplant* 10 (1995) 1901-1904
- 14 Charra B, Laurent G, Chazot C, Calemard E, Terrat JC, Vanel T, Jean G, Ruffet M: Clinical assessment of dry weight. *Nephrol Dial Transplant* 11 [Suppl 2] (1996) 16-19
- 15 Chazot C, Charra B, Laurent G, Didier C, Vo Van C, Terrat JC, Calemard E, Vanel T, Ruffet M: Interdialysis blood pressure control by long haemodialysis sessions. *Nephrol Dial Transplant* 10 [Suppl 6] (1995) 831-837
- 16 Cheriex EC, Leunissen KML, Janssen JHA, Mooy JMV, van Hooff JP: Echography of the inferior vena cava is a simple and reliable tool for estimation of "dry weight" in haemodialysis patients. *Nephrol Dial Transplant* 4 (1989) 563-568
- 17 Chertow GM, Lowrie EG, Wilmore DW, Gonzalez J, Lew NL, Ling J, Leboff MS, Gottlieb MN, Huang W, Zebrowski B: Nutritional assessment with bioelectrical impedance analysis in maintenance hemodialysis patients. *J Am Soc Nephrol* 6 (1995) 75-81

- 18 Detsky AS, Baker JP, Mendelson RA, Wolman SL, Wesson DE, Jeejeebhoy KN: Evaluating the accuracy of nutritional assessment techniques applied to hospitalized patients: methodology and comparisons. *J Parenter Enteral Nutr* 8 (1984) 153-159
- 19 Don C, Burns KD, Levine DZ: Body fluid volume status in hemodialysis patients: The value of the chest Radiograph. *J Can Assoc Radiol* 41 (1990) 123-126
- 20 Fiedler R, Saile P, Steudte R, Mall M, Osten B: Klinischer Ernährungsstatus bei Hämodialysepatienten im Vergleich zu laborchemischen Parametern und zur bioelektrischen Impedanz. *Nieren- und Hochdruckkrankheiten* 35 (2006) 178-184
- 21 Fleischman E, Teal N, Dudley J, May W, Bower JD, Salahudeen A: Overweight in hemodialysis and its influence on mortality and hospital stay: Data from 1,346 patients. *Kidney Int* 55 (1999) 1560-1567
- 22 Friedl W, Mair J, Thomas S, Pichler M, Puschendorf B: Natriuretic peptides and cyclic guanosine monophosphate in asymptomatic and symptomatic left ventricular dysfunction. *Heart* 76 (1996) 129-136
- 23 Fießl HS, Middeke MRF: Anamnese und klinische Untersuchung. Georg Thieme, Stuttgart, 1998, S. 172-175
- 24 Fießl HS, Middeke MRF: Anamnese und klinische Untersuchung. Georg Thieme, Stuttgart, 1998, S. 334-335
- 25 Gladisch R: Einführung in die sonographische Diagnostik. In: Gladisch R. (Hrsg): *Praxis der abdominalen Ultraschalldiagnostik*. Schattauer, Stuttgart, 1992, S. 2-28
- 26 Gudivaka R, Schoeller D, Kushner RF: Effect of skin temperature on multi-frequency bioelectrical impedance analysis. *J Appl Physiol* 81 (1996) 838-845

- 27 Guigoz Y, Vellas B, Garry PJ: Mini nutritional assessment: a practical assessment tool for grading the nutritional state of elderly patients: Facts Res Gerontol [Suppl 2] (1994) 15-59
- 28 Hakim RM, Lazarus JM: Initiation of dialysis. J Am Soc Nephrol 6 (1995) 1319-1328
- 29 Hakim RM, Levin N: Malnutrition in Hemodialysis Patients. Am J Kidney Dis 21 (1993) 125-137
- 30 Hartmann A, Eide TC, Fauchald P, Bentdal Q, Herbert J, Gallimore JR, Pepys MB : Serum amyloid A protein is a clinically useful indicator of acute renal allograft rejection. Nephrol Dial Transplant 12 (1997) 161-166
- 31 Hauck G: Pathophysiologie der Ödembildung. Perfusion. Springer, 1990, S. 1-5
- 32 Haug C, Metzeler A, Steffgen J, Kochs M, Hombach V, Grunert A: Increased brain natriuretic peptide and atrial natriuretic peptide plasma concentrations in dialysis dependent chronic renal failure and in patients with elevated left ventricular filling pressure. Clin Investig 72 (1994) 430-434
- 33 Henderson LW, Thomson GE, Waterhouse K, McDonald HP, Friedman EA, Kinet JP, Soyeur D, Balland N, Saint-Remy M, Collignon P, Godon JP: Symptomatic hypotension during hemodialysis. Kidney Int 17 (1980) 571-576
- 34 Hoenich NA, Levin NW: Can technology solve the clinical problem of „dry weight“? Nephrol Dial Transplant 18 (2003) 647-650
- 35 Hofmann W, Guder WG: A diagnostic programme for quantitative analysis of proteinuria. J Clin Chem Clin Biochem 27 (1989) 589-600
- 36 Hogarth MB, Gallimore R, Savage P, Palmer AJ, Starr, JM, Bulpitt CJ, Pepys MB: Acute phase proteins, C-reactive protein and serum amyloid A protein, as prognostic markers in the elderly inpatient. Age Ageing 26 (1997) 153-158

- 37 Igarashi Y, Suzuki H, Imafuku T: Hypertension in patients on chronic hemodialysis: the role of the renin-angiotensin system. *Jap Circulat J* 51 (1987) 479-484
- 38 Ikizler TA, Greene JH, Wingard RL, Parker RA, Hakim RM: Spontaneous dietary protein intake during progression of chronic renal failure. *J Am Soc Nephrol* 6 (1995) 1386-1391
- 39 Ishizaka Y, Yamamoto Y, Fukunaga T, Yokota N, Kida O, Kitamura K, Kangawa K, Minamino N, Matsuo H, Eto T: Plasma concentration of human brain natriuretic peptide in patients on hemodialysis. *Am J Kidney Dis* 24 (1994) 461-472
- 40 Jaeger JQ, Mehta RL: Assessment of dry weight in hemodialysis. *J Am Soc Nephrol* 10 (1999) 392-403
- 41 John E: Morley *Proceedings of the Nutrition Survey* 57 (1998) 587-592
- 42 Johnson DW, Herzig KA, Purdie DM, Chang W, Brown AM, Rigby RJ, Campbell, SB, Nicol DL, Hawley, CM: Is obesity a favorable prognostic factor in peritoneal dialysis patients? *Perit Dial Int* 20 (2000) 715-721
- 43 Kalantar-Zadeh K, Block G, Humphreys MH, Kopple JD : Reverse epidemiology of cardiovascular risk factors in maintenance dialysis patients. *Kidney Int* 63 (2003) 793-808
- 44 Kalantar-Zadeh K, Kopple JD, Block G, Humphreys MH: A malnutrition-inflammation score is correlated with morbidity and mortality in maintenance hemodialysis patients. *Am J Kidney Dis* 38 (2001) 1251-1263
- 45 Kalantar-Zadeh K, Kopple JD, Humphreys MH, Block G: Comparing outcome predictability of markers of malnutrition-inflammation complex syndrom in haemodialysis patients. *Nephrol Dial Transplant* 19 (2004) 1507-1519

- 46 Kathenelson R, Perel A, Berkenstadt H, Preisman S, Kogan S, Sternik L, Segal E: Accuracy of transpulmonary thermodilution versus gravimetric measurement of extravascular lung water. *Critical Care Medicine* 32 (2004) 1550-1554
- 47 Katzarski KS, Nisell J, Randmaa I, Danielsson A, Freyschuss U, Bergström J: A critical evaluation of ultrasound measurement of inferior vena cava diameter in assessing dry weight in normotensive and hypertensive hemodialysis patients. *Am J Kidney Diseases* 30 (1997) 459-465
- 48 Kaysen GA, Kumar V: Inflammation in ESRD: causes and potential consequences. *J Ren Nutr* 13 (2003) 158-160
- 49 Kondrup J, Rasmussen HH, Hamburg O, Stanga Z, Ad Hoc ESPEN Working Group: Nutritional Risk Screening (NRS 2002): A new method based on analysis of controlled clinical trials. *Clin Nutr* 22 (2003) 321-336
- 50 Krahenbuhl B, Restellini A, Frangos A: Peripheral venous pulsatility detected by Doppler method for diagnosis of right heart failure. *Cardiology* 71 (1984) 173-176
- 51 Krause I, Birk E, Davidovits M, Cleper R, Blieden L, Pinhas L, Gamzo Z, Eisenstein B: Inferior vena cava diameter: a useful method for estimation of fluid status in children on haemodialysis. *Nephrol Dial Transplant* 16 (2001) 1203-1206
- 52 Krautzig S, Janssen U, Koch KM, Granolleras C, Shaldon S: Dietary salt restriction and reduction of dialysate sodium to control hypertension in maintenance haemodialysis patients. *Nephrol Dial Transplant* 13 (1998) 552-553
- 53 Kuhn C, Kuhn A, Rykow K, Osten B: Extravascular lung water index: a new method to determine dry weight in chronic hemodialysis patients. *Hemodial Int* 10 (2006) 68-72
- 54 Kusaba T, Yamagucci K, Oda H: Echography of the inferior vena cava for estimating fluid removal from patients undergoing hemodialysis. *Jpn J Nephrol* 38 (1996) 119-123

- 55 Kushner RF, de Vries PM, Gudivaka R: Use of bioelectrical impedance analysis measurements in the clinical management of patients undergoing dialysis. *Am J Clin Nutr* 64 [Suppl 3] (1996) 503-509
- 56 Lauster F, Fülle HJ, Gerzer R, Schiffel H: The postdialytic plasma cyclic guanosine 3`5`- monophosphate level as a measure of fluid overload in chronic hemodialysis. *J Am Soc Nephrol* 2 (1992) 1451-1454
- 57 Lee SW, Song JH, Kim GA, Lim HJ, Kim MJ: Plasma brain natriuretic peptide concentration on assessment of hydration status in hemodialysis patient. *Am J Kidney Dis* 41 (2003) 1257-1266
- 58 Lowrie EG, Lew NL: Death risk in hemodialysis patients: the predictive value of commonly measured variables and an evaluation of death rate differences between facilities. *Am. J Kidney Dis* 15 (1990) 458-482
- 59 Löser C, Keymling M: Praxis der enteralen Ernährung: Indikation-Technik-Nachsorge. Georg Thieme, Stuttgart, 2001
- 60 Lukaski HC: Methods for assessment of human body composition: traditional and new. *Am J Clin Nutr* 50 (1987) 537-543
- 61 Malle E, De Beer FC: Human serum amyloid A (SAA) protein: a prominent acute-phase reactant for clinical practice. *Eur J Clin Invest* 26 (1996) 427-435
- 62 Mandelbaum A, Link A, Wambach G, Ritz E: Vena-cava-Sonographie zur Beurteilung des Hydratationszustandes bei Niereninsuffizienz. *Dtsch Med Wschr* 118 (1993) 1309-1315
- 63 Mandelbaum A, Ritz E: Vena cava diameter measurement for estimation of dry weight in haemodialysis patients. *Nephrol Dial Transplant* 11 [Suppl 2] (1996) 24-27

- 64 Marhaug G, Dowton SB: Serumamyloid A: apolipoprotein and precursor of AA amyloid. In: Husby G (Ed): Reactive Amyloidosis and the Acute Phase Response. Ballieres Clinical Rheumatology Vol 8 Balliere Tindall, London, 1994, S. 533-537
- 65 Martinez-Maldonado M: Hypertension in end-stage renal disease. *Kidney Int [Suppl 68]* (1998) 67-72
- 66 Müller MJ, Selberg O: Ernährungsmedizinische Untersuchungen. In: Müller MJ (Hrsg): Ernährungsmedizinische Praxis: Methoden-Prävention-Behandlung. Springer, Berlin, Heidelberg, 1998, S. 29-202
- 67 Naganuma T, Sugimura K, Wada S, Yasumoto R, Sugimura T, Masuda C, Uchida J, Nakatani T: The prognostic role of brain natriuretic peptides in hemodialysis patients. *Am J Nephrol* 22 (2002) 437-444
- 68 National Kidney Foundation: K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification and stratification. *Am J Kidney Dis* 39 (2002) 1-266
- 69 Natori H, Tamaki S, Kira S: Ultrasonographic evaluation of ventilatory effect on inferior vena cava configuration. *Am Rev Respir Dis* 120 (1979) 421-427
- 70 Niebauer J, Volk HD, Kemp M, Dominguez M, Schumann RR, Rauchhaus M, Poole-Wilson PA, Coats AJS, Anker SD: Endotoxin and immune activation in chronic heart failure: a prospective cohort study. *Lancet* 353 (1999) 1838-1842
- 71 NIH (1996): NIH Consensus statement. Bioelectrical impedance analysis in body composition measurement. National Institutes of Health Technology Assessment Conference Statement. *Nutrition* 12 (1994) 749-762
- 72 Oe B de Fijter WM, de Fijter CWH, Straver B, Oe PL Stevens P, de Vries PMJM: Detection of hydration status by total body bioelectrical impedance analysis (BIA) in patients on hemodialysis. *Int J Art Org* 20 (1997) 371-374

- 73 Pepys M, Hirschfield GM: C-reactive protein: a critical update. *J Clin Invest* 111 (2003) 1805-1812
- 74 Perez-Garcia R, Lopez-Gomez JM, Jofre R, Junco E, Valderrabano F: Haemodialysis dose, extracellular volume control and arterial hypertension. *Nephrol Dial Transplant* 16 [Suppl 1] (2001) 98-101
- 75 Pietrobelli A, Formica C, Wang Z, Heymsfield SB: Dual-energy x-ray absorptiometry body composition model: review of physical concepts. *Am J Physiol Endocrinol Metab* 271 (1996) E941-E951
- 76 Pirlich M, Luhmann N, Schütz T, Plauth M, Lochs H: Mangelernährung bei Klinikpatienten: Diagnostik und klinische Bedeutung. *Akt Ernähr Med* 24 (1999) 260-266
- 77 Pollock CA, Ibels LS, Zhu FY, Warnant M, Caterson RJ, Waugh DA, Mahony JF: Protein intake in renal disease. *J Am Soc Nephrol* 8 (1997) 777-783
- 78 Rahman M, Dixit A, Donley V, Gupta S, Hanslik T, Lacson E, Ogundipe A, Weigel K, Smith MC: Factors associated with inadequate blood pressure control in hypertensive hemodialysis patients. *Am J Kidney Dis* 33 (1999) 498-506
- 79 Rauchhaus M, Doehner W, Francis DP, Davos C, Kemp M, Liebenthal C, Niebauer J, Volk HD, Coats AJ, Anker SD: Plasma cytokine parameters and mortality in patients with chronic heart failure. *Circulation* 102 (2000) 3060-3067
- 80 Reibe A: Die Dopplersonographie der Vena femoralis – ein Beitrag zur Bestimmung des Optimalgewichtes bei Dialysepatienten. Diss, Halle, 1997
- 81 Reibe A, Langer T, Rösch R, Osten B: Die Doppler-Sonographie der Vena femoralis. *Nieren- und Hochdruckkrankheiten* 27 (1998) 368-372
- 82 Ridker PM: Role of inflammatory biomarkers in prediction of coronary heart disease. *Lancet* 358 (2001) 946-948

- 83 Romero-Corral A, Montori VM, Somers VK, Korinek J, Thomas RJ, Allison TG, Mookadam F, Lopez-Jimenez F: Association of bodyweight with total mortality and with cardiovascular events in coronary artery disease: a systematic review of cohort studies. *Lancet* 368 (2006) 666-678
- 84 Rosanelli K: Present diagnostic value of IgM determination in newborns. In: Betke K, Riegel K, Belohradsky BH (Eds): *Diagnostics in Peritoneal Infections*. Die medizinische Verlagsgesellschaft, Marburg, 1984, S. 174-178
- 85 Santoro A, Mancini E, Paolini F, Zuccelli P: Blood volume monitoring and control. *Nephrol Dial Transplant* 11 [Suppl 2] (1996) 42-47
- 86 Sasaki T; Kubo T, Miyamoto T, Komamura K, Honda K, Miyatake K: Clinical significance of measuring inferior vena cava dimension in patients with acute exacerbation of chronic heart failure. *J Cardiol* 37 (2001) 309-313
- 87 Schoenfeld PY, Henry RR, Laird NM, Roxe DM: Assessment of nutritional status of the national cooperative dialysis study population. *Kidney Int* 23 (1983) 80-88
- 88 Schönweiß G: *Dialysefibel*. 2. Aufl. Abakiss, Bad Kissingen, 1996, S. 36-37
- 89 Schönweiß G: *Dialysefibel*. 2. Aufl. Abakiss, Bad Kissingen, 1996, S. 329-336
- 90 Stenvinkel P: The role of inflammation in the anaemia of end-stage renal disease. *Nephrol Dial Transplant* 16 (2001) 36-40
- 91 Stenvinkel P, Heimbürger O, Lindholm B, Kaysen GA, Bergström, J: Are there two types of malnutrition in chronic renal failure? *Nephrol Dial Transplant* 15 (2000) 953-960
- 92 Stenvinkel P, Heimbürger O, Paultre F, Diczfalusy U, Wang T, Berglund L, Jogestrand T: Strong association between malnutrition, inflammation and atherosclerosis in chronic renal failure. *Kidney Int* 55 (1999) 1899-1911

- 93 Stroh S: Methoden zur Erfassung der Körperzusammensetzung. Ernährungs-
umschau 42 (1995) 88-94
- 94 Thomas L: β_2 -Mikroglobulin. In: Thomas L (Hrsg): Clinical Laboratory
Diagnostics. TH-Books Verlagsgesellschaft, Frankfurt, 1998, S. 685-688
- 95 Thomas L: Immunglobuline. In: Thomas L (Hrsg): Labor und Diagnose. TH-
Books Verlagsgesellschaft, Frankfurt, 1998, S. 684-695
- 96 Thomas L: Labor und Diagnose. 5. Aufl. TH-Books Verlagsgesellschaft,
Frankfurt, 1998, S. 116-120
- 97 Venugopal SK, Devaraj S, Jialal I: Effect of C-reactive protein on vascular cells:
evidence for a pro-inflammatory, proatherogenic role. Curr Opin Nephrol
Hypertens 14 (2005) 33-37
- 98 Verma S, Li SH, Badiwala MV, Weisel RD, Fedak PW, Li RK, Dhillon B,
Mickle DA: Endothelin antagonism and interleukin-6 inhibition attenuate the
proatherogenic effects of C-reactive protein. Circulation 105 (2002) 1890-1896
- 99 Vorderwinkler KP, Artner-Dworzak E, Jakob G, Mair J, Diensti F, Pichler M,
Puschendorf B: Release of cyclic guanosine monophosphate evaluated as a
diagnostic tool in cardiac disease. Clin Chem 37 (1991) 186-190
- 100 Webster ADB: Laboratory investigation of primary deficiency of the lymphoid
system. Clinics Immunology and Allergy 5 (1985) 447
- 101 Wei CM, Heublein DM, Perrella MA, Lerman A, Rodeheffer RJ, McGregor CG,
Edwards WD, Schaff HV, Burnett Jr JC: Natriuretic peptide system in human
heart failure. Circulation 88 (1993) 1004-1009
- 102 Whicher J, Spence C: When is serum albumin worth measuring? Ann Clin
Biochem 24 (1987) 572-580
- 103 WHO: Obesity. Preventing and managing the global epidemic. Report of a
WHO Consultation. World Health Organ Tech Rep Ser 894 (2000) 1-253

- 104 Wolbink GJ, Brouwer MC, Buysmann S, ten Berge IJ, Hack CE: CRP-mediated activation of complement in vivo: assessment by measuring circulating complement-C-reactive protein complexes. *J Immunol* 157 (1996) 473-479
- 105 Wolfram G, Sitter T, Gottsman M, Gerzer R, Schiffel H: Assessment of dry weight in hemodialysis patients by the volume markers ANP and cGMP. *Nephrol Dial Transplant* 11 [Suppl 2] (1986) 28-30
- 106 Yamada T: Serum Amyloid A (SAA): a concise review of biology, assay methods and clinical usefulness. *Clin Chem Lab Med* 37 (1999) 381-388
- 107 Yoshimura M, Yasue H, Okumura K, Ogawa H, Jougasaki M, Mukoyama M, Nakao K, Imura H: Different secretion patterns of atrial natriuretic peptide and brain natriuretic peptide in patients with congestive heart failure. *Circulation* 87 (1993) 464-469
- 108 Zaluska WT, Schneditz D, Kaufman AM, Morris AT, Levin NW: Relative underestimation of fluid removal during hemodialysis hypotension measured by whole body bioimpedance. *ASAIO J* 44 (1998) 823-827
- 109 Zhu F, Schneditz D, Wang E, Martin K, Morris AT, Levin NW: Validation of changes in extracellular volume measured during hemodialysis using a segmental bioimpedance technique. *ASAIO J* 44 (1998) 541-545
- 110 Zimmermann J, Schramm L, Metzger T, Wanner C: Malnutrition, Inflammation und Arteriosklerose bei Niereninsuffizienz. *Nieren- und Hochdruckkrankheiten* 32 (2003) 401-407